

4-1-2000

The Age of the Woolly Rhino from Dream Cave, Derbyshire, UK

Donald A. McFarlane

Claremont McKenna College; Pitzer College; Scripps College

Joyce Lundberg

Carleton University

Derek C. Ford

McMaster University

Recommended Citation

McFarlane, D.A., J. Lundberg, and D.C. Ford. "The Age of the Woolly Rhino from Dream Cave, Derbyshire, UK." *Cave and Karst Science* 27.1 (2000): 25-28.

This Article is brought to you for free and open access by the W.M. Keck Science Department at Scholarship @ Claremont. It has been accepted for inclusion in WM Keck Science Faculty Papers by an authorized administrator of Scholarship @ Claremont. For more information, please contact scholarship@cuc.claremont.edu.

The Age of the Woolly Rhino from Dream Cave, Derbyshire, UK

Donald A McFARLANE¹, Joyce LUNDBERG² and Derek C FORD³

¹ W M Keck Science Center, The Claremont Colleges, 925 North Mills Avenue, Claremont, CA 91711, USA

² Department of Geography and Environmental Studies, Carleton University, Carleton, Ontario, K1S 5B6, Canada

³ Departments of Geography and Geology, McMaster University, Hamilton, Ontario L8S 4K1, Canada



Abstract: The Dream Cave woolly rhinoceros, *Coelodonta antiquitatis*, is a 'classic' specimen of a 'cold-stage' fossil fauna from central England. The find was illustrated and described by Dean William Buckland in his seminal tome *Reliquiae Diluvianae* (1823) during the first half of the 19th century, and made a significant contribution to the development of Buckland's views on the origin of extinct and extirpated fossil vertebrates. Here we present the first, albeit indirect, radiometric dates on the specimen, and argue that the animal fell into the cave just before 37,000 years before present, during the middle of Marine Isotope Stage 3 Interstadial (39-41 ka).

(Received 25 April 2000; Accepted 22 May 2000)

BACKGROUND AND HISTORICAL INTRODUCTION

Dream Cave first came to attention in December 1822, as a result of lead mining activity near the hamlet of Callow, near Wirksworth, Derbyshire. Miners sinking an exploratory shaft just north of Sprink Wood (NGR 275 530) breached a natural cave, completely (or almost completely) filled with clastic sediments. Whilst excavating this material they encountered the bones of 'ante Diluvian' animals. William Buckland, first Professor of Geology at Oxford University, learned of the discovery through the property owner, Phillip Gell, and a mutual friend, the Reverend D Stacy. Buckland made haste to the site "... for the purpose of examining all [the discovery's] circumstances..." (Buckland, 1823, p.61). The principal discovery was of the skeletal remains of an adult woolly rhinoceros, *Coelodonta antiquitatis*, (Fig.1), together with fragmentary remains of horse, bear, and deer. Continued excavation of the fill exposed a natural fissure entrance, which forms both the original and modern entrance to the cave. Buckland figured the cave in a copperplate illustration (Fig.2), drawn by T Webster from Buckland's original notes.

Despite the prominence afforded to Dream Cave in *Reliquiae Diluvianae*, the site has not been re-examined critically in more recent times. The presence of *Coelodonta* is characteristic of a cold-stage fauna, so the site has been generally assumed to be later Devensian in age (e.g. Sutcliffe, 1983). All radiometrically-dated *Coelodonta* specimens from Britain fall in the range 35,000 – 22,350 ¹⁴C years BP, but records are scarce (Table 1). Thus, a possibility that *Coelodonta* persisted through the Devensian maximum at ~ 18,000 ¹⁴C years BP cannot be discounted. It might even have extended into the Younger Dryas (~ 11,000-10,000 ¹⁴C years BP; chronozones from Mangerud *et al.*, 1974), which supported a fauna of reindeer (*Rangifer tarandus*) and steppe pika (*Ochotona pusilla*). The latest European continental record of *Coelodonta* is c.12,500 ¹⁴C years BP (skeletal remains and late Magdalenian engravings; Bosinski, 1978; 1981). If the species was eliminated locally during the Devensian maximum it may have failed to recolonise Britain as the ice front retreated.




Figure 1. Buckland's Dream Cave *Coelodonta* specimen, courtesy of the Oxford University Museum.




Figure 2. Copperplate illustration of Dream cave in 1822 (Buckland, 1823).

DISCUSSION OF THE CURRENT WORK

Examination of the *Coelodonta* specimen in the collections of the University Museum, Oxford University, uncovered the presence of a hitherto un-noted block of flowstone. The block carried the notation, ten in Indian Ink in Buckland's handwriting. "from out the Dream Cave". As indicated by the moulds of elastic fragments and the ochre-coloured clay adhering to the lower surface, this flowstone overlay the soft sediment that preserved the bones of the woolly rhinoceros. Unfortunately none of this flowstone or equivalent material remains at the site today. The specimen consists of a layer of dense, laminated flowstone, several centimetres thick, whose underside preserves traces of a yellowish, argillaceous sediment and moulds of rounded clasts 1

to 5mm in diameter. A sketch of the sub-sample taken from Buckland's block is shown in Fig. 3.

Uranium series disequilibrium dating by thermal ionisation mass spectrometry was conducted at McMaster University. Two samples of ~1g each were taken, and laboratory preparation and dating followed standard procedures (see for example, Lauritzen and Lundberg, 1999). Isotope measurement was carried out using a VG354 thermal ionization mass spectrometer. Results are presented in Table 2. The low value for the activity ratios of $^{230}\text{Th}/^{232}\text{Th}$ suggests the presence of some detrital or non-radiogenic ^{230}Th . Correction for this would reduce the ages a little. However, the very low absolute concentrations of thorium (below 0.1 ppm) and the relatively high value of $^{234}\text{U}/^{232}\text{Th}$

| | |
|---------------------------------|-------------------------------|
| 22,350 620 | Ogof Yr Ychen, Caldey Island. |
| 22,500 700 | Pin Hole Cave, England. |
| 24,372 153 (weighted mean of 5) | Earls Barton, Northampton. |
| 27,550 + 1370 / -1680 | Bishopbriggs, Scotland. |
| 28,160 435 | Kent's Cavern, Devon. |
| 29,300 480 | Robin Hood's Cave, England. |
| 29,450 350 | Leadenhall Street, London. |
| 30,729 405 (weighted mean of 2) | Ash Tree Cave, England. |
| 32,180 580 | Tornewton Cave, Somerset. |
| 33,200 650 | Pontnewydd Caves, Wales. |
| 34,101 779 (weighted mean of 3) | Conningbrook, Kent. |
| 34,559 573 (weighted mean of 2) | Kent's Cavern, Devon. |
| 40,900 1800 | Ash Tree Cave, England. |
| 42,700 2200 | Pin Hole Cave, England. |
| 42,900 2400 | Robin Hood's Cave, England. |

Table 1. Radiometric dates on *Coelodonta antiquitatis* (Gowlett, et al., 1987; Hedges et al., 1989; 1994; 1996)




Figure 3. Diagrammatic sketch of the flowstone from Buckland's Dream Cave collection.

suggest that the detrital component is insignificant, so that use of the corrected value is probably not justified.

Sample B was cut from the basal layer, separated from the next layer, which provided sample LM, by an obvious erosional hiatus, marked by unconformable growth layers. A date of 36,451 -1262 +1277 ka on the basal sample of calcite just above the sediment fill indicates that calcite deposition began towards the end of isotope stage 3. The hiatus probably represents the Late Glacial Maximum, isotope stage 2, when conditions would have been cold and dry. A date of 19,600 -690 +695 just above the hiatus level suggests that calcite deposition resumed immediately after local deglaciation. Higher layers of the sample have not been dated; the topmost few millimetres show a distinct change in colour and several clay-lined, but not erosional, hiatuses, one of which is the top layer. Dissolutional cavities shown in the basal layer in Fig. 3 developed upwards from the base, but the hiatus level limited their penetration. This dissolutional etching probably occurred as water penetrated the sediment-calcite interface after calcite deposition had ceased.

Because its lower surface demonstrates clearly that the flowstone was deposited on top of the cave sediment, the rhino must have fallen into the cave before calcite deposition began, i.e. before 36 ka. *C. antiquitatis* is known from remains dated (stratigraphically) as 'early Devensian' (c.70 - 54 ka) in the Crayforth brickearths of Kent (West, *et al.*, 1964), and from Tornewton Cave, Somerset (Sutcliffe and Zeuner, 1962). So, the species undoubtedly entered Britain following the Eemian Interglacial (marine isotope stage 5). However, the steppe-tundra climate of central England during marine isotope stage 4, 'early Devensian', time was interrupted between about 43 - 41 ka by a warm interstadial, evidenced by a fossil beetle fauna (Huizjer and Isarin, 1996). It seems likely that, had the Dream Cave *Coelodonta* deposit been in place, interstadial conditions would have initiated speleothem growth and capped the deposit. The flowstone shows no evidence of deposition at this time. Therefore, it is most parsimonious to conclude

Table 2. U-series data. All errors quoted are 2 σ . All ratios are activity ratios.

| Sample Lab code | mm from base | Age (years before 1997) | U conc (ppm) | Th conc (ppm) | $^{230}\text{Th}/^{234}\text{U}$ | $^{234}\text{U}/^{238}\text{U}$ | $^{230}\text{Th}/^{232}\text{Th}$ | Initial $^{234}\text{U}/^{238}\text{U}$ | $^{234}\text{U}/^{232}\text{Th}$ |
|-----------------|--------------|-------------------------|------------------|----------------|----------------------------------|---------------------------------|-----------------------------------|---|----------------------------------|
| 94 BUCK LM2 | 2-5 | 19,600 - 690 +695* | 1.272 ±0.0030 | 0.0411 ±0.0002 | 0.165 ±0.005329 | 1.084 ±0.003407 | 17.67 ±0.566 | 1.08924 ±0.00018 | 107.14 ±4.84 |
| 94 BUCK B2 | 5-8 | 36,451 -1262 +1277* | 1.539 ±0.0034 | 0.0892 ±0.0005 | 0.287 ±0.008459 | 1.196 ±0.003883 | 18.92 ±0.551 | 1.21778 ±0.00078 | 65.88 ±0.11 |

*Correction for detrital thorium using an initial $^{230}/^{232}$ activity ratio of 1.7 gives an age of 17,895 -680 +684 and 33,726 -1232 +1246 respectively.




Figure 4. Marine isotope chronology of the Devensian and the Dream Cave deposits.

that the rhino and the sediment deposit were emplaced during the sharp, cold stadial (Heinrich 4 event), which prevailed from about 41-39 ka (Fig.4). Sediment infilling must have been rapid, because the rhino was preserved intact (Appendix 1; Fig.1), which could not have occurred if the carcass had remained exposed for many seasons under such harsh conditions. An amelioration of climate at 39-36 ka followed the Heinrich H4 stadial, and the timing of this mild interstadial correlates well with the deduced date of initiation of speleothem growth over the sediment infill at Dream Cave.

ACKNOWLEDGEMENTS

Richard Gledhill and Anna Hubbard provided assistance with the fieldwork. Mr Phillip Powell of the University Museum, Oxford, kindly provided access to the specimen. Uranium series dating was performed in the McMaster University Speleothem Dating Laboratory, supported by the Natural Sciences and Engineering Research Council of Canada grant to D C Ford.

REFERENCES

- Bosinski, G (Editor), 1978. Geowissenschaftliche Untersuchungen in Gönnersdorf. [Wiesbaden.]
- Bosinski G, 1981. Gönnersdorf; Eiszeitjäger am Mittelrhein. 258pp. [Koblenz: Rhenania. Steiner.]
- Buckland, W. 1823. Reliquiae Diluvianae: or, observations on the organic remains contained in caves, fissures, and diluvial gravel, and on other geological phenomena, attesting to the action of a universal deluge. viii + 303 pp. [London: Murray.]
- Gowlett, J A, Hedges, R E M, Law, I A and Perry, C, 1987. Radiocarbon dates from the Oxford AMS system: Archaeometry Datelist 5. *Archaeometry*, Vol.29(1),125-155

- Hedges, R E M, Housley, R A, Law, I A and Bronk, C R, 1989. Radiocarbon dates from the Oxford AMS system: *Archaeometry Datalist 9*. *Archaeometry*, Vol.31(2), 207-234.
- Hedges, R E M, Housley, R A, Ramsey, C R and von Klinken, G J, 1994. Radiocarbon dates from the Oxford AMS system: *Archaeometry Datalist 18*. *Archaeometry*, Vol.36(2), 337-344.
- Hedges, R E M, Pettit, P B, Ramsey, C R and von Klinken, G J, 1996. Radiocarbon dates from the Oxford AMS system: *Archaeometry Datalist 22*. *Archaeometry*, Vol.38(2), 391-415.
- Huijzer, A S and Isarin R F B. 1996. Reconstruction of past climates using multi-proxy evidence: example of the Weichselian pleniglacial in NW and central Europe. *Quaternary Science Reviews*, Vol.16, 513-533.
- Lauritzen, S-E and Lundberg, J. 1999. Calibration of the speleothem delta function: an absolute temperature record for the Holocene in northern Norway. *The Holocene*, Vol.9, 6, 659-669.
- Mangerud, J, Andersen, S T, Berglund, B E and Donner, J J, 1974. Quaternary stratigraphy of Norden: a proposal for terminology and classification. *Boreas*, Vol.3, 109-127.
- Stuart, A J, 1983. Pleistocene bone caves in Britain and Ireland: a short review. *Studies in Speleology*, Vol.4(9), 9-36.
- Sutcliffe, A J and Zeuner, F E. 1962. Excavations in the Torbryan Caves, Devonshire. 1. Tornewton Cave. *Proceedings of the Devon Archaeological and Exploration Society*, Vol 5, 127-145.
- West, R G, Lambert, C A and Sparks, B W, 1964. Interglacial deposits at Ilford, Essex. *Philosophical Transactions of the Royal Society*, B247, 185-212.

APPENDIX I

| | |
|---------------|---|
| Skull and Jaw | |
| | Nasal region with septum Partial maxilla 5 detached upper molariform teeth Left mandible Right mandible 1 detached lower molariform tooth |
| Trunk | Atlas 3 rd and 4 th cervical vertebrae 2 dorsal vertebrae 3 lumbar vertebrae Sacrum 1 rib fragment Partial left scapula |
| Limbs | Left and right humerus, ulna, radius Left and right scaphoid, lunar, cuneiform (left only), magnum(left only) and unciform carpals Left and right II, III, and IV forelimb metacarpals Left and right portions of pelvis Left and right femora and tibia Left and right patella Left and right astragalus, calcaneum, and cuboid tarsals Right navicular tarsal II, II, IV hind metatarsals |