

1-1-2012

# On the Closure of the Complex Symmetric Operators: Compact Operators and Weighted Shifts

Stephan Ramon Garcia  
*Pomona College*

Daniel E. Poore '11  
*Pomona College*

---

## Recommended Citation

Garcia, Stephan Ramon and Poore, Daniel E. '11, "On the Closure of the Complex Symmetric Operators: Compact Operators and Weighted Shifts" (2012). *Pomona Faculty Publications and Research*. 244.  
[http://scholarship.claremont.edu/pomona\\_fac\\_pub/244](http://scholarship.claremont.edu/pomona_fac_pub/244)

This Article - preprint is brought to you for free and open access by the Pomona Faculty Scholarship at Scholarship @ Claremont. It has been accepted for inclusion in Pomona Faculty Publications and Research by an authorized administrator of Scholarship @ Claremont. For more information, please contact [scholarship@cuc.claremont.edu](mailto:scholarship@cuc.claremont.edu).

**ON THE CLOSURE OF THE COMPLEX SYMMETRIC  
OPERATORS: COMPACT OPERATORS  
AND WEIGHTED SHIFTS**

STEPHAN RAMON GARCIA AND DANIEL E. POORE

ABSTRACT. We study the closure  $\overline{CSO}$  of the set  $CSO$  of all complex symmetric operators on a separable, infinite-dimensional, complex Hilbert space. Among other things, we prove that every compact operator in  $\overline{CSO}$  is complex symmetric. Using a construction of Kakutani as motivation, we also describe many properties of weighted shifts in  $\overline{CSO} \setminus CSO$ . In particular, we show that weighted shifts which demonstrate a type of approximate self-similarity belong to  $\overline{CSO} \setminus CSO$ . As a byproduct of our treatment of weighted shifts, we explain several ways in which our result on compact operators is optimal.

1. INTRODUCTION

Throughout the following, we let  $\mathcal{H}$  denote a separable, infinite-dimensional complex Hilbert space. Recall that  $T \in \mathcal{B}(\mathcal{H})$  is a *complex symmetric operator* if there exists a *conjugation*  $C$  (i.e., a conjugate-linear, isometric involution on  $\mathcal{H}$ ) such that  $T = CT^*C$ . We remark that the term *complex symmetric* stems from the fact that  $T$  is a complex symmetric operator if and only if  $T$  is unitarily equivalent to a symmetric (i.e., self-transpose) matrix with complex entries, regarded as an operator acting on a  $\ell^2$ -space of the appropriate dimension [15, Sect. 2.4]. The general study of complex symmetric operators was undertaken by the first author, M. Putinar, and W.R. Wogen in [16, 18, 20, 21, 23, 24], although much of the theory has classical roots in the matrix-oriented work of N. Jacobson [28], T. Takagi [38], C.L. Siegel [37], and I. Schur [34]. A number of other authors have recently made significant contributions to the study of complex symmetric operators [7, 25, 29–31, 39, 40, 42, 44], which has proven particularly relevant to the study of truncated Toeplitz operators [6, 8, 9, 22, 35, 36], a rapidly growing branch of function-theoretic operator theory stemming from the seminal work of D. Sarason [33].

In the following, we let  $CSO$  denote the set of all complex symmetric operators on  $\mathcal{H}$ . We remark that the set  $CSO$  is neither closed under addition nor under multiplication, although it is closed under the adjoint operation and the Aluthge transform [17, Thm. 1], a remarkable nonlinear mapping on  $\mathcal{B}(\mathcal{H})$  which has been much studied in recent years [1–5, 13, 14, 27, 41]. Lately there has been some interest in the study of  $CSO$  itself as a subset of  $\mathcal{B}(\mathcal{H})$  [19, 23, 44]. Along these lines we begin by examining the closure of  $CSO$  in several of the most common topologies

---

2010 *Mathematics Subject Classification.* 47A05, 47B35, 47B99.

*Key words and phrases.* Complex symmetric operator, unitary orbit, Hilbert space, compact operator, strong-\* topology, strong operator topology, weak operator topology, Kakutani shift, self-similarity, palindrome, shift operator, weight sequence, unilateral shift, irreducible.

Partially supported by National Science Foundation Grant DMS-1001614.

on  $\mathcal{B}(\mathcal{H})$  (Section 2). In particular, we prove that the closure of  $CSO$  in the strong- $*$  topology is all of  $\mathcal{B}(\mathcal{H})$ . Among other things, this immediately implies that the strong-operator and weak-operator closures of  $CSO$  are both  $\mathcal{B}(\mathcal{H})$ . This contrasts sharply with the situation for the norm topology, which we discuss in significant detail below.

Let us denote by  $\overline{CSO}$  the closure of  $CSO$  with respect to the operator norm on  $\mathcal{B}(\mathcal{H})$ . We first remark that  $\overline{CSO}$  is a proper subset of  $\mathcal{B}(\mathcal{H})$ . Although this will be clear from what follows, we should mention that a simple example of an operator in  $\mathcal{B}(\mathcal{H}) \setminus \overline{CSO}$  can be constructed by taking the direct sum of the matrix [24, Ex. 1] with 0 and then applying [23, Lem. 1].

The so-called *norm closure problem* for complex symmetric operators asked whether or not  $\overline{CSO} = CSO$  [23, p. 1260]. Although it appeared in print only in 2009, this question had been circulating around the community for some years prior. Recently, S. Zhu, C.G. Li, and Y.Q. Ji demonstrated that a particular weighted shift operator belongs to  $\overline{CSO} \setminus CSO$ , thereby settling the norm-closure problem in the negative [44]. Shortly thereafter, the authors of this note constructed a completely different counterexample using a certain infinite direct sum of multiples of the unilateral shift and its adjoint [19]. These examples indicate that the structure of  $\overline{CSO}$  is much richer than previously expected. In particular, a complete description of the set  $\overline{CSO}$  is now much desired.

**Compact operators.** Our first main result (Theorem 4) asserts that every compact operator in  $\overline{CSO}$  belongs to  $CSO$ . In some sense, this complements the results of [19] and [44], since none of the examples of operators in  $\overline{CSO} \setminus CSO$  described there are compact. Let us also remark that Theorem 4 furnishes a simple proof that  $\overline{CSO} \neq \mathcal{B}(\mathcal{H})$ . Indeed, if  $T$  is an irreducible weighted unilateral shift whose weights tend to zero, then  $T$  is compact by [11, Cor. 4.27.5]. However,  $\dim \ker T = 0 \neq 1 = \dim \ker T^*$  for such an operator whence  $T$  is not complex symmetric by [20, Prop. 1].

It turns out that our result about compact operators is sharp in the following sense. Our proof relies heavily upon the fact that the spectrum  $\sigma(|T|)$  of the *modulus*  $|T| = \sqrt{T^*T}$  of a compact operator consist of 0 along with a decreasing sequence of positive eigenvalues of finite multiplicity. As we will see, it is possible to construct operators in  $\overline{CSO} \setminus CSO$  such that  $\sigma(|T|) \setminus \{0, 1, \frac{1}{2}, \frac{1}{4}, \dots\}$  consists only of eigenvalues, each of multiplicity one (Example 12). On the other hand, the so-called *Kakutani shift* (discussed at length below) belongs to  $\overline{CSO} \setminus CSO$  and satisfies  $\sigma(|T|) = \{0\} \cup \{\frac{1}{2^n} : n = 0, 1, 2, \dots\}$ , each nonzero eigenvalue being of infinite multiplicity. In light of these examples, it is difficult to envision a stronger version of Theorem 4.

**Weighted shifts.** As the preceding comments suggest, the study of  $\overline{CSO}$  leads naturally to the consideration of weighted shifts. Consequently, a substantial portion of this article is dedicated to this topic. We say that  $T \in \mathcal{B}(\mathcal{H})$  is a *unilateral weighted shift* (or simply a *weighted shift*) if there is an orthonormal basis  $\{e_n\}_{n=1}^\infty$  of  $\mathcal{H}$  and a sequence of scalars  $\{\alpha_n\}_{n=1}^\infty$  (the *weight sequence*) such that  $Te_n = \alpha_n e_{n+1}$  for  $n \geq 1$ . Since the weighted shift having weight sequence  $\{\alpha_n\}_{n=1}^\infty$  is unitarily equivalent to the unilateral shift with weight sequence  $\{|\alpha_n|\}_{n=1}^\infty$  (see [11, Prop. 4.27.2] or [26, Prob. 89]), we henceforth assume that  $\alpha_n \geq 0$  for all  $n \geq 1$ . We maintain this convention and the preceding notation in what follows.



satisfies  $\|T - T'\| < \epsilon$  and is complex symmetric since it is unitarily equivalent to a direct sum of matrices of the form (1) having palindromic weights.

Let us make a few remarks about the preceding construction which will help motivate our results. First of all, observe that the weight sequence  $\{\alpha_n\}_{n=1}^\infty$  of the Kakutani shift (2) has a subsequence which converges to zero and another which tends to  $\sup\{\alpha_n\}_{n=1}^\infty$ . As we will see, this behavior is typical of all irreducible weighted shifts belonging to  $\overline{CSO}$  (Theorem 7). In particular, no irreducible weighted shift in  $\overline{CSO}$  is compact. This agrees with our earlier remarks about compact operators.

It turns out that every irreducible weighted shift which is *approximately Kakutani*, in a sense made precise in Section 5, belongs to  $\overline{CSO} \setminus CSO$  (Theorem 10). Among other things, this allows us to construct operators in  $\overline{CSO} \setminus CSO$  whose moduli have desired spectral properties.

## 2. CLOSURES IN THE WEAK, STRONG, AND STRONG-\* TOPOLOGIES

Although we are primarily interested in studying the closure  $\overline{CSO}$  of  $CSO$  with respect to the operator norm, let us first say a few words about the closure of  $CSO$  with respect to several other standard topologies on  $\mathcal{B}(\mathcal{H})$ . In particular, we consider the *weak operator topology* (WOT), *strong operator topology* (SOT), and *strong-\* topology* (SST). We say that

- (i)  $T_n \rightarrow T$  means that  $\lim_{n \rightarrow \infty} \|T - T_n\| = 0$ ,
- (ii)  $T_n \rightarrow T$  (WOT) means that  $\lim_{n \rightarrow \infty} \langle (T - T_n)x, y \rangle = 0$  for all  $x, y \in \mathcal{H}$ ,
- (iii)  $T_n \rightarrow T$  (SOT) means that  $\lim_{n \rightarrow \infty} \|(T - T_n)x\| = 0$  for all  $x \in \mathcal{H}$ ,
- (iv)  $T_n \rightarrow T$  (SST) means that  $T_n \rightarrow T$  (SOT) and  $T_n^* \rightarrow T^*$  (SOT).

Observe that the containments

$$\overline{CSO} \subseteq \overline{CSO}^{(\text{SST})} \subseteq \overline{CSO}^{(\text{SOT})} \subseteq \overline{CSO}^{\text{WOT}} \subseteq \mathcal{B}(\mathcal{H})$$

hold trivially. The superscripts in the preceding chain indicate the topology with respect to which the closure of  $CSO$  is taken, the absence of a superscript being reserved for the norm topology. Let us also remark that the preceding notions have obvious analogues for conjugate-linear operators, which we employ without further comment.

To proceed, we require the following well-known lemma [10, Prop. IX.1.3.d]:

**Lemma 2.** *Suppose that  $\{e_i\}_{i=1}^\infty$  is an orthonormal basis of  $\mathcal{H}$  and that  $T_n$  is a bounded sequence in  $\mathcal{B}(\mathcal{H})$ . If  $T_n e_i \rightarrow T e_i$  for all  $i$ , then  $T_n \rightarrow T$  (SOT).*

We are now ready to prove the main result of this section. Our approach is inspired by the arguments of [23, Ex. 2], which were in turn inspired by a matrix trick first observed in [12, p. 793].

**Theorem 3.**  $\overline{CSO}^{(\text{SST})} = \overline{CSO}^{(\text{SOT})} = \overline{CSO}^{\text{WOT}} = \mathcal{B}(\mathcal{H})$ .

*Proof.* Since the strong and weak operator topologies are both weaker than the strong-\* topology, it suffices to prove that  $\overline{CSO}^{(\text{SST})} = \mathcal{B}(\mathcal{H})$ . Let  $T \in \mathcal{B}(\mathcal{H})$ , fix an orthonormal basis  $\beta = \{e_1, e_2, \dots\}$  of  $\mathcal{H}$ , and let  $\mathcal{H}_n = \text{span}\{e_1, e_2, \dots, e_n\}$ . Define  $A_n \in \mathcal{B}(\mathcal{H}_n)$  by insisting that  $\langle A_n e_k, e_j \rangle = \langle T e_k, e_j \rangle$  for  $1 \leq j, k \leq n$ . In other words,  $A_n$  is simply the upper-left  $n \times n$  principal submatrix of the matrix representation of  $T$  with respect to the basis  $\beta$ . Let  $C_n$  be an arbitrary conjugation

on  $\mathcal{H}_n$  and observe that the operator  $T_n = A_n \oplus C_n A_n^* C_n \oplus 0$  on  $\mathcal{H}$  is complex symmetric by [12, p. 793]. Since  $n > i$  implies that

$$\|Te_i - T_n e_i\|^2 = \sum_{j=n+1}^{\infty} |\langle Te_i, e_j \rangle|^2, \quad (3)$$

it follows that  $T_n e_i \rightarrow Te_i$  for each fixed  $i$ . Since  $\|T_n\| = \|A_n\| \leq \|T\|$  by construction, it follows from Lemma 2 that  $T_n \rightarrow T$  (SOT). An estimate similar to (3) and another appeal to Lemma 2 confirm that  $T_n^* \rightarrow T^*$  (SOT) as well, whence  $T_n \rightarrow T$  (SST).  $\square$

Among other things, the preceding theorem implies that

$$\overline{CSO} \subsetneq \overline{CSO}^{(\text{SST})} = \overline{CSO}^{(\text{SOT})} = \overline{CSO}^{\text{WOT}} = \mathcal{B}(\mathcal{H}).$$

In particular, the closure of  $CSO$  is only of interest if the closure is taken respect to the norm topology on  $\mathcal{B}(\mathcal{H})$ .

### 3. COMPACT OPERATORS

Having seen that  $\overline{CSO}$  is a proper subset of  $\mathcal{B}(\mathcal{H})$ , one naturally wishes to know how various well-studied classes of operators intersect  $\overline{CSO}$ . It turns out that a complete answer can be given in the case of compact operators.

**Theorem 4.** *If  $T \in \overline{CSO}$  and  $T$  is compact, then  $T \in CSO$ .*

Before proving Theorem 4, we require two simple lemmas.

**Lemma 5.** *The set of all conjugations on  $\mathcal{H}$  is SOT closed.*

*Proof.* Let  $C_n$  be a sequence of conjugations on  $\mathcal{H}$  such that  $C_n \rightarrow C$  (SOT). For each  $x \in \mathcal{H}$  we have

$$\begin{aligned} \|C^2 x - x\| &= \|C^2 x - C_n^2 x\| \\ &\leq \|C^2 x - C_n C x\| + \|C_n C x - C_n^2 x\| \\ &= \|(C - C_n)C x\| + \|(C - C_n)x\|, \end{aligned}$$

which tends to zero by hypothesis whence  $C^2 = I$ . Next observe that

$$|\|Cx\| - \|x\|| = |\|Cx\| - \|C_n x\|| \leq \|Cx - C_n x\| \rightarrow 0,$$

from which it follows that  $C$  is isometric. Since  $C$  is obviously conjugate-linear, we conclude that  $C$  is a conjugation on  $\mathcal{H}$ .  $\square$

**Lemma 6.** *If  $T \in \overline{CSO}$ , then there exist conjugations  $C_n$  such that  $C_n T^* C_n \rightarrow T$ .*

*Proof.* If  $T_n$  is a sequence of operators such that  $T_n \rightarrow T$  and  $C_n$  is a sequence of conjugations such that  $T_n = C_n T_n^* C_n$ , then

$$\begin{aligned} \|T - C_n T^* C_n\| &\leq \|T - C_n T_n^* C_n\| + \|C_n T_n^* C_n - C_n T^* C_n\| \\ &= \|T - T_n\| + \|T_n^* - T^*\| \\ &= 2 \|T - T_n\| \end{aligned}$$

tends to zero.  $\square$

*Pf. of Theorem 4.* Suppose that  $T$  belongs to  $\overline{CSO}$ . Let  $\mathcal{K} = \overline{\text{ran } T + \text{ran } T^*}$  so that  $\mathcal{K}^\perp \subseteq \ker T \cap \ker T^*$  is a reducing subspace upon which  $T$  vanishes. In other words, with respect to the orthogonal decomposition  $\mathcal{H} = \mathcal{K} \oplus \mathcal{K}^\perp$  we have  $T = T|_{\mathcal{K}} \oplus 0$ . By [23, Lem. 1] we know that  $T \in \mathcal{B}(\mathcal{H})$  is complex symmetric if and only if  $T|_{\mathcal{K}} \in \mathcal{B}(\mathcal{K})$  is complex symmetric. Without loss of generality, we therefore assume that

$$\overline{\text{ran } T + \text{ran } T^*} = \mathcal{H}. \quad (4)$$

Let us briefly discuss our main approach. Since  $T$  belongs to  $\overline{CSO}$ , there exist conjugations  $C_n$  such that  $C_n T^* C_n \rightarrow T$  by Lemma 6. It suffices to prove that there exists a subsequence  $C_{n_j}$  of the  $C_n$  which converges pointwise on both  $\text{ran } T$  and  $\text{ran } T^*$ . Indeed, the uniform boundedness of the  $C_{n_j}$  and the assumption (4) will then ensure that  $C_{n_j}$  is SOT convergent on all of  $\mathcal{H}$ , whence there exists a conjugation  $C$  such that  $C_{n_j} \rightarrow C$  (SOT) by Lemma 5. The desired conclusion  $T = CT^*C$  will then follow by a simple limiting argument.

By [10, VIII.3.11], there exists a partial isometry  $U$  with  $\ker T = \ker U = \ker |T|$  such that  $T = U|T|$ . Here  $|T|$  denotes the compact selfadjoint operator  $\sqrt{T^*T}$ . Let  $\lambda_1 \geq \lambda_2 \geq \dots > 0$  denote the nonzero eigenvalues of  $|T|$ , repeated according to their multiplicity, and let  $x_1, x_2, \dots$  denote corresponding orthonormal eigenvectors. The vectors  $y_i = Ux_i$  for  $i = 1, 2, \dots$  are also orthonormal and we see that

$$\overline{\text{span}\{y_1, y_2, \dots\}} = \overline{\text{ran } T}, \quad \overline{\text{span}\{x_1, x_2, \dots\}} = \overline{\text{ran } T^*}. \quad (5)$$

By the remarks of the preceding paragraph, it suffices to find a subsequence  $C_{n_j}$  of  $C_n$  such that  $\lim_{j \rightarrow \infty} C_{n_j} x_m$  and  $\lim_{j \rightarrow \infty} C_{n_j} y_m$  exist for  $m = 1, 2, \dots$

By compactness, the eigenspaces for  $|T|$  corresponding to its nonzero eigenvalues are finite-dimensional and hence we may define a sequence  $\ell_k$  of indices such that  $\lambda_{\ell_k} > \lambda_{\ell_{k+1}}$  and  $\lambda_{\ell_k} = \lambda_i$  for  $\ell_k \leq i < \ell_{k+1}$ . Let  $P_k$  denote the orthogonal projection onto the spectral subspace

$$\ker(|T| - \lambda_{\ell_k} I) = \text{span}\{x_{\ell_k}, x_{\ell_k+1}, \dots, x_{\ell_{k+1}-1}\} \quad (6)$$

and observe that  $Q_k = UP_k U^*$  is the orthogonal projection onto

$$\text{span}\{y_{\ell_k}, y_{\ell_k+1}, \dots, y_{\ell_{k+1}-1}\}.$$

The most difficult step in the proof of Theorem 4 is the verification of the following claim:

**Claim.** For  $\ell_k \leq m < \ell_{k+1}$ , we have

$$\lim_{n \rightarrow \infty} \|P_k C_n y_m\| = 1, \quad (7)$$

$$\lim_{n \rightarrow \infty} \|Q_k C_n x_m\| = 1. \quad (8)$$

*Pf. of Claim.* We proceed by induction on  $k$ . Suppose for our inductive hypothesis that

$$\lim_{n \rightarrow \infty} \|P_j C_n y_i\| = \lim_{n \rightarrow \infty} \|Q_j C_n x_i\| = 1 \quad (9)$$

holds for all  $\ell_j \leq i < \ell_{j+1}$  whenever  $0 < j < k$ . Observe that this is trivially true if  $k = 1$  since no corresponding  $j$  exist.

Let  $\ell_k \leq m < \ell_{k+1}$  and for each fixed  $i < \ell_k$  let  $j$  be the unique index  $j < k$  such that  $\ell_j \leq i < \ell_{j+1}$ . By the inductive hypothesis (9), we see that

$$\lim_{n \rightarrow \infty} (I - P_j) C_n y_i = 0 \quad (10)$$

since  $C_n y_i$  is a unit vector. For  $i < \ell_k$  it follows from (10) that

$$\begin{aligned}
\lim_{n \rightarrow \infty} \langle C_n x_m, y_i \rangle &= \lim_{n \rightarrow \infty} \langle C_n y_i, x_m \rangle \\
&= \lim_{n \rightarrow \infty} \langle P_j C_n y_i, x_m \rangle + \lim_{n \rightarrow \infty} \langle (I - P_j) C_n y_i, x_m \rangle \\
&= \lim_{n \rightarrow \infty} \langle C_n y_i, 0 \rangle + \langle 0, x_m \rangle \\
&= 0
\end{aligned} \tag{11}$$

since  $P_j x_m = 0$  by definition.

Suppose that  $0 < \epsilon < 4\lambda_{\ell_k}^2$ . Since  $C_n T^* C_n \rightarrow T$  there exists some  $N_1$  so that

$$n \geq N_1 \quad \Rightarrow \quad \|C_n T^* C_n x_m - T x_m\| < \frac{\epsilon}{4\lambda_{\ell_k}}$$

holds whenever  $\ell_k \leq m < \ell_{k+1}$ . On the other hand, by (11) there exists some  $N_2$  such that

$$n \geq N_2 \quad \Rightarrow \quad \sum_{i=1}^{\ell_k-1} \lambda_i^2 |\langle C_n x_m, y_i \rangle|^2 < \frac{\epsilon}{2}. \tag{12}$$

Since  $\|T x_m\| = \lambda_{\ell_k}$ , for  $n \geq N = \max\{N_1, N_2\}$  it follows that

$$\begin{aligned}
\lambda_{\ell_k}^2 - \frac{\epsilon}{2} &< \lambda_{\ell_k}^2 - \frac{\epsilon}{2} + \frac{\epsilon^2}{16\lambda_{\ell_k}^2} \\
&< \left( \lambda_{\ell_k} - \frac{\epsilon}{4\lambda_{\ell_k}} \right)^2 \\
&\leq (\|T x_m\| - \|C_n T^* C_n x_m - T x_m\|)^2 \\
&\leq \|C_n T^* C_n x_m\|^2 \\
&= \|T^* C_n x_m\|^2 \\
&= \sum_{i=1}^{\infty} |\langle T^* C_n x_m, x_i \rangle|^2 && \text{by (5)} \\
&= \sum_{i=1}^{\infty} |\langle T | U^* C_n x_m, x_i \rangle|^2 \\
&= \sum_{i=1}^{\infty} \lambda_i^2 |\langle U^* C_n x_m, x_i \rangle|^2 \\
&= \sum_{i=1}^{\infty} \lambda_i^2 |\langle C_n x_m, y_i \rangle|^2 \\
&< \frac{\epsilon}{2} + \sum_{i=\ell_k}^{\infty} \lambda_i^2 |\langle C_n x_m, y_i \rangle|^2 && \text{by (12)} \\
&< \frac{\epsilon}{2} + \lambda_{\ell_k}^2 \underbrace{\sum_{i=\ell_k}^{\ell_{k+1}-1} |\langle C_n x_m, y_i \rangle|^2}_{d_{n,m}} + \lambda_{\ell_{k+1}}^2 \sum_{i=\ell_{k+1}}^{\infty} |\langle C_n x_m, y_i \rangle|^2 \\
&\leq \frac{\epsilon}{2} + \lambda_{\ell_k}^2 d_{n,m} + \lambda_{\ell_{k+1}}^2 (1 - d_{n,m}).
\end{aligned}$$



The final inequality follows since  $C_n x_m$  is a unit vector and  $y_{\ell_k}, y_{\ell_k+1}, \dots$  is an orthonormal set. Rearranging things somewhat we see that

$$n \geq N \quad \Rightarrow \quad 0 \leq \underbrace{(\lambda_{\ell_k}^2 - \lambda_{\ell_k+1}^2)}_{>0} (1 - d_{n,m}) < \epsilon, \quad (13)$$

whence

$$\lim_{n \rightarrow \infty} d_{n,m} = 1 \quad (14)$$

whenever  $\ell_k \leq m < \ell_{k+1}$ . Since

$$d_{n,m} = \sum_{i=\ell_k}^{\ell_{k+1}-1} |\langle C_n x_m, Q_k y_i \rangle|^2 = \|Q_k C_n x_m\|^2,$$

we obtain the second condition (8) of the claim. To conclude the induction, we need to establish the first condition (7).

Summing (14) over  $\ell_k \leq i < \ell_{k+1}$  we obtain

$$\begin{aligned} \ell_{k+1} - \ell_k &= \lim_{n \rightarrow \infty} \sum_{i=\ell_k}^{\ell_{k+1}-1} d_{n,i} \\ &= \lim_{n \rightarrow \infty} \sum_{i=\ell_k}^{\ell_{k+1}-1} \sum_{m=\ell_k}^{\ell_{k+1}-1} |\langle C_n x_i, y_m \rangle|^2 \\ &= \sum_{m=\ell_k}^{\ell_{k+1}-1} \left( \lim_{n \rightarrow \infty} \sum_{i=\ell_k}^{\ell_{k+1}-1} |\langle C_n y_m, x_i \rangle|^2 \right) \\ &\leq \sum_{m=\ell_k}^{\ell_{k+1}-1} 1 \\ &= \ell_{k+1} - \ell_k \end{aligned}$$

by applying Bessel's inequality to the unit vector  $C_n y_m$  and using the fact that set  $x_{\ell_k}, x_{\ell_k+1}, \dots, x_{\ell_{k+1}-1}$  is orthonormal. In particular, the preceding tells us that

$$\lim_{n \rightarrow \infty} \|P_k C_n y_m\|^2 = \lim_{n \rightarrow \infty} \sum_{i=\ell_k}^{\ell_{k+1}-1} |\langle C_n y_m, P_k x_i \rangle|^2 = 1,$$

which is the first condition (7) of the claim. This concludes the proof of the claim.  $\square$

We now wish to prove that there exists a subsequence  $C_{n_j}$  of  $C_n$  such that  $C_{n_j} x_m$  and  $C_{n_j} y_m$  converge for each  $m = 1, 2, 3, \dots$ . For each fixed  $m$  there exists a unique  $k$  such that  $\ell_k \leq m < \ell_{k+1}$ . Since the sets  $\{P_k C_n y_m : n = 1, 2, \dots\}$  and  $\{Q_k C_n x_m : n = 1, 2, \dots\}$  are bounded subsets of the finite-dimensional spaces  $\text{ran } P_k$  and  $\text{ran } Q_k$ , respectively, it follows from a standard subsequence refinement argument that there exists vectors  $y'_m \in \text{ran } P_k$  and  $x'_m \in \text{ran } Q_k$  and a subsequence  $C_{n_j}$  of  $C_n$  such that

$$\lim_{j \rightarrow \infty} P_k C_{n_j} y_m = y'_m, \quad (15)$$

$$\lim_{j \rightarrow \infty} Q_k C_{n_j} x_m = x'_m. \quad (16)$$

Putting this all together we find that

$$\begin{aligned}
\lim_{j \rightarrow \infty} \|C_{n_j} y_m - y'_m\| &\leq \lim_{j \rightarrow \infty} \|(I - P_k)C_{n_j} y_m\| + \lim_{j \rightarrow \infty} \|P_k C_{n_j} y_m - y'_m\| \\
&= \lim_{j \rightarrow \infty} \sqrt{\|C_{n_j} y_m\|^2 - \|P_k C_{n_j} y_m\|^2} + 0 \\
&= \sqrt{1 - \lim_{j \rightarrow \infty} \|P_k C_{n_j} y_m\|^2} \\
&= 0
\end{aligned}$$

by (15) and (7), respectively. Thus  $\lim_{j \rightarrow \infty} C_{n_j} y_m = y'_m$ , as desired. An analogous argument confirms that  $\lim_{j \rightarrow \infty} C_{n_j} x_m = x'_m$  as well. This concludes the proof of Theorem 4.  $\square$

#### 4. WEIGHTED SHIFTS

We turn our attention now toward weighted shifts. It turns out that many features of the Kakutani shift (2) are typical of irreducible weighted shifts which belong to  $\overline{CSO}$ . For instance, consider the following theorem.

**Theorem 7.** *If  $T \in \overline{CSO}$  is an irreducible weighted shift with weights  $\{\alpha_n\}_{n=1}^\infty$ , then*

- (i) *there exists a subsequence of  $\{\alpha_n\}_{n=1}^\infty$  which tends to zero,*
- (ii) *there exists a subsequence of  $\{\alpha_n\}_{n=1}^\infty$  which tends to  $\alpha_+ = \sup\{\alpha_n\}_{n=1}^\infty$ .*

*In particular, 0 and  $\alpha_+$  belong to the essential spectrum  $\sigma_e(|T|)$  of  $|T| = \sqrt{T^*T}$ .*

Recalling that a weighted shift  $T$  with weights  $\{\alpha_n\}_{n=1}^\infty$  is compact if and only if  $\alpha_n \rightarrow 0$  [11, Cor. 4.27.5], we see that Theorem 7 asserts that there are no compact irreducible weighted shifts in  $\overline{CSO}$ , in agreement with Theorem 4.

The remainder of this section is devoted to developing the tools required to prove Theorem 7. In particular, we prove statements (i) and (ii) separately since they call for completely different methods. The final statement of Theorem 7, however, can easily be justified since statements (i) and (ii) imply that neither 0 nor  $\alpha_+$  is an isolated eigenvalue of  $|T|$  of finite multiplicity [10, Prop. 4.6].

Before proving the first portion of Theorem 7, we need to introduce a few useful facts about the spectral theory of operators in  $CSO$  and its closure. Recall that a complex number  $\lambda$  belongs to the *approximate point spectrum*  $\sigma_{\text{ap}}(T)$  of  $T$  if and only if  $T - \lambda I$  is not bounded below. In other words,  $\lambda$  belongs to  $\sigma_{\text{ap}}(T)$  if and only if there exists a sequence of unit vectors  $x_n$  such that  $(T - \lambda I)x_n \rightarrow 0$ . This is equivalent to asserting that  $T$  is not left invertible in  $\mathcal{B}(\mathcal{H})$  [11, p. 116] (see also [10, Prop. VII.6.4, Ex. VII.3.4]). Although in general, one only has

$$\sigma_{\text{ap}}(T) \cup \overline{\sigma_{\text{ap}}(T^*)} = \sigma(T), \quad (17)$$

(see [26, Pr. 73]) for a complex symmetric operator one obtains something significantly stronger [30, Lem. 4.1]. Indeed, if  $T = CT^*C$  for some conjugation  $C$ , then observe that  $\|(T - \lambda I)x\| = \|(T^* - \bar{\lambda}I)Cx\|$  for all  $x \in \mathcal{H}$  whence

$$\sigma_{\text{ap}}(T) = \overline{\sigma_{\text{ap}}(T^*)}.$$

Putting the preceding together with (17) we find that

$$\sigma(T) = \sigma_{\text{ap}}(T). \quad (18)$$

It turns out that (18) also holds under the weaker assumption that  $T$  is a norm limit of complex symmetric operators.

**Theorem 8.** *If  $T \in \overline{CSO}$ , then  $\sigma(T) = \sigma_{ap}(T) = \overline{\sigma_{ap}(T^*)}$ .*

*Proof.* By Lemma 6 there exists conjugations  $C_n$  such that  $C_n T^* C_n \rightarrow T$ . If  $\lambda$  belongs to  $\sigma_{ap}(T)$ , then there exists a sequence  $x_n$  of unit vectors such that  $(T - \lambda I)x_n \rightarrow 0$ . Thus

$$\begin{aligned} \|(T^* - \bar{\lambda}I)C_n x_n\| &= \|C_n T^* C_n x_n - \lambda x_n\| \\ &\leq \|(C_n T^* C_n - T)x_n\| + \|(T - \lambda I)x_n\|, \end{aligned}$$

which tends to zero. We therefore conclude that  $\sigma_{ap}(T) \subseteq \overline{\sigma_{ap}(T^*)}$ . The proof of the reverse containment is similar.  $\square$

The preceding theorem gives us a simple criterion for excluding certain operators from  $\overline{CSO}$ . For instance, the unilateral shift  $S$  is not a norm limit of complex symmetric operators since  $\sigma(S) = \mathbb{D}$  but  $\sigma_{ap}(S) = \partial\mathbb{D}$  [26, Prob. 82] (of course there are more direct ways to prove this, see [15, Ex. 2.14], or [16, Cor. 7]).

With these preliminaries in hand, the proof of part (i) of Theorem 7 is now quite simple.

*Pf. of Theorem 7, (i).* Since  $T^* e_1 = 0$  it follows that  $0 \in \sigma_{ap}(T^*)$  whence  $0 \in \sigma_{ap}(T)$  by Theorem 8. Thus there exists a sequence of unit vectors  $x_n$  such that  $T x_n \rightarrow 0$ . Suppose toward a contradiction that there exists some  $\delta > 0$  such that  $\alpha_n > \delta$  for all  $n$ . This implies that

$$\|T x_n\|^2 = \sum_{i=1}^{\infty} \alpha_i^2 |\langle x_n, e_i \rangle|^2 \geq \delta^2 \sum_{i=1}^{\infty} |\langle x_n, e_i \rangle|^2 = \delta^2 \|x_n\|^2 = \delta^2,$$

which contradicts the fact that  $T x_n \rightarrow 0$ .  $\square$

Before proving the second part of Theorem 7, we require a somewhat lengthy technical lemma and the following definition.

**Definition.** If  $T \in \mathcal{B}(\mathcal{H})$  and  $x \in \mathcal{H}$ , then we say that  $x$  is *shift-cyclic* for  $T$  if

$$\overline{\text{span}\{x, Tx, T^2x, \dots, T^*x, T^{*2}x, \dots\}} = \mathcal{H}.$$

The motivation for the preceding definition lies in the fact if  $T$  is an irreducible weighted shift, then each corresponding basis vector  $e_n$  of  $\mathcal{H}$  is shift-cyclic for  $T$ .

**Lemma 9.** *Suppose that  $T \in \overline{CSO}$ . If  $C_n$  is a sequence of conjugations such that  $C_n T^* C_n \rightarrow T$ ,  $x$  is a shift-cyclic vector for  $T$ , and  $C_n x$  converges, then  $T \in CSO$ .*

*Proof.* Let  $v \in \mathcal{H}$  and  $\epsilon > 0$ . Without loss of generality we may assume that  $\|T\| \leq 1$ ,  $\|x\| = 1$ , and  $\|v\| = 1$ . Since  $x$  is a shift-cyclic vector for  $T$ , there exists constants  $a_0, a_1, \dots, a_m$  and  $b_0, b_1, \dots, b_m$  such that

$$\left\| v - \left( \sum_{k=0}^m a_k T^k x + b_k T^{*k} x \right) \right\| < \frac{\epsilon}{6}.$$

Since each  $C_n$  is a conjugation it follows that

$$\left\| C_n v - \left( \sum_{k=0}^m \overline{a_k} C_n T^k + \overline{b_k} C_n T^{*k} \right) x \right\| < \frac{\epsilon}{6}. \quad (19)$$

Since  $\Delta_n = T - C_n T^* C_n \rightarrow 0$  by hypothesis, there exists  $N_1$  such that

$$n \geq N_1 \quad \Rightarrow \quad \|\Delta_n\| < \min \left\{ 1, \frac{\epsilon}{6M2^m} \right\},$$

where

$$M = \sum_{k=0}^m (|a_k| + |b_k|).$$

In particular,  $n \geq N_1$  implies that

$$\begin{aligned} \left\| C_n T^k - T^{*k} C_n \right\| &= \left\| T^k - C_n T^{*k} C_n \right\| \\ &= \left\| (C_n T^* C_n + \Delta_n)^k - (C_n T^* C_n)^k \right\| \\ &\leq \sum_{j=1}^k \binom{k}{j} \|C_n T^* C_n\|^{k-j} \|\Delta_n\|^j \\ &< \|\Delta_n\| \sum_{j=1}^k \binom{k}{j} \|T\|^{k-j} \|\Delta_n\|^{j-1} \\ &< \|\Delta_n\| 2^k \\ &< \frac{\epsilon}{6M2^{m-k}} \\ &< \frac{\epsilon}{6M} \end{aligned}$$

holds for  $0 \leq k \leq m$ . Thus for  $n \geq N_1$  we have

$$\begin{aligned} &\left\| \sum_{k=0}^m (\overline{a_k} C_n T^k + \overline{b_k} C_n T^{*k}) x - \sum_{k=0}^m (\overline{a_k} T^{*k} C_n + \overline{b_k} T^k C_n) x \right\| \\ &\leq \left\| \sum_{k=0}^m \overline{a_k} (C_n T^k - T^{*k} C_n) x \right\| + \left\| \sum_{k=0}^m \overline{b_k} (C_n T^{*k} - T^k C_n) x \right\| \\ &\leq \sum_{k=0}^m (|a_k| + |b_k|) \left\| C_n T^k - T^{*k} C_n \right\| \\ &< M \cdot \frac{\epsilon}{6M} \\ &= \frac{\epsilon}{6}. \end{aligned} \tag{20}$$

Since  $C_n x$  converges to some  $y$  by assumption, there exists  $N_2$  such that

$$n \geq N_2 \quad \Rightarrow \quad \|C_n x - y\| < \frac{\epsilon}{6M}.$$

Therefore  $n \geq N_2$  implies that

$$\begin{aligned} &\left\| \sum_{k=0}^m (\overline{a_k} T^{*k} C_n + \overline{b_k} T^k C_n) x - \sum_{k=0}^m (\overline{a_k} T^{*k} + \overline{b_k} T^k) y \right\| \\ &\leq \sum_{k=0}^m \left\| \overline{a_k} T^{*k} + \overline{b_k} T^k \right\| \|C_n x - y\| \\ &< M \cdot \frac{\epsilon}{6M} \end{aligned}$$

$$= \frac{\epsilon}{6}. \quad (21)$$

Putting this all together, if  $n \geq N = \max\{N_1, N_2\}$  we find that

$$\left\| C_n v - \left( \sum_{k=0}^m \overline{a_k} T^{*k} + \overline{b_k} T^k \right) y \right\| < \frac{\epsilon}{2}$$

by (19), (20), and (21). In particular,

$$n, n' \geq N \quad \Rightarrow \quad \|C_n v - C_{n'} v\| < \epsilon$$

whence  $C_n v$  is Cauchy and therefore converges. By Lemma 5, there exists a conjugation  $C$  such that  $C_n \rightarrow C$  (SOT).

Now consider  $T - CT^*C$ . Fix  $u \in \mathcal{H}$  and observe that

$$\begin{aligned} \|(T - CT^*C)u\| &= \|(CT - T^*C)u\| \\ &\leq \|(C - C_n)Tu\| + \|C_n Tu - T^*C_n u\| + \|T^*(C_n - C)u\| \\ &\leq \|(C - C_n)Tu\| + \|T - C_n T^* C_n\| + \|(C_n - C)u\|. \end{aligned}$$

Since  $C_n \rightarrow C$  (SOT), the first and third terms tend to zero. The second term tends to zero by hypothesis whence  $T = CT^*C$  so that  $T \in CSO$ .  $\square$

Now armed with Lemma 9, we complete the proof of Theorem 7.

*Pf. of Theorem 7, (ii).* Suppose toward a contradiction that no such subsequence exists. Letting  $\alpha_+ = \sup\{\alpha_n\}_{n=1}^\infty$ , it follows that there exists an index  $\ell$  such that  $\alpha_\ell = \alpha_+$ . Additionally, there exist  $\delta > 0$  and  $N \in \mathbb{N}$  such that

$$n \geq N \quad \Rightarrow \quad 0 < \alpha_n \leq \alpha_+ - \delta. \quad (22)$$

Since  $T \in \overline{CSO}$ , there exist conjugations  $C_n$  such that  $C_n T^* C_n \rightarrow T$  by Lemma 6. We now write

$$C_n e_\ell = \underbrace{\sum_{k=1}^N \langle C_n e_\ell, e_k \rangle e_k}_{x_n} + \underbrace{\sum_{k=N+1}^\infty \langle C_n e_\ell, e_k \rangle e_k}_{y_n}$$

and observe that  $\|x_n\|^2 + \|y_n\|^2 = 1$  whence

$$\begin{aligned} \|C_n T^* C_n e_\ell\|^2 &= \|T^*(x_n + y_n)\|^2 \\ &\leq \alpha_+^2 \|x_n\|^2 + (\alpha_+ - \delta)^2 \|y_n\|^2 \\ &\leq \alpha_+^2 \|x_n\|^2 + (\alpha_+^2 - 2\alpha_+ \delta + \delta^2) \|y_n\|^2 \\ &= \alpha_+^2 - \delta(2\alpha_+ - \delta) \|y_n\|^2 \\ &\leq \alpha_+^2 - \delta \alpha_+ \|y_n\|^2 \end{aligned}$$

since  $\delta < \alpha_+$  by (22). In particular, the preceding tells us that  $\|C_n T^* C_n e_\ell\| \leq \alpha_+$  from which it follows that

$$\begin{aligned} 0 &\leq \delta \alpha_+ \|y_n\|^2 \\ &\leq \alpha_+^2 - \|C_n T^* C_n e_\ell\|^2 \\ &= \|T e_\ell\|^2 - \|C_n T^* C_n e_\ell\|^2 \\ &= (\|T e_\ell\| + \|C_n T^* C_n e_\ell\|)(\|T e_\ell\| - \|C_n T^* C_n e_\ell\|) \\ &\leq 2\alpha_+(\|T e_\ell\| - \|C_n T^* C_n e_\ell\|) \end{aligned}$$

$$\leq 2\alpha_+ \|(T - C_n T^* C_n) e_\ell\|.$$

Since the preceding tends to zero, we conclude that  $y_n \rightarrow 0$ .

Now observe that the vectors  $x_n$  belong to unit ball of the finite-dimensional space  $\mathcal{H}_N = \text{span}\{e_i\}_{i=1}^N$ . Thus there exists a subsequence  $x_{n_k}$  of the  $x_n$  which converges to some  $x \in \mathcal{H}_N$ . Therefore

$$\|C_{n_k} e_\ell - x\| = \|x_{n_k} + y_{n_k} - x\| \leq \|x_{n_k} - x\| + \|y_{n_k}\| \rightarrow 0,$$

whence  $C_{n_k} e_\ell \rightarrow x$ . Since  $e_\ell$  is a shift-cyclic vector for  $T$  and  $C_{n_k} T^* C_{n_k} \rightarrow T$ , we conclude from Lemma 9 that  $T \in \overline{CSO}$ . However, this contradicts Lemma 1.  $\square$

## 5. APPROXIMATELY KAKUTANI SHIFTS

As we saw in Section 4, the Kakutani shift (2) demonstrates behavior which is typical of irreducible weighted shifts in  $\overline{CSO}$ . While Theorem 7 addresses some of the large-scale structure of the weight sequence  $\{\alpha_n\}_{n=1}^\infty$ , it sheds little light on the small-scale behavior of the weights. For instance, the Kakutani shift possesses a remarkable self-similarity in the sense that certain palindromic sequences are repeated infinitely often in its weight sequence. Remarkably, it turns out that an irreducible weighted shift which demonstrates some approximate level of self-similarity must belong to  $\overline{CSO}$ .

**Theorem 10.** *If  $T$  is an irreducible weighted shift with weights  $\{\alpha_n\}_{n=1}^\infty$  such that for each  $n \in \mathbb{N}$  and  $\epsilon > 0$  there exists an index  $c_{n,\epsilon} \geq n$  such that*

$$0 < \alpha_{c_{n,\epsilon}} < \epsilon, \tag{23}$$

and

$$1 \leq k \leq n \quad \Rightarrow \quad |\alpha_k - \alpha_{c_{n,\epsilon}-k}| < \epsilon, \tag{24}$$

then  $T \in \overline{CSO}$ .

Since the proof of the preceding theorem is somewhat long and involved, we defer it until the end of this section. We instead prefer to focus on a related conjecture and several consequences of our theorem.

Let us call an irreducible weighted shift  $T$  satisfying the hypotheses of Theorem 10 *approximately Kakutani*. We conjecture that this property is also necessary for an irreducible weighted shift to belong to  $\overline{CSO}$ .

**Conjecture 1.** *Every irreducible weighted shift in  $\overline{CSO}$  is approximately Kakutani.*

Among other things, the following corollary asserts that an irreducible weighted shift whose weight sequence is a suitable perturbation of the Kakutani sequence (2) also belongs to  $\overline{CSO}$ . In particular, this permits us to construct weighted shifts in  $\overline{CSO} \setminus CSO$  whose moduli have desired spectral properties.

**Corollary 11.** *If  $T$  is an irreducible weighted shift with weights  $\{\alpha_n\}_{n=1}^\infty$  such that*

- (i)  $\lim_{n \rightarrow \infty} \alpha_{2^n} = 0$ ,
- (ii)  $\lim_{n \rightarrow \infty} \sup\{|\alpha_k - \alpha_{2^n - k}| : 1 \leq k \leq 2^n\} = 0$ ,

then  $T$  belongs to  $\overline{CSO}$ .

*Proof.* Fix  $n$  and let  $\epsilon > 0$ . By (i), there exists  $K_1$  such that  $0 < \alpha_{2^k} < \epsilon$  holds whenever  $k \geq K_1$ . Letting

$$A_n = \sup\{|\alpha_k - \alpha_{2^n - k}| : 1 \leq k \leq 2^n\},$$

we obtain from (ii) that there is a  $K_2$  such that  $k \geq K_2$  implies that  $0 \leq A_k < \epsilon$ . Now let

$$c_{n,\epsilon} = 2^{K_1 + K_2 + n}.$$

Since  $K_1 + K_2 + n > K_1$  we have  $0 < \alpha_{c_{n,\epsilon}} < \epsilon$ , which is condition (23) of Theorem 10. Moreover, since  $K_1 + K_2 + n > K_2$ , we also have

$$|\alpha_k - \alpha_{c_{n,\epsilon} - k}| < \epsilon \quad (25)$$

for  $k < c_{n,\epsilon}$ , which is condition (24) from Theorem 10. Finally, since  $c_{n,\epsilon} > n$  we see that (25) holds whenever  $1 \leq k \leq n$ . By Theorem 10, we conclude that  $T$  belongs to  $\overline{CSO}$ .  $\square$

**Example 12.** Consider the weight sequence  $\{\alpha_n\}_{n=1}^\infty$  whose first few terms are

$$\begin{aligned} \alpha_1 &= 1, & \alpha_9 &= 1 + \frac{1}{3^7} + \frac{1}{3^8}, \\ \alpha_2 &= \frac{1}{2}, & \alpha_{10} &= \frac{1}{2} + \frac{1}{3^6} + \frac{1}{3^{10}}, \\ \alpha_3 &= 1 + \frac{1}{3^3}, & \alpha_{11} &= 1 + \frac{1}{3^3} + \frac{1}{3^5} + \frac{1}{3^{11}}, \\ \alpha_4 &= \frac{1}{4}, & \alpha_{12} &= \frac{1}{4} + \frac{1}{3^{12}}, \\ \alpha_5 &= 1 + \frac{1}{3^3} + \frac{1}{3^5}, & \alpha_{13} &= 1 + \frac{1}{3^3} + \frac{1}{3^{13}}, \\ \alpha_6 &= \frac{1}{2} + \frac{1}{3^6}, & \alpha_{14} &= \frac{1}{2} + \frac{1}{3^{14}}, \\ \alpha_7 &= 1 + \frac{1}{3^7}, & \alpha_{15} &= 1 + \frac{1}{3^{15}}, \\ \alpha_8 &= \frac{1}{8}, & \alpha_{16} &= \frac{1}{16}. \end{aligned}$$

In other words, the weights are defined inductively according to the following rules. Let  $\alpha_{2^n} = \frac{1}{2^n}$  and, having previously defined  $\alpha_1, \alpha_2, \dots, \alpha_{2^n - 1}$ , set

$$\alpha_{2^n + j} = \alpha_{2^n - j} + \frac{1}{3^{2^n + j}}. \quad (1 \leq j \leq 2^n)$$

By construction, the weight sequence  $\{\alpha_n\}_{n=1}^\infty$  satisfies the hypotheses of Corollary 11 and hence the corresponding weighted shift  $T$  belongs to  $\overline{CSO}$ . Moreover, a simple number-theoretic argument reveals that the  $\alpha_i$  are distinct whence  $\sigma(|T|) = \{0\} \cup \{\alpha_i\}_{i=1}^\infty$  where each  $\alpha_i$  which is not a power of two is an eigenvalue of multiplicity one. The essential spectrum  $\sigma_e(|T|)$  of  $|T|$  is simply  $\{0, 1, \frac{1}{2}, \frac{1}{4}, \dots\}$ .

Returning briefly to the subject of compact operators, we remark that the preceding example demonstrates that the fact that the eigenvalues of  $|T|$  tend to zero is essential in the proof of Theorem 4. We remark that this fact is used explicitly in equation (13). If the eigenvalues of  $|T|$  are allowed to accumulate elsewhere, then behavior such as that exhibited in Example 12 is possible. It is therefore difficult to conceive of a way in which the proof of Theorem 4 could be generalized to include certain classes non-compact operators.

Having made our remarks about Theorem 10, we now proceed to its proof.

*Pf. of Theorem 10.* Since this proof is somewhat long and intricate, let us first describe the general strategy. Using an iterative procedure, we first approximate the original irreducible weighted shift  $T$  by a certain direct sum  $T'$  of finite-dimensional

matrices of the form (1). In general,  $T'$  itself will not be a complex symmetric operator since there is no reason to believe that the matrices (1) produced will have any palindromic structure. We therefore approximate  $T'$  with a complex symmetric weighted shift  $T''$  constructed using an index juggling scheme.

Our first task is to select a strictly increasing sequence  $\{m_k\}_{k=0}^{\infty}$  of indices so that the weighted shift  $T'$  having the weight sequence  $\{\beta_i\}_{i=1}^{\infty}$  defined by

$$\beta_i = \begin{cases} \alpha_i & \text{if } i \neq m_k \text{ for all } k, \\ 0 & \text{if } i = m_k \text{ for some } k, \end{cases} \quad (26)$$

approximates  $T$  well in the operator norm while also being itself well-approximated by a complex symmetric weighted shift.

Given  $\epsilon > 0$ , find an index  $N$  such that

$$0 < \alpha_N < \frac{\epsilon}{4}. \quad (27)$$

This is made possible by the assumption (23). Now inductively define sequences  $\{\delta_k\}_{k=0}^{\infty}$  and  $\{m_k\}_{k=0}^{\infty}$  by setting

$$m_{-1} = 0, \quad m_0 = m_1 = N, \quad (28)$$

and

$$m_{2k+3} = c_{3m_{2k}, \delta_k} - m_{2k-1}, \quad (29)$$

$$m_{2k+2} = m_{2k+3} - m_{2k} + m_{2k-1}, \quad (30)$$

and

$$\delta_k = \frac{1}{8} \min \left\{ \alpha_1, \alpha_2, \dots, \alpha_{3m_{2k}}, \frac{\epsilon}{2^k} \right\}. \quad (31)$$

Unfortunately, it is not clear that the sequence  $\{m_k\}_{k=1}^{\infty}$  is strictly increasing. We must therefore establish the following claim.

**Claim.** *The sequence  $m_1, m_2, m_3, \dots$  is strictly increasing.*

*Pf. of Claim.* We induct on  $k$  in the statement

$$0 < m_{2k-1} < m_{2k} < m_{2k+1} < m_{2k+2}. \quad (32)$$

Let us first verify the base case  $k = 1$ , which is the statement

$$m_1 < m_2 < m_3 < m_4. \quad (33)$$

First observe that

$$3m_{2k} \leq c_{3m_{2k}, \delta_k} \quad (34)$$

for  $k \geq 0$ . Substituting (29) into (30) and using (34) then yields

$$2m_{2k} \leq m_{2k+2} \quad (35)$$

for  $k \geq 0$ . Therefore

$$\begin{aligned} m_0 &= m_1 = N && \text{by (28)} \\ &< 2m_0 = 3m_0 - m_0 \\ &\leq c_{3m_0, \delta_0} - m_0 = m_2 && \text{by (34)} \\ &= m_3 - m_0 < m_3 && \text{by (30)} \\ &= m_2 + m_0 < 2m_2 && \text{by (30)} \\ &\leq m_4. && \text{by (35)}. \end{aligned}$$



This establishes the base case (33).

Suppose now that (32) holds for some  $k \geq 1$ . Under this hypothesis, we wish to show that

$$m_{2k+1} < m_{2k+2} < m_{2k+3} < m_{2k+4}. \quad (36)$$

First note that  $m_{2k+1} < m_{2k+2}$  is already part of the induction hypothesis (32). The middle inequality of (36) follows from (30) since

$$m_{2k+2} = m_{2k+3} - (m_{2k} - m_{2k-1}) < m_{2k+3}$$

holds by the lower inequality in (32). To complete the induction, we need only verify the upper inequality in (36). This is established as follows:

$$\begin{aligned} m_{2k+3} &< m_{2k+3} + m_{2k-1} \\ &= m_{2k+2} + m_{2k} && \text{by (30)} \\ &< 2m_{2k+2} && \text{by (32)} \\ &\leq m_{2k+4}. && \text{by (35)} \end{aligned}$$

This completes the proof of the claim.  $\square$

Having constructed the desired sequence  $\{m_k\}_{k=0}^{\infty}$  of indices, we consider the weighted shift  $T'$  whose weight sequence is defined by (26). To prove that  $T'$  is a good approximation to  $T$  with respect to the operator norm, we must establish that each omitted weight  $\alpha_{m_k}$  is small. This is our next task.

In light of (27) and (28) we have

$$0 < \alpha_{m_0} = \alpha_{m_1} = \alpha_N < \frac{\epsilon}{4}. \quad (37)$$

Since  $m_3 = c_{3N, \delta_0}$  it follows from (23) and (31) that

$$0 < \alpha_{m_3} < \delta_0 < \frac{\epsilon}{4}. \quad (38)$$

By (29) and (30) we see that

$$m_{2k+2} + m_{2k} = m_{2k+3} + m_{2k-1} = c_{3m_{2k}, \delta_k}$$

which yields

$$\begin{aligned} |\alpha_{m_{2k+3}} - \alpha_{m_{2k-1}}| &< \delta_k < \frac{\epsilon}{2^{k+3}}, \\ |\alpha_{m_{2k+2}} - \alpha_{m_{2k}}| &< \delta_k < \frac{\epsilon}{2^{k+3}}, \end{aligned}$$

by (24). Using the triangle inequality and summing a finite geometric series yields

$$\begin{aligned} |\alpha_{m_{2k+2}} - \alpha_{m_0}| &< \frac{\epsilon}{4}, && \text{for } k \geq 0, \\ |\alpha_{m_{2k+3}} - \alpha_{m_1}| &< \frac{\epsilon}{4}, && \text{if } 2 \nmid k, \\ |\alpha_{m_{2k+3}} - \alpha_{m_3}| &< \frac{\epsilon}{4}, && \text{if } 2 \mid k. \end{aligned}$$

Since  $\alpha_{m_0}, \alpha_{m_1}, \alpha_{m_3} < \frac{\epsilon}{4}$  by (37) and (38), we conclude from the preceding that

$$0 < \alpha_{m_k} < \frac{\epsilon}{2}$$

for  $k \geq 0$ . This implies that  $\|T - T'\| < \frac{\epsilon}{2}$ .

Unfortunately, there is no reason to believe that  $T'$  belongs to  $CSO$ . Therefore our next task is to approximate  $T'$  with a complex symmetric weighted shift  $T''$ . At this point, it becomes more convenient to write  $T' = \bigoplus_{k=1}^{\infty} A_k$  where

$$A_1 = \begin{pmatrix} 0 & & & & & \\ \alpha_1 & 0 & & & & \\ & \alpha_2 & 0 & & & \\ & & \ddots & \ddots & & \\ & & & \alpha_{m_1-1} & 0 & \end{pmatrix}$$

and

$$A_k = \begin{pmatrix} 0 & & & & & \\ \alpha_{m_{k-1}+1} & 0 & & & & \\ & \alpha_{m_{k-1}+2} & 0 & & & \\ & & \ddots & \ddots & & \\ & & & \alpha_{m_k-1} & 0 & \end{pmatrix}$$

for  $k \geq 2$ . To make certain formulas work out, we let  $A_0 = A_1$ . Let

$$A'_k = \begin{pmatrix} 0 & & & & & \\ \alpha_{m_k-1} & 0 & & & & \\ & \alpha_{m_k-2} & 0 & & & \\ & & \ddots & \ddots & & \\ & & & \alpha_{m_{k-1}+1} & 0 & \end{pmatrix}$$

denote the matrix obtained from  $A_k$  by reversing the order of the weights along the first subdiagonal.

Consider the relationship between the matrices  $A'_{2k+3}$  and  $A_{2k}$ . The  $\ell$ th subdiagonal entry of  $A'_{2k+3}$  is  $\alpha_{m_{2k+3}-\ell}$  while the  $\ell$ th subdiagonal entry of  $A_{2k}$  is  $\alpha_{m_{2k-1}+\ell}$ . Since the sum of these indices is

$$(m_{2k+3} - \ell) + (m_{2k-1} + \ell) = m_{2k+3} + m_{2k-1} = c_{3m_{2k}, \delta_k}$$

by (29), it follows from (24) that

$$|\alpha_{m_{2k+3}-\ell} - \alpha_{m_{2k-1}+\ell}| < \delta_k < \frac{\epsilon}{2}.$$

In particular, this tells us that

$$\|A'_{2k+3} - A_{2k}\| < \frac{\epsilon}{2}.$$

Since  $A_0 = A_1$  we observe that

$$T' = \bigoplus_{k=1}^{\infty} A_k = A_0 \oplus \bigoplus_{k=2}^{\infty} A_k \cong \bigoplus_{k=0}^{\infty} (A_{2k+3} \oplus A_{2k}) = S'.$$

Finally define  $T''$  and  $S''$  by

$$T'' = A'_3 \oplus \left( \bigoplus_{j=2}^{\infty} \begin{cases} A_j & \text{if } 2 \nmid j, \\ A'_{j+3} & \text{if } 2 \mid j, \end{cases} \right) \cong \bigoplus_{k=0}^{\infty} \underbrace{(A_{2k+3} \oplus A'_{2k+3})}_{\in CSO} = S''.$$

The operator  $S''$  belongs to  $CSO$  since it is a direct sum of matrices  $A_{2k+3} \oplus A'_{2k+3}$  of the form (1) whose entries on the first subdiagonal are palindromic. Therefore

$$\|T' - T''\| = \|S' - S''\| < \frac{\epsilon}{2}$$

whence

$$\|T - T''\| \leq \|T - T'\| + \|T' - T''\| < \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon.$$

Thus  $T$  belongs to  $\overline{CSO}$ , as claimed.  $\square$

#### ACKNOWLEDGMENTS

We wish to thank W.R. Wogen for his numerous comments and suggestions. We also greatly appreciate the careful eye of the anonymous referee, who pointed out a number of minor mistakes in the original manuscript.

#### REFERENCES

- [1] Tsuyoshi Ando. Aluthge transforms and the convex hull of the spectrum of a Hilbert space operator. In *Recent advances in operator theory and its applications*, volume 160 of *Oper. Theory Adv. Appl.*, pages 21–39. Birkhäuser, Basel, 2005. 1
- [2] Jorge Antezana, Enrique R. Pujals, and Demetrio Stojanoff. The iterated Aluthge transforms of a matrix converge. *Adv. Math.*, 226(2):1591–1620, 2011. 1
- [3] C. Benhida and E. H. Zerouali. Backward Aluthge iterates of a hyponormal operator and scalar extensions. *Studia Math.*, 195(1):1–10, 2009. 1
- [4] Fernanda Botelho and James Jamison. Elementary operators and the Aluthge transform. *Linear Algebra Appl.*, 432(1):275–282, 2010. 1
- [5] Gilles Cassier and Jérôme Verliat. Stability for some operator classes by Aluthge transform. In *Operator theory live*, volume 12 of *Theta Ser. Adv. Math.*, pages 51–67. Theta, Bucharest, 2010. 1
- [6] I. Chalendar, E. Fricain, and D. Timotin. On an extremal problem of Garcia and Ross. *Oper. Matrices*, 3(4):541–546, 2009. 1
- [7] Nicolas Chevrot, Emmanuel Fricain, and Dan Timotin. The characteristic function of a complex symmetric contraction. *Proc. Amer. Math. Soc.*, 135(9):2877–2886 (electronic), 2007. 1
- [8] J. A. Cima, W. T. Ross, and W. R. Wogen. Truncated Toeplitz operators on finite dimensional spaces. *Oper. Matrices*, 2(3):357–369, 2008. 1
- [9] Joseph A. Cima, Stephan Ramon Garcia, William T. Ross, and Warren R. Wogen. Truncated Toeplitz operators: spatial isomorphism, unitary equivalence, and similarity. *Indiana Univ. Math. J.*, 59(2):595–620, 2010.
- [10] John B. Conway. *A course in functional analysis*, volume 96 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 1990. 4, 6, 9
- [11] John B. Conway. *A course in operator theory*, volume 21 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2000. 2, 3, 9
- [12] Jeffrey Danciger, Stephan Ramon Garcia, and Mihai Putinar. Variational principles for symmetric bilinear forms. *Math. Nachr.*, 281(6):786–802, 2008. 4, 5
- [13] Ken Dykema and Hanne Schultz. Brown measure and iterates of the Aluthge transform for some operators arising from measurable actions. *Trans. Amer. Math. Soc.*, 361(12):6583–6593, 2009. 1
- [14] George R. Exner. Aluthge transforms and  $n$ -contractivity of weighted shifts. *J. Operator Theory*, 61(2):419–438, 2009. 1
- [15] Stephan Ramon Garcia. Conjugation and Clark operators. In *Recent advances in operator-related function theory*, volume 393 of *Contemp. Math.*, pages 67–111. Amer. Math. Soc., Providence, RI, 2006. 1, 10
- [16] Stephan Ramon Garcia. Means of unitaries, conjugations, and the Friedrichs operator. *J. Math. Anal. Appl.*, 335(2):941–947, 2007. 1, 10
- [17] Stephan Ramon Garcia. Aluthge transforms of complex symmetric operators. *Integral Equations Operator Theory*, 60(3):357–367, 2008. 1
- [18] Stephan Ramon Garcia. The eigenstructure of complex symmetric operators. In *Recent advances in matrix and operator theory*, volume 179 of *Oper. Theory Adv. Appl.*, pages 169–183. Birkhäuser, Basel, 2008. 1
- [19] Stephan Ramon Garcia and Daniel E. Poore. On the norm closure problem for complex symmetric operators. *Proc. Amer. Math. Soc.* to appear. 1, 2

- [20] Stephan Ramon Garcia and Mihai Putinar. Complex symmetric operators and applications. *Trans. Amer. Math. Soc.*, 358(3):1285–1315 (electronic), 2006. 1, 2, 3
- [21] Stephan Ramon Garcia and Mihai Putinar. Complex symmetric operators and applications. II. *Trans. Amer. Math. Soc.*, 359(8):3913–3931 (electronic), 2007. 1
- [22] Stephan Ramon Garcia and William T. Ross. A non-linear extremal problem on the Hardy space. *Comput. Methods Funct. Theory*, 9(2):485–524, 2009. 1
- [23] Stephan Ramon Garcia and Warren R. Wogen. Complex symmetric partial isometries. *J. Funct. Anal.*, 257(4):1251–1260, 2009. 1, 2, 4, 6
- [24] Stephan Ramon Garcia and Warren R. Wogen. Some new classes of complex symmetric operators. *Trans. Amer. Math. Soc.*, 362(11):6065–6077, 2010. 1, 2
- [25] T. M. Gilbreath and Warren R. Wogen. Remarks on the structure of complex symmetric operators. *Integral Equations Operator Theory*, 59(4):585–590, 2007. 1
- [26] Paul Richard Halmos. *A Hilbert space problem book*, volume 19 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 1982. Encyclopedia of Mathematics and its Applications, 17. 2, 3, 9, 10
- [27] Huajun Huang and Tin-Yau Tam. Aluthge iteration in semisimple Lie group. *Linear Algebra Appl.*, 432(12):3250–3257, 2010. 1
- [28] N. Jacobson. Normal Semi-Linear Transformations. *Amer. J. Math.*, 61(1):45–58, 1939. 1
- [29] S. Jung, E. Ko, and J. Lee. On scalar extensions and spectral decompositions of complex symmetric operators. *J. Math. Anal. Appl.* preprint. 1
- [30] S. Jung, E. Ko, M. Lee, and J. Lee. On local spectral properties of complex symmetric operators. *J. Math. Anal. Appl.*, 379:325–333, 2011. 1, 9
- [31] Chun Guang Li, Sen Zhu, and Ting Ting Zhou. Foguel operators with complex symmetry. preprint. 1
- [32] Charles E. Rickart. *General theory of Banach algebras*. The University Series in Higher Mathematics. D. van Nostrand Co., Inc., Princeton, N.J.-Toronto-London-New York, 1960. 3
- [33] D. Sarason. Algebraic properties of truncated Toeplitz operators. *Oper. Matrices*, 1(4):491–526, 2007. 1
- [34] Issai Schur. Ein Satz ueber quadratische Formen mit komplexen Koeffizienten. *Amer. J. Math.*, 67:472–480, 1945. 1
- [35] N. Sedlock. Algebras of truncated Toeplitz operators. *Oper. Matrices* (to appear) <http://arxiv.org/abs/1011.3425>. 1
- [36] Nicholas Alexander Sedlock. *Properties of truncated Toeplitz operators*. ProQuest LLC, Ann Arbor, MI, 2010. Thesis (Ph.D.)–Washington University in St. Louis. 1
- [37] Carl Ludwig Siegel. Symplectic geometry. *Amer. J. Math.*, 65:1–86, 1943. 1
- [38] Teiji Takagi. On an algebraic problem related to an analytic theorem of Carathéodory and Fejér and on an allied theorem of Landau. *Japan J. Math.*, 1:83–93, 1925. 1
- [39] James E. Tener. Unitary equivalence to a complex symmetric matrix: an algorithm. *J. Math. Anal. Appl.*, 341(1):640–648, 2008. 1
- [40] Xiao Huan Wang and Zong Sheng Gao. Some equivalence properties of complex symmetric operators. *Math. Pract. Theory*, 40(8):233–236, 2010. 1
- [41] Xiaohuan Wang and Zongsheng Gao. A note on Aluthge transforms of complex symmetric operators and applications. *Integral Equations Operator Theory*, 65(4):573–580, 2009. 1
- [42] Sergey M. Zagorodnyuk. On a  $J$ -polar decomposition of a bounded operator and matrix representations of  $J$ -symmetric,  $J$ -skew-symmetric operators. *Banach J. Math. Anal.*, 4(2):11–36, 2010. 1
- [43] Sen Zhu and Chun Guang Li. Complex symmetric weighted shifts. preprint. 3
- [44] Sen Zhu, Chun Guang Li, and You Qing Ji. The class of complex symmetric operators is not norm closed. *Proc. Amer. Math. Soc.* to appear. 1, 2, 3

DEPARTMENT OF MATHEMATICS, POMONA COLLEGE, CLAREMONT, CALIFORNIA, 91711, USA  
*E-mail address:* [Stephan.Garcia@pomona.edu](mailto:Stephan.Garcia@pomona.edu)  
*URL:* <http://pages.pomona.edu/~sg064747>