

1-1-1971

# Measurement of Nuclear Magnetic Dipole Moment of Li-8 by Implantation in Metal Foils

Richard C. Haskell  
*Harvey Mudd College*

R. L. Williams Jr.

Leon Madansky  
*Johns Hopkins University*

---

## Recommended Citation

Haskell, Richard C., R. L. Williams Jr., and L. Madansky. "Measurement of nuclear magnetic dipole moment of Li-8 by implantation in metal foils." *Bulletin of the American Physical Society* 16.12 (1971): 1417.

This Abstract is brought to you for free and open access by the HMC Faculty Scholarship at Scholarship @ Claremont. It has been accepted for inclusion in All HMC Faculty Publications and Research by an authorized administrator of Scholarship @ Claremont. For more information, please contact [scholarship@cuc.claremont.edu](mailto:scholarship@cuc.claremont.edu).

structing high energy scattering amplitudes. The effect of long-range correlations has been estimated by using a macroscopic collective model to describe the low-lying  $2^+(4.44 \text{ MeV})$ ,  $0^+(7.6 \text{ MeV})$ , and  $3^-(9.6 \text{ MeV})$  states. These states make a correction to the optical potential through the second-order correlation function which usually includes only Pauli and center-of-mass correlations. These corrections considerably improve the agreement between theory and experiment. In addition, the differential cross-section for scattering and excitation of the low-lying states has been calculated using the same model and is in substantial agreement with experiment.

FF 2 Partial-Wave Contribution in  $\pi$ -Nucleus Scatterings Near the 3-3 Resonance. RYOICHI SEKI, San Fernando Valley State College.--Within the frame work of the multiple scattering formalism which we proposed previously, we have examined how much each ( $\pi$ -nucleus) partial wave contributes to the  $\pi$ -nucleus total cross sections near the 3-3 resonance. It is found that the appreciable contribution comes from partial waves up to a little larger than  $kR$ , where  $k$  is the pion wave number and  $R$  is the geometric radius of the nucleus seen by the pion, and also found that the largest contribution comes from partial waves  $\sim kr^{-1/2}$ , where  $r$  is the RMS radius of the nucleus: The  $\pi$ -nucleus scatterings show remarkably well a feature of the simple black-body scatterings caused by formation of the 3-3 resonance in the nucleus<sup>1</sup>. The result of our calculation seems to be in agreement with a partial wave analysis of the  $\pi$ -C<sup>12</sup> scatterings by use of an impact parameter method<sup>2</sup>.

<sup>1</sup> R. Seki, Bull. Am. Phys. Soc. 14, 52 (1969) and Phys. Rev. C 3, 454 (1971).

<sup>2</sup> J. Beiner and P. Huguenin, Helv. Phys. Acta 43, 421 (1970).

FF 3 Deuteron Wave Function at Large  $r$ . H. WONG\*, C. BURNAP\*†, and J.S. LEVINGER, Rensselaer Polytech. -- We substitute analytical expressions for the deuteron wave function ( $r \geq 4 \text{ F.}$ ) into the Schrödinger equation to find the central and tensor potentials.<sup>1</sup> We compare with OBE<sup>2</sup> and Reid. We consider: i) Iwadare's wave function<sup>3</sup> which gives a poor fit; ii) modified Iwadare, varying 4 of his parameters (to  $B = 0.350$ ,  $\beta = 1.235 \text{ F}^{-1}$ ,  $D = 2.65 \text{ F}^2$ ,  $\gamma = 0.640 \text{ F}^{-1}$ ) to obtain a good fit to Reid's potentials; iii) Hulthen-Sugawara wave function<sup>3</sup> which gives a poor fit.

\* National Science Foundation Undergraduate Research Participant.

† Now at Harvard University.

<sup>1</sup> Burnap et al. Phys. Lett. 33B, 337 (1970).

<sup>2</sup> Iwadare et al. Prog. Theor. Phys. 16, 455 (1956).

<sup>3</sup> Hulthen and Sugawara, Physics Handbook, 39, 91 (1957).

FF 4 Photodisintegration of <sup>13</sup>C Leading to Excited States of <sup>12</sup>C and <sup>12</sup>B. E. J. WINHOLD, Rensselaer Polytechnic Institute and AERE Harwell, UK, E. M. BOWEY, B. H. PATRICK, and J. M. REID\*, AERE Harwell, UK.-- Ge(Li) detector measurements were made of gamma ray spectra from the <sup>13</sup>C( $\gamma, n^*$ ) reactions. These reactions were initiated by bremsstrahlung from the Harwell linac, and excitation functions for the production of particular gamma lines were obtained as a function of bremsstrahlung end-point energy over the range from 15 to 40 MeV. The T=0 4.44 MeV state and the T=1 15.1 MeV state of <sup>12</sup>C are both strongly populated, as Murray has observed. However the cross section for ( $\gamma, n$ ) leading to the 15.1 MeV level is peaked at 25 MeV, while that leading to the 4.44 MeV level peaks below 15 MeV. The 0.95 MeV state in <sup>12</sup>B is weakly

excited. These results appear generally consistent with <sup>13</sup>C giant resonance calculations<sup>2</sup> which predict a substantial isospin splitting of the resonance.

\*Permanent Address: University of Glasgow.

<sup>1</sup> K. M. Murray, Nuovo Cimento Letters 1, 571 (1971).

<sup>2</sup> B. R. Easlea, Phys. Letters 1, 163 (1962);

Wessday et al, Nucl. Phys. 61, 269 (1965).

FF 5 Observation of Quadrupole Splitting of <sup>12</sup>B in a Single Crystal.\* R. L. WILLIAMS, JR., R. C. HASKELL† and L. MADANSKY, The Johns Hopkins University--The quadrupole coupling of <sup>12</sup>B implanted in <sup>9</sup>Be has been observed using a single crystal of Be. One sees a narrow resonance line, the location of which depends in the normal way on the orientation of the crystalline c-axis with respect to the external magnetic field direction. The coupling constant is given by  $e^2qQ/h = 54.9(6) \text{ kHz}$ . This is consistent with our previous measurement<sup>1</sup> using a Be foil. Using the field gradient at <sup>9</sup>Be lattice sites, calculated by Pomerantz and Das,<sup>2</sup> one finds  $Q(^{12}\text{B}) = 34.6 \text{ mb}$ .

\*Work supported by U. S. Atomic Energy Commission.

†NSF Predoctoral Fellow.

<sup>1</sup> R. L. Williams, Jr., L. Pfeiffer, J. C. Wells, Jr and L. Madansky, Phys. Rev. C2, 1219 (1970).

<sup>2</sup> M. Pomerantz and T. P. Das, Phys. Rev. 119, 70 (1960).

FF 6 Comparison of the Precessions of Angular Correlations Produced by Magnetic Dipole and Axially Symmetric Quadrupole Interactions.\* / O. KLEPPER, MIT -

According to the semi-classical vector model a magnetic dipole interaction or an axially symmetric quadrupole interaction can result in spin precessions, but in the latter case the spin precesses in opposite directions for +M and -M substates. Developing the attenuation factors  $G_{K, K_1, K_2}^{\mu, \mu_1, \mu_2}(t)$  for the electric case analogously to the magnetic, one can show that a polarization of the nucleus results in a net rotation of the angular correlation. Generally, one gets a superposition of correlations precessing with different frequencies  $n\omega$ , and a non-rotating part due to the alignment of the nucleus.<sup>1</sup> Experiments at the Rutgers-Bell-Tandem will be mentioned which use Coulomb excitation with scattering angles  $\theta < 180^\circ$  to excite, polarize, and implant nuclei into single crystals and allow measurement of the sign of  $\omega$ .

\* Submitted by A.M. Bernstein

† Work supported in part by the U.S. Atomic Energy Comm.

<sup>1</sup> L. Grodzins and O. Klepper, Phys. Rev. C3 (1971), 1019.

FF 7 Measurement of the Nuclear Magnetic Dipole Moment of <sup>6</sup>Li by Implantation in Metal Foils.\* R. C. HASKELL†, R. L. WILLIAMS, JR., and L. MADANSKY, The Johns Hopkins University--Polarized <sup>6</sup>Li nuclei have been produced through the <sup>7</sup>Li(d,p) reaction using the 3.5-MeV Van de Graaff accelerator at Brookhaven National Laboratory. The observed polarization was a slowly-varying function of deuteron energy over the range 1.3-2.9 MeV, reaching a maximum of about +1.6%. The recoiling nuclei were stopped in Au, Pt and Pd foils and the effective dipole moments were measured by a resonant depolarization method. The results were  $1.65362(22)\mu_N$ ,  $1.65288(20)\mu_N$  and  $1.65270(30)\mu_N$ , respectively. These are consistent with the work of Connor, who found  $\mu(^6\text{Li}) = 1.6530(8)\mu_N$  in a LiF crystal. An upper limit for the <sup>6</sup>Li quadrupole moment will also be discussed.

\*Work supported by the U.S. Atomic Energy Commission.

†NSF Predoctoral Fellow.

<sup>1</sup> D. Connor, Phys. Rev. Letters 3, 429 (1959).