


2007

Index Volume 23

Follow this and additional works at: <http://scholarship.claremont.edu/aliso>

Recommended Citation

(2007) "Index Volume 23," *Aliso: A Journal of Systematic and Evolutionary Botany*: Vol. 23: Iss. 1, Article 47.
Available at: <http://scholarship.claremont.edu/aliso/vol23/iss1/47>



Index

Primary indexing terms include plant taxa, contributing authors, major subjects, and most key words. Additional salient topics are featured if deemed appropriate, including geographic regions, genes and genetic loci, analytical software and concepts, selected informal plant groups, and vegetation types. For reasons of brevity, taxonomic names at the rank of species and below are omitted, except where particular species constitute the focus of research. New taxa and combinations are in **boldface**. Organisms other than monocotyledonous plants (e.g., outgroup taxa, model organisms, and pollinators, etc.) are listed where they augment the discussion of monocot comparative biology. First authors of papers are cited in conjunction with the title of their article, followed by coauthor names (“et al.” follows the first author’s name in contributions having three or more authors). Coauthors are entered without the article title. Numbers precede letters in the alphabetical order of the index, unless they are part of a composite term where a subsequent component determines placement: e.g., 18S rDNA is arranged under DNA. The connectors “and,” “as,” “in,” “or,” and “versus” are ignored for the purpose of alphabetical placement. Indexing terms are located to pages where their discussion assumes significant importance and not where they represent merely a passing mention.

Abildgaardia

- in Abildgaardieae phylogeny 149–164
- in Cyperaceae phylogeny 79
- 5.8S rDNA insertion in 87, 94
- embryo type of 150
- geographic distribution of 149
- ITS analysis of 155–156, 158–160
- morphology of 150
- photosynthetic pathways in (C_3 and C_4) 101–102, 111, 161
- rbcL* analysis of 79
- relationship with *Bulbostylis* 150, 161
- relationship with *Crosslandia* 161
- relationship with *Fimbristylis* 104, 150, 161–162
- trnL*–*F* analysis of 155–158, 161–162
- trnL*–*F* and ITS analysis of, combined 151–154, 159–162

Abildgaardieae

- clades of 150–151
- in Cyperaceae phylogeny 73, 79, 81, 149–150
- in Cyperales phylogeny 68
- 5.8S rDNA insertion in 87, 94, 96
- embryo types of 149–150
- ITS analysis of 151–156, 158–162
- photosynthetic pathways in 161
- phylogeny of 149–164
- polytomy of 73
- rbcL* analysis of 73, 79, 81, 149–151
- relationship with Arthrostylideae 81, 149–162
- relationship with Eleocharideae 150
- relationship with Fuireneae 150
- relationship with Scirpeae 150
- sister relationships in 150–151

trnL–*F* analysis of 151–162

trnL–*F* and ITS analysis of, combined 151–154, 159–162

Abolboda

- androecium of 230
- gynoecium of 231
- inflorescence of 227–233
- perianth of 228–230

Abromeitiella

- in Bromeliaceae phylogeny 7–9, 14–17
- CAM photosynthesis by 11
- epiphytism of 12
- geographic origin and distribution of 10–11, 19
- leaf and trichome anatomy of 13–14
- ndhF* analysis of 7–9, 14–17
- pollination of 13, 21

Acamptocladus

- in Chloridoideae phylogeny 574, 576
- relationship with *Eragrostis* 595–596, 599, 602

Acanthostachys

- in Bromeliaceae phylogeny 34, 36
- trnL*, *trnL*–*F*, and *trnT*–*F* analysis of, combined 34
- trnL* and *trnL*–*F* analysis of, combined 36

Achlaena

- biogeography of 509–510, 516
- New World 505, 507, 509–510, 516
- in Panicoideae phylogeny 503, 505, 507, 516
- photosynthetic pathway of 516

Achlyphila

- androecium of 230
- gynoecium of 231
- inflorescence of 227–233
- perianth of 228–230

Achnatherum

- geographic distribution of 349, 355
- ITS analysis of 351, 355–357
- matK* analysis of 342
- ndhF* analysis of 342
- ndhH* analysis of 342
- in Poaceae phylogeny 342
- rbcL* analysis of 342
- relationship with *Anemathelle* 355
- relationship with *Jarava* 349–350, 355, 359
- relationship with *Oryzopsis* 350
- in Stipeae phylogeny 349–351, 355–357

Acostia

- biogeography of 509–510, 521
- New World 505, 508–510, 521
- in Panicoideae phylogeny 505, 508, 521
- photosynthetic pathway of 521

Acrachne

- in Chloridoideae phylogeny 581
- New World 581

Acroceras

- biogeography of 511, 519
- New World 505, 508, 511, 519
- in Panicoideae phylogeny 505, 508, 519
- photosynthetic pathway of 519
- in Poaceae phylogeny 254

Actinocladum

- branching of 308
- buds of 308

Actinoschoenus

- in Cyperaceae phylogeny 81, 149
- geographic distribution of 149
- ITS analysis of 156
- photosynthetic pathway in 102, 111
- trnL*–*F* analysis of 157–158
- trnL*–*F* and ITS analysis of, combined 159

Actinoscirpus

- absence of 5.8S rDNA insertion in 88, 92, 94, 96
- in Cyperaceae phylogeny 79, 81
- photosynthetic pathway in 102, 111
- rbcL* analysis of 79
- relationship with *Schoenoplectus* 81

Adaptive radiation, of Bromeliaceae 5, 12–14, 19–22

adh1, in Poaceae analysis 252

Aechmea

- AFLP analysis of 37–39
- in Bromeliaceae (Bromelioideae) phylogeny 8–9, 27, 29, 34–39, 41
- CAM photosynthesis by 11
- epiphytism of 12
- geographic origin and distribution of 10
- leaf and trichome anatomy of 14, 39–40
- pollination of 13
- trnL*, *trnL*–*F*, and *trnT*–*F* analysis of, combined 34
- trnL* and *trnL*–*F* analysis of, combined 35–36

Aechmea subgen. *Aechmea* 29, 41

Aechmea subgen. *Gravisia* 39

Aechmea subgen. *Lamprococcus* 29, 35, 39, 41

Aechmea subgen. *Macrochordion* 29

Aechmea subgen. *Ortgiesia* 29, 35, 39

Aechmea subgen. *Platyaechmea* 29

Aechmea subgen. *Podaechmea* 29, 38–39

Aechmea subgen. *Pothuava* 29, 38–39, 41

Aegilops

- chloroplast DNA analysis of 366, 368
- DMCI* analysis of 365, 367
- EF-G* analysis of 363, 367
- mitochondrial genome of 366
- morphological analysis of 365, 368
- total evidence (combined) analysis of 367–369
- in Triticeae phylogeny 362–363, 365–369

Aegilops triuncialis

- bet-hedging strategy of 296, 298
- dimorphism in 296–298
- inflorescence of 296
- invasion by 295–300
- maternal effect on germination 296–298
- phenotypic plasticity in 295–300
- seed dormancy in 296–298
- seed mass of 296–297
- seed production in 298
- sibling effect on germination 296–298
- spikelets of 296

Aegopogon

- biogeography of 587, 593
- breeding system of 608–610
- in Chloridoideae phylogeny 566, 570–571, 573–574, 577, 581, 584
- ITS analysis of 566, 571
- ITS and *trnL*–*F* analysis of, combined 573
- morphological analysis of 584–585
- New World 581, 584, 587, 593
- trnL*–*F* analysis of 566, 570

Africa

- Bromeliaceae origin and distribution in 5, 7, 10–12
- Chloridoideae in 585

Afrotrilepis, photosynthetic pathway in 102, 111

Agenium

- biogeography of 509, 511, 515
- New World 504–505, 509, 511, 515
- in Panicoideae phylogeny 504–505, 515
- photosynthetic pathway of 515

Agmatoploidy

- of *Carex* 63–64, 193
- of Cyperales 63–64, 66–67
- of *Rhynchospora* 193

Agnesia

- biogeography of 310
- in Olyrinae (Olyreae) phylogeny 310

Agropyron

- chloroplast DNA analysis of 366, 368
- DMCI* analysis of 366–367, 369–370

- EF-G* analysis of 366–367, 369–370
 mitochondrial genome of 366
 morphological analysis of 365
 relationship with *Elymus* 372, 374, 377
 total evidence (combined) analysis of 367–369
 in Triticeae phylogeny 365–370
- Agrostideae, in Pooideae phylogeny 380
- Agrostidinae
 as outgroup in *Poa* study 422
 in Pooideae phylogeny 346
- Agrostis*
 as outgroup in *Poa* study 422
 in Poaceae phylogeny 256
 in Pooideae phylogeny 343, 346–347
- Aira*, in Pooideae phylogeny 343, 346–347
- Airinae
 in Pooideae phylogeny 346–347
 relationship with *Poa* 421–422, 427
- Alatae 194
- Alcantarea*
 in Bromeliaceae phylogeny 28, 34
trnL, *trnL-F*, and *trnT-F* analysis of, combined 34
- Alinula*
 in Cyperaceae phylogeny 80
 photosynthetic pathway in 102, 111
rbcL analysis of 80
- Allolepsis*
 biogeography of 587, 593
 in Chloridoideae phylogeny 566, 570–571, 573, 581
 ITS analysis of 566, 571
 ITS and *trnL-F* analysis of, combined 573
 New World 581, 587, 593
trnL-F analysis of 566, 570
- Allopolyploids
 of *Bromus* 468–471
 of *Elymus* 372–379
 of *Eragrostis* 596, 602
- Alloteropsis*
 New World 505, 508
 in Panicoideae phylogeny 505, 508
- Alopecurinae
 placement of genera in 423–424
 in Poaceae phylogeny 421
 in Pooideae phylogeny 346
 relationship with *Poa* 421–427, 431–433
- Alopecurus*
 in Poaceae phylogeny 421
 in Pooideae phylogeny 343, 346
 relationship with *Poa* 421, 424, 427, 431, 433
 tribal/subtribal placement of 423
- Altai Mountain region, variability of *Poa* in 435, 448
- Altoparadisium*
 biogeography of 509, 511, 516
 New World 505, 507, 509, 511, 516
 in Panicoideae phylogeny 505, 507, 516
 photosynthetic pathway of 516
- Alves, Marccus V. 72–83
- Alvimia*
 in Arthrostylidiinae (Bambuseae) phylogeny 308
 biogeography of 308
 branching of 308
 buds of 308
 fruit of 308
 spikelets of 308
- Amazon Basin, Bromeliaceae origin and distribution in 5, 7, 10–12
- Amazonia, Bambusoideae in 311
- Amazonian Region, Panicoideae of 510–511
- Amblyopyrum*
 chloroplast DNA analysis of 366, 368
DMC1 analysis of 365, 367
EF-G analysis of 367
 mitochondrial genome of 366
 morphological analysis of 365–366
 total evidence (combined) analysis of 367–369
 in Triticeae phylogeny 365–369
- Ammophila*, in Pooideae phylogeny 346
- Ampelodesmeae, in Poaceae phylogeny 252–253
- Ampelodesmos*
 ITS analysis of 351–356
matK analysis of 342
ndhF analysis of 342
ndhH analysis of 342
 in Poaceae phylogeny 242–244, 342
 in Pooideae phylogeny 345–346
rbcL analysis of 342
 in Stipeae phylogeny 350–356
- Amphibromus*, in Pooideae phylogeny 343, 346
- Amphicarpum*
 biogeography of 509–510, 521
 cleistogamy and chasmogamy in 287–293
 New World 505, 508–510, 521
 in Panicoideae phylogeny 505, 508, 521
 photosynthetic pathway of 521
- Amphilophis*, in Andropogoneae phylogeny 532
- Amphiphyllum* 6
- Amphipogon*
matK analysis of 342
ndhF analysis of 342
ndhH analysis of 341–342, 344
 in Poaceae phylogeny 242–244, 341–342
rbcL analysis of 342
- Amphiscirpus*, photosynthetic pathway in 102, 111
- Amplified fragment length polymorphism (AFLP)
 in Bromeliaceae (Bromelioideae) analysis 29–33, 35–39, 41
 in *Carex* sect. *Ovales* analysis 193–203
- Ananas*
 in Bromeliaceae phylogeny 8–9, 34–36
 CAM photosynthesis by 11
 epiphytism of 12
 geographic origin and distribution of 10, 12

- leaf and trichome anatomy of 14
- morphology of 35
- ndhF* analysis of 8–9
- trnL*, *trnL*-F, and *trnT*-F analysis of, combined 34
- trnL* and *trnL*-F analysis of, combined 35–36
- Anarthriaceae, relationship with Bromeliaceae 27–28
- Anatherostipa*, nomenclature for 350
- Andean Region, Panicoideae of 510–512
- Andes
 - Bambusoideae in 311
 - Bromeliaceae origin and distribution in 5, 7, 10–12, 18–19
- Androdioecy/andromonoecy 605–614
- Androlepis*
 - in Bromeliaceae phylogeny 34, 36
 - trnL*, *trnL*-F, and *trnT*-F analysis of, combined 34
 - trnL* and *trnL*-F analysis of, combined 36
- Andropogon*
 - in Andropogoneae phylogeny 262, 530–532, 534, 537–542
 - biogeography of 514, 541
 - in Centothecoideae + Panicoideae phylogeny 494–496
 - cleistogamy in 286
 - Cymbopogon* inclusion in 542
 - ITS analysis of 534, 538, 541
 - ITS and *trnL*-F analysis of, combined 537, 540–542
 - morphological analysis of 494, 541
 - ndhF* analysis of 494–496
 - New World 504–505, 514
 - in Panicoideae phylogeny 504–505, 514
 - photosynthetic pathways in 514
 - PHYB* analysis of 541
 - rpl16* analysis of 494, 496
 - spikelets of 286
 - trnL*-F analysis of 534, 539, 541
- Andropogon* sect. *Gymnanthelia* 532, 542
- Andropogon* subgen. *Cymbopogon* 532, 542
- Andropogon* subgen. *Heterozygi* 532
- Andropogon* subgen. *Isozygi* 532
- Andropogoneae
 - awned versus awnless 262
 - base chromosome number of 504
 - biogeography of 509, 511, 514–516
 - in Centothecoideae + Panicoideae phylogeny 487, 491–500
 - chloroplast DNA analysis of 530–544
 - core 262
 - core clade of 542
 - economic and ecological importance of 530
 - inflorescence of 530–531
 - ITS analysis of 262, 530–544
 - ITS and *trnL*-F analysis of, combined 537, 540–543
 - morphological analysis of 504
 - ndhF* analysis of 262, 490–500, 530–544
 - New World 504–506, 509, 511, 514–516, 530
 - Old World 530
 - in Panicoideae phylogeny 262, 503–505, 514–516
 - photosynthetic pathway of 262, 504, 514–516, 530–531
 - PHYB* analysis of 531, 537–543
 - phylogeny of 262, 530–544
 - plastid genome analysis of 262
 - in Poaceae phylogeny 245, 252–254
 - rames of 262
 - rbcL* analysis of 533
 - relationship with *Arundinella* 262, 531, 533, 537–542
 - relationship with Arundinelleae 537–541
 - relationship with Paniceae 262
 - rpl16* analysis of 490–500
 - spikelets of 261–262, 504, 530–531, 543
 - tb1* analysis of 531, 541–543
 - trnL* analysis of 530–544
 - trnL*-F analysis of 262, 530–544
- Andropogoninae
 - in Andropogoneae phylogeny 262, 531–532, 538–540, 542
 - biogeography of 514–515
 - ITS analysis of 538
 - ITS and *trnL*-F analysis of, combined 540
 - New World 504, 506, 514–515
 - in Panicoideae phylogeny 503–504, 514–515
 - photosynthetic pathways in 514–515
 - trnL*-F analysis of 539
- Andropogonodae, in Andropogoneae phylogeny 531
- Andropogonoideae, in Panicoideae phylogeny 503
- Androtrichum*, photosynthetic pathway in 102, 111
- Anemanthele*
 - hilum of 355
 - ITS analysis of 355–357
 - relationship with *Achnatherum* 355
 - relationship with *Austrostipa* 355, 358
 - stamen of 355
 - in Stipeae phylogeny 350, 355–357
- Animal pollination, of Hypolytreae 77
- Aniselytron*, tribal/subtribal placement of 423
- Anisopogon*
 - ITS analysis of 351, 355–356
 - matK* analysis of 342
 - ndhF* analysis of 342
 - ndhH* analysis of 342
 - in Poaceae phylogeny 239, 242–244, 342
 - in Pooideae phylogeny 345–346
 - rbcL* analysis of 342
 - in Stipeae phylogeny 350–351, 355–356
- Anomochloa*
 - matK* analysis of 342
 - ndhF* analysis of 240, 242, 248–249, 341–342, 344
 - ndhH* analysis of 341–342
 - in Poaceae phylogeny 239–244, 248–249, 251, 254, 341–342, 344
 - rbcL* analysis of 342
- Anomochloae
 - in Anomochlooideae phylogeny 303

- in Poaceae phylogeny 252–253
- Anomochlooideae
matK analysis of 342
ndhF analysis of 242, 248–249, 342, 344
ndhH analysis of 342
 New World, classification and biogeography of 304–305, 311
 phylogeny of 303
 in Poaceae phylogeny 238–243, 248–255, 260, 303–304, 341–342, 344
rbcL analysis of 342
- Anthaenantia*
 biogeography of 509, 521
 New World 505, 508–509, 521
 in Panicoideae phylogeny 505, 508, 521
 photosynthetic pathway of 521
- Anthaenantiopsis*
 biogeography of 509, 511, 521–522
 New World 505, 508–509, 511, 521–522
 in Panicoideae phylogeny 505, 508, 521–522
 photosynthetic pathway of 521–522
 relationship with *Paspalum* and *Thrasya* 547–548, 550–551, 553
- Antheophora*
 New World 505, 507
 in Panicoideae phylogeny 503, 505, 507
 in Poaceae phylogeny 254
- Anthephoreae, in Panicoideae phylogeny 503
- Anthephorinae, in Panicoideae phylogeny 503
- Anthistiriinae
 in Andropogoneae phylogeny 262, 532, 538–542
 biogeography of 515
 ITS analysis of 538
 ITS and *trnL*–F analysis of, combined 540
 New World 504–506
 in Panicoideae phylogeny 504–505, 515
 photosynthetic pathways in 515
trnL–F analysis of 539
- Anthochloa*
 in *Poa* phylogeny 421, 429–431
 tribal/subtribal placement of 423
- Anthoxanthum*
 geographic distribution of 472
Hierochloë inclusion in 472
 morphological/anatomical analysis of species 472–484
 as outgroup in *Poa* study 422
 in Pooideae phylogeny 343, 346
- Anthoxanthum alpinum*
 geographic distribution of 472
 glume length of 474–475, 480
 height of 474–475
 leaf anatomy of 473–480
 morphological characters of 473–474, 482–484
 relationship with *Anthoxanthum odoratum* 472–484
 spikelets of 474–477, 480
- Anthoxanthum odoratum*
 diploids of 472–484
 geographic distribution of 472
 glume length of 474–475, 480
 height of 474–475
 leaf anatomy of 473–480
 morphological characters of 473–474, 482–484
 relationship with *Anthoxanthum alpinum* 472–484
 spikelets of 474–477, 480
- Apluda*
 in Andropogoneae phylogeny 530, 532, 534, 538–540
 inflorescence of 530
 ITS analysis of 534, 538
 ITS and *trnL*–F analysis of, combined 540
 New World 504–505
 in Panicoideae phylogeny 504–505
trnL–F analysis of 534, 539
- Apoclada*
 biogeography of 309
 in Guaduinae (Bambuseae) phylogeny 309
- Araeococcus*
 in Bromeliaceae phylogeny 36
trnL and *trnL*–F analysis of, combined 36
- Aratitiopea*
 geographic distribution of 227
 inflorescence of 227–233
 in Xyridaceae phylogeny 227, 232
- Aratitiopea lopezii*
 androecium of 230
 calyx of 228–230
 corolla of 228
 gynoecium of 230–232
 inclusion in *Navia* 227
 inflorescence architecture and floral morphology of 227–233
 perianth of 228–230
 in Xyridaceae phylogeny 227
- Archambault, Annie 420–434
- Arctagrostis*
 in Poae phylogeny 421
 in Pooideae phylogeny 343, 346
 relationship with *Poa* 421, 424, 427, 431, 433
 tribal/subtribal placement of 423
- Arctophila*
 relationship with *Poa* 421, 427, 431
 tribal/subtribal placement of 423
- Arctopoa*
 in *Poa* phylogeny 421, 429, 431
 tribal/subtribal placement of 423
- Argentina
 Chloridoideae in 583, 585–588
 Panicoideae of 510–511
- Aristida*
 ITS analysis of 567, 571
 ITS and *trnL*–F analysis of, combined 573
 morphological analysis of 493–494

- ndhF* analysis of 493–496
- as outgroup in Chloridoideae study 567
- in Poaceae phylogeny 242–244
- rpl16* analysis of 493–494, 496
- trnL*-F analysis of 567, 570
- Aristideae, in Poaceae phylogeny 252–253
- Aristidoideae
 - leaf anatomy of 261
 - matK* analysis of 342
 - morphological analysis of 494, 497
 - ndhF* analysis of 342, 490, 494–497
 - ndhH* analysis of 342
 - in Poaceae phylogeny 237, 242–243, 250–254, 260–261, 335, 341–342
 - rbcL* analysis of 342
 - rpl16* analysis of 490, 494, 496–497
 - synapomorphy of 261
- Arriaga, Mirta 349–361
- Arthraxon*
 - in Andropogoneae phylogeny 532, 534, 538–542
 - ITS analysis of 534, 538
 - ITS and *trnL*-F analysis of, combined 540
 - morphological analysis of 541
 - New World 504–505
 - in Panicoideae phylogeny 504–505
 - relationship with *Arundinella* 541
 - trnL*-F analysis of 534, 539
- Arthroxoninae, in Andropogoneae phylogeny 262
- Arthropogon*
 - biogeography of 509, 511, 517
 - New World 505, 507, 509, 511, 517
 - in Panicoideae phylogeny 503, 505, 507, 517
 - photosynthetic pathway of 517
- Arthropogoninae
 - biogeography of 510–511, 516–517
 - New World 504–507, 510–511, 516–517
 - in Panicoideae phylogeny 504–505, 507, 516–517
 - photosynthetic pathway of 507, 516–517
 - spikelets of 507
- Arthrotyliidae
 - buds of 308
 - in Cyperaceae phylogeny 81, 149
 - ITS analysis of 156
 - photosynthetic pathway of 161
 - relationship with Abildgaardieae 81, 149–162
 - relationship with Schoeneae 160
- Arthrotyliidiinae
 - in Andes 311
 - in Bambuseae phylogeny 308–309, 315–316, 323–330
 - biogeography of 308–309, 311
 - branching of 308
 - culms of 308–309
 - in eastern Brazil 311
 - fruit of 308
 - in Guayana Highlands, Venezuela 311
 - internode lengths of 308
 - leaf blade of 308
 - morphological analysis of 323, 326, 329
 - ndhF* analysis of 308, 329
 - phylogeny of 308
 - relationship with Chusqueinae 316
 - relationship with Guaduinae 308–309
 - rhizomes of 308
 - rpl16* analysis of 308, 327, 329
 - rpl16* and morphological analysis of, combined 328
 - spikelets of 308
 - synflorescence of 308
- Arthrotylidium*
 - in Andes 311
 - in Arthrotyliidiinae phylogeny 308–309
 - in Bambuseae phylogeny 308–309, 317, 326–328
 - biogeography of 308–309, 311
 - branching of 308
 - buds of 308
 - internode lengths of 308
 - morphological analysis of 324, 326
 - rpl16* analysis of 327
 - rpl16* and morphological analysis of, combined 328
- Arthrotylis*
 - in Cyperaceae phylogeny 79, 81, 149
 - geographic distribution of 149
 - ITS analysis of 156
 - photosynthetic pathway in 102, 111
 - rbcL* analysis of 79
 - trnL*-F analysis of 81, 157–158
 - trnL*-F and ITS analysis of, combined 159
- Arundinaria*
 - in Arundinariinae phylogeny 307
 - in Bambuseae phylogeny 307, 326–328
 - biogeography of 307
 - morphological analysis of 324, 326, 494
 - ndhF* analysis of 494–496
 - in Poaceae phylogeny 237–239, 242–246
 - rpl16* analysis of 322, 327, 494, 496
 - rpl16* and morphological analysis of, combined 328
- Arundinariinae
 - in Bambuseae phylogeny 306–307, 315–317, 326–328
 - biogeography of 307
 - morphological analysis of 326
 - phylogeny of 307
 - rhizomes of 307
 - rpl16* analysis of 327
 - rpl16* and morphological analysis of, combined 328
- Arundineae
 - in Poaceae phylogeny 252–253
 - Thysanolaena* placement in 509
- Arundinella*
 - in Centothecoideae + Panicoideae phylogeny 487, 494–497
 - ITS analysis of 534, 538, 541
 - ITS and *trnL*-F analysis of, combined 540–541
 - morphological analysis of 494

- ndhF* analysis of 494–496
 New World 504–505
 in Panicoideae phylogeny 504–505
 relationship with Andropogoneae 262, 531, 533–534, 538–540, 542
rpl16 analysis of 494, 496
 spikelets of 262
trnL–F analysis of 534, 539, 541
- Arundinelleae**
 biogeography of 516
 in Centothecoideae + Panicoideae phylogeny 487, 491–500
 geographic distribution of 504
ndhF analysis of 490–500
 New World 504–506, 516
 in Panicoideae phylogeny 503–506, 516
 photosynthetic pathway of 504, 516
 in Poaceae phylogeny 252–254
 relationship with Andropogoneae 537–541
rpl16 analysis of 490–500
Thysanolaena placement in 509
- Arundinoideae**
 in Centothecoideae + Panicoideae phylogeny 499
 Eriachneae placement in 503
matK analysis of 342
 morphological analysis of 494, 497
ndhF analysis of 342, 490, 497
ndhH analysis of 342, 494–496
 in Panicoideae phylogeny 504
 in Poaceae phylogeny 237, 241–243, 248, 250–254, 260, 335, 342
rbcL analysis of 342
 relationship with Centothecoideae 488
rpl16 analysis of 490, 494, 496–497
 Steyermarkochloae placement in 503, 508
- Arundo**
 culms of 325
 ITS analysis of 567, 571
 ITS and *trnL–F* analysis of, combined 573
matK analysis of 342
 morphological analysis of 494
ndhF analysis of 342, 494–496
ndhH analysis of 342
 as outgroup in Chloridoideae study 567
 in Poaceae phylogeny 242–244, 342
rbcL analysis of 342
 relationship with *Steyermarkochloa* 508
rpl16 analysis of 494, 496
trnL–F analysis of 567, 570
- Arundoclaytonia**
 biogeography of 509–510, 529
 in Centothecoideae + Panicoideae phylogeny 494–498, 500
 morphological analysis of 494
ndhF analysis of 493–498, 508
 New World 505, 508–510, 529
- in Panicoideae phylogeny 505, 508–509, 529
 photosynthetic pathway of 529
rpl16 analysis of 494, 496, 498
- Ascolepis**
 absence of 5.8S rDNA insertion in 87, 94
 in Cyperaceae phylogeny 80
 photosynthetic pathway in 102, 111
rbcL analysis of 80
- Asterochaete**, photosynthetic pathway in 102, 111
- Athroostachys**
 branching of 308
 buds of 308
- Atlantic Province, Panicoideae of 511
- atpB**
 in Bromeliaceae analysis 19
 in Poaceae analysis 248
 in *Rhodocoma* analysis 213–214, 217–218
- Atractantha**
 in Arthrostylidiinae phylogeny 308
 in Bambuseae phylogeny 308, 326–328
 biogeography of 308
 branching of 308
 buds of 308
 florets of 308
 morphological analysis of 324, 326
rpl16 analysis of 327
rpl16 and morphological analysis of, combined 328
 spikelets of 308
- Aulonemia**
 in Andes 311
 in Arthrostylidiinae phylogeny 308–309
 in Bambuseae phylogeny 308–309, 326–328
 biogeography of 308–309, 311
 branching of 308
 buds of 308
 internode lengths of 308
 morphological analysis of 324, 326
rpl16 analysis of 322, 327
rpl16 and morphological analysis of, combined 328
- Austral blackbird pollination, of *Puya* 13
- Australia, Chloridoideae in 585
- “Australopoa” group, of *Poa* 421, 430, 432
- Australopyrum**
 chloroplast DNA analysis of 366, 368
DMC1 analysis of 366–367, 370
EF-G analysis of 366–367, 370
 mitochondrial genome of 366
 morphological analysis of 365
 total evidence (combined) analysis of 367–369
 in Triticeae phylogeny 362, 365–370
 W genome transfer to *Elymus* 372
- Austrochloa**, in Chloridoideae phylogeny 572
- Austrodanthonia**, in Danthonioideae phylogeny 261
- Austrofestuca**
 ITS analysis of 414–415
 in *Poa* phylogeny 421, 429–431

- relationship with *Festuca* 407–408
tribal/subtribal placement of 423
- Austrostipa*
geographic distribution of 355
ITS analysis of 351, 354–359
relationship with *Anemathelle* 355, 358
relationship with *Hesperostipa* 358
in Stipeae phylogeny 350–351, 354–360
subgeneric classification of 359
- Austrostipa* subgen. *Arbuscula* 352, 359
Austrostipa subgen. *Aulax* 352
Austrostipa subgen. *Austrostipa* 352, 359
Austrostipa subgen. *Bambusina* 352, 359
Austrostipa subgen. *Ceres* 352
Austrostipa subgen. *Eremophilae* 352
Austrostipa subgen. *Falcateae* 353, 359
Austrostipa subgen. *Lancea* 353
Austrostipa subgen. *Lanterna* 353
Austrostipa subgen. *Lobatae* 353, 359
Austrostipa subgen. *Longiaristatae* 353, 359
Austrostipa subgen. *Petaurista* 353
Austrostipa subgen. *Tuberculatae* 353, 359
- Avena*
ndhF analysis of 341
in Poaceae phylogeny 242–244, 341
in Pooideae phylogeny 343, 346
- Aveneae
as outgroup in Loliinae study 382
in Poaceae phylogeny 251–253, 255–256, 421
in Pooideae phylogeny 335, 345–347, 380–381
- Avenella*, in Pooideae phylogeny 343
- Aveninae
as outgroup in *Poa* study 422
in Pooideae phylogeny 346
- Avenula*, in Pooideae phylogeny 347
- Axonopus*
biogeography of 509–511
New World 504–506, 508–511
in Panicoideae phylogeny 504–505, 508
photosynthetic pathway of 511
- Ayensua*
in Bromeliaceae phylogeny 3–4, 7–9, 14–16, 28, 34, 39
in Bromelioideae phylogeny 34
geographic origin and distribution of 9–10
leaf and trichome anatomy of 13–14
ndhF analysis of 3–4, 7–9, 14–16
pollination of 13
trnL, *trnL-F*, and *trnT-F* analysis of, combined 34
trnL analysis of 28
- Bacterial artificial chromosomes (BACs) 64, 68
- Bagnall, Neil 349–361
- Bambusa*
in Bambuseae phylogeny 326–329
matK analysis of 342
morphological analysis of 324, 326
ndhF analysis of 342
ndhH analysis of 342
Old World taxa, in New World 309
in Poaceae phylogeny 238–239, 242–245, 341–342
rbcL analysis of 342
rpl16 analysis of 322, 327
rpl16 and morphological analysis of, combined 328
- Bambuseae
Arundinariinae + Shibataeinae clade of 306–307
in Bambusoideae phylogeny 303, 306–307, 315
branch patterns of 315, 320
branching of 321–322, 325
buds of 320–322, 325
culms of 306, 315, 320, 325
Euchusquea clade of 329
flowers of 306, 315, 321, 325
foliage leaves of 315, 320, 325
foliar anatomy of 321
foliar micromorphology of 321
fruit of 321
GBSSI analysis of 330
geographic distribution of 315
glumes of 320
Guaduiniae + Arthrotylidiinae clade of 308–309, 311, 315–316, 323–325, 329–330
in Guayana Highlands, Venezuela 311
Laurasian distribution of 307
lemmas of 320
life cycle of 320
monophyly of 306, 315, 323–325
morphological analysis of 317, 320–326
morphology of 306
ndhF analysis of 315–316, 323–325, 329–330
Neotropical clade of 307, 315, 323
New World, classification and biogeography of 306–310
north temperate clade of 306–307, 309, 315–317, 323, 329–330
Old World 306
Old World taxa, in New World 309–310
one-flowered, determinate genera of 315–332
origins of 307
Paleotropical clade of 307, 315, 323, 329–330
phylogeny of 306–310, 315–332
plastid genome analysis of 316
in Poaceae phylogeny 245–246, 251–254
rbcL analysis of 330
rhizomes of 306, 315, 320
rpl16 analysis of 315–316, 319–330
rpl16 and morphological analysis of, combined 328
spikelets of 306, 320–321, 325
synflorescences of 320
trnL-F analysis of 329–330
- Bambusinae
in Bambuseae phylogeny 316, 323, 326–328, 330
morphological analysis of 326

- Old World taxa, in New World 309
 relationship with Hickelinae 316
rpl16 analysis of 327, 330
rpl16 and morphological analysis of, combined 328
- Bambusoideae**
 in Amazonia 311
 in Andes 311
 arm cells of 306, 315
 Bambuseae one-flowered, determinate genera 315–332
 Centothecae placement in 503
 core 303
 in eastern Brazil 311
 foliage leaves of 306
 fusoid cells of 303, 306
 in Guayana Highlands, Venezuela 311
 leaf blades of 303
matK analysis of 342
 in Mesoamerican mountains 311
 monophyly of 315
 morphology of 303, 306
ndhF analysis of 242, 248–249, 342, 490
ndhH analysis of 342
 New World, classification and biogeography of 306–311
 papillae of 306
 phylogeny of 303, 306–311, 323
 in Poaceae phylogeny 237, 239, 242–243, 245, 248–253, 255, 260, 303–304, 315, 335, 342, 345
 pseudospikelets of 303
rbcL analysis of 342
 relationship with Ehrhartoideae 305, 307
 relationship with Steyermarkochloaeae 508
rpl16 analysis of 490
 synflorescences of 303
- BAPO clade, of *Poa* 421, 427–429, 432
- Barbacenia**
 as outgroup in Bromeliaceae (Bromelioideae) study 33
trnL, *trnL*-F, and *trnT*-F analysis of, combined 34
- Barbed goatgrass. *See Aegilops triuncialis*
- Barkworth, Mary 349–361
- Bat pollination, of Bromeliaceae 13
- Baumea**
 in Cyperaceae phylogeny 78
 photosynthetic pathway in 102, 111
rbcL analysis of 78
- Bayer, Randall 349–361
- Bealia**
 biogeography of 593
 in Chloridoideae phylogeny 566, 570–571, 573–575, 581, 584
 ITS analysis of 566, 571
 ITS and *trnL*-F analysis of, combined 573
 morphological analysis of 575, 584–585
 New World 581, 584, 593
trnL-F analysis of 566, 570
- Beckmannia**
 in Poeae phylogeny 421
 in Pooideae phylogeny 343, 346
 relationship with *Poa* 421, 427, 433
 tribal/subtribal placement of 423
- Becquerelia**
 in Cyperaceae phylogeny 78
 5.8S rDNA insertion in 88, 94
 photosynthetic pathway in 102, 111
rbcL analysis of 78
- Bee pollination, of Bromeliaceae 13
- Bell, Hester L. 565–579
- Bellardiochloa**
 in Poeae phylogeny 380, 421
 in Pooideae phylogeny 343, 346
 relationship with *Poa* 421, 424, 427, 431, 433
 tribal/subtribal placement of 423
- BEP clade, of Poaceae 237, 239, 241, 246, 248, 255, 260, 263–264, 303–304, 335, 341–342, 345, 487
- Berry, Paul E. 3–26, 193–203
- Billbergia**
 in Bromeliaceae phylogeny 34, 36
trnL, *trnL*-F, and *trnT*-F analysis of, combined 34
trnL and *trnL*-F analysis of, combined 36
- Bird pollination, of Bromeliaceae 5, 11–13, 21–22
- Bisboeckelera**, photosynthetic pathway in 102, 111
- Bisboeckelereae**
 in Cyperaceae phylogeny 73, 77–78
 5.8S rDNA insertion in 88, 94
 monophyly of 73
rbcL analysis of 73, 77–78
 relationship with Sclerieae 77
- Blackbird pollination, of *Puya* 13
- “Bladder sedges” 179–180, 186–187
- Blepharidachne**
 biogeography of 593
 in Chloridoideae phylogeny 566, 570–571, 573, 581
 ITS analysis of 566, 571
 ITS and *trnL*-F analysis of, combined 573
 New World 581, 593
trnL-F analysis of 566, 570
- Blepharoneuron**
 biogeography of 593
 in Chloridoideae phylogeny 566, 570–571, 573–575, 581, 584
 ITS analysis of 566, 571
 ITS and *trnL*-F analysis of, combined 573
 morphological analysis of 575, 584–585
 New World 581, 584, 593
trnL-F analysis of 566, 570
- Bluegrass. *See Poa*
- Blasmus**
 in Cyperaceae phylogeny 79
 photosynthetic pathway in 102, 111
rbcL analysis of 79
- Boivinella**, in Panicoideae phylogeny 503

- Boivinelleae, in Panicoideae phylogeny 503
- Boivinellinae, in Panicoideae phylogeny 503
- Bolboschoenus*
- in Cyperaceae phylogeny 79, 81
 - 5.8S rDNA insertion in 88, 92, 94, 96
 - ITS analysis of 156
 - photosynthetic pathway in 102, 112
 - rbcL* analysis of 79, 81
 - trnL*-F analysis of 157–158
 - trnL*-F and ITS analysis of, combined 159
- Bothriochloa*
- amphitropical disjuncts in 511
 - in Andropogoneae phylogeny 262, 532, 534, 537–542
 - biogeography of 515
 - ITS analysis of 534, 538, 541
 - ITS and *trnL*-F analysis of, combined 537, 540–542
 - morphological analysis of 541
 - ndhF* analysis of 541
 - New World 504–505, 515
 - in Panicoideae phylogeny 504–505, 515
 - photosynthetic pathways in 515
 - PHYB* analysis of 541
 - tb1* analysis of 541
 - trnL*-F analysis of 534, 539, 541
- Bouchenak-Khelladi, Yanis 248–258
- Bouteloua*
- biogeography of 585, 587–589, 593
 - in Chloridoideae phylogeny 566, 568, 570–571, 573, 575, 577–578, 580–581, 583–584
 - dicliny (breeding systems) in 605–614
 - inflorescence of 606
 - ITS analysis of 566, 568, 571, 577, 606
 - ITS and *trnL*-F analysis of, combined 573
 - morphological analysis of 575, 584
 - New World 581, 583–585, 587–589, 593
 - spikelets of 606–613
 - trnL* analysis of 606
 - trnL*-F analysis of 566, 570, 577, 606
- Boutelouinae
- biogeography of 587, 590, 593
 - in Chloridoideae phylogeny 575, 578, 581–584
 - morphological analysis of 584
 - New World 581–584, 587, 590, 593
- Brachiaria*
- New World 507
 - in Panicoideae phylogeny 507
- Brachyelytrae
- in Poaceae phylogeny 252–253
 - in Pooideae phylogeny 345
- Brachyelytrum*
- ITS analysis of 351, 355–356
 - matK* analysis of 342
 - ndhF* analysis of 342
 - ndhH* analysis of 342
 - in Poaceae phylogeny 242–244, 342
 - in Pooideae phylogeny 341
 - rbcL* analysis of 342
 - relationship with Stipeae 350–351, 355–356
- Brachypodieae
- as outgroup in Loliinae study 382
 - in Pooideae phylogeny 345–346
- Brachypodium*
- molecular analysis of 390–392, 414–415
 - morphological analysis of 387, 394
 - morphological-molecular analysis of, combined 397
 - ndhF* analysis of 341, 343–344
 - as outgroup in *Festuca* study 407, 414–415
 - as outgroup in Loliinae study 382, 387, 390–392, 394, 397
 - in Poaceae phylogeny 242–244, 341
 - in Pooideae phylogeny 343
- Bract(s). *See also specific taxa*
- of Juncaceae 55, 60
- Brazil
- Chloridoideae in 583, 585–588
 - Panicoideae in 509–511
- Brazilian Region, Panicoideae of 511
- Brazilian Shield, Bromeliaceae origin and distribution in 5, 7, 10–12
- Breeding systems
- in *Bouteloua* 605–614
 - evolution of dioecy 605–614
 - phylogenetic study of 606
- Brewcaria*
- in Bromeliaceae phylogeny 4, 7–9, 14–18, 28
 - CAM photosynthesis by 11
 - epiphytism of 12
 - geographic origin and distribution of 10–11
 - leaf and trichome anatomy of 14
 - matK-rps16* analysis of, combined 28
 - ndhF* analysis of 4, 7–9, 14–18
 - pollination of 13
- Briza*, in Pooideae phylogeny 343, 346
- Brizinae, in Poaceae (Pooideae) phylogeny 380
- Brocchinia*. *See also Sequencia*
- adaptive radiation of 7
 - in Bromeliaceae phylogeny 3–4, 7–9, 14–16, 21, 28, 34, 39
 - CAM photosynthesis by 11
 - epiphytism of 12, 19
 - geographic origin and distribution of 9–10, 18–19
 - growth form of 15, 22
 - leaf and trichome anatomy of 14, 20
 - matK-rps16* analysis of, combined 28
 - ndhF* analysis of 3–4, 7–9, 14–16, 21
 - nutrient capture in 22
 - pollination of 13
 - trnL*, *trnL*-F, and *trnT*-F analysis of, combined 34
 - trnL* analysis of 28
- Brocchinia serrata*. *See Sequencia serrata*
- Brocchinia uaipanensis*** 15
- in Bromeliaceae phylogeny 15

- growth form of 15
- morphology of 15
- ndhF* analysis of 15
- Brochnioideae** 15
 - in Bromeliaceae phylogeny 9, 15–16
 - CAM photosynthesis by 11
 - epiphytism of 12
 - geographic origin and distribution of 10
 - leaf and trichome anatomy of 14
 - morphology of 15
 - ndhF* analysis of 9, 15–16
 - pollination of 13, 21
- Bromeae**
 - Bromus* position in 450
 - in Poaceae phylogeny 252–253
 - in Pooideae phylogeny 345
 - relationship with Triticeae 450
- Bromelia*
 - in Bromeliaceae phylogeny 8–9, 36
 - CAM photosynthesis by 11
 - epiphytism of 12
 - geographic origin and distribution of 10, 12
 - leaf and trichome anatomy of 14
 - ndhF* analysis of 8–9
 - pollination of 13
 - trnL* and *trnL-F* analysis of, combined 36
- Bromeliaceae**
 - adaptive radiation of 5, 12–14, 19–22
 - AFLP analyses of 29–33, 35–39, 41
 - atmospheric 20–21, 44
 - atpB* analysis of 19
 - CAM photosynthesis by 4–5, 11–12, 16, 19–20, 28, 41
 - carnivory among 21
 - cladogenetic events within 4–5, 9
 - concerted convergence in 14, 16, 22
 - dispersion and range size of 21
 - 18S rDNA analysis of 19
 - drought adaptations among 5, 12, 19–21
 - ecological types of 20–21, 39
 - epiphytism among 4–5, 12, 19–21, 28
 - fossil record of 5
 - fruit of 3, 8, 12, 21
 - generic delimitation in 27
 - growth forms of 22
 - historical biogeography of 5, 9–12, 18–19
 - hypodermal sclerenchyma among 13–14
 - internal water-storage tissue among 13–14, 19–21, 39–40
 - ladderlike organization of clades in 7–8
 - leaf traits of 5, 12–14, 19–22, 33, 39–41
 - matK* analysis of 3–4, 15, 28
 - matK-rps16* analysis of, combined 28
 - molecular analysis of 3, 27–28
 - molecular clock analysis of 5, 9–12
 - morphological analysis of 3
 - multigene analysis of, need for 17, 20
 - ndhF* analysis of 3–26, 28
 - nomenclature/nomenclature changes for 15–18, 26
 - phylogeny of 3–9, 14–18, 27–43
 - in Poales phylogeny 4, 19, 27–28
 - pollination of 5, 11–13, 21–22
 - rbcL* analysis of 19, 28
 - rps16* analysis of 4
 - seeds of, winged versus plumose 3, 8, 21
 - subfamilies of 3, 15–18
 - tank habits among 20–21
 - trichome anatomy of 5, 13–14, 19–22, 27, 44–52
 - trnL*, *trnL-F*, and *trnT-F* analysis of, combined 34
 - trnL* analysis of 16, 28, 33–35
 - trnL* and *trnL-F* analysis of, combined 33–36
 - trnL-F* analysis of 29, 33–35
 - trnT-L* analysis of 29, 33–35
 - type I–IV 20–21, 39
 - undifferentiated chlorenchyma among 13–14
 - xeromorphic genera of 7–8
- “Bromeliad revolution” 20
- Bromelioideae**
 - AFLP analyses of 29–33, 35–39, 41
 - ancestral stock of 41
 - in Bromeliaceae phylogeny 3–4, 7–9, 15, 28–29, 34, 35, 40–41
 - CAM photosynthesis by 11–12, 19
 - core group of 29, 35, 41
 - epiphytism of 12, 19–21
 - evolution of 28–29
 - fleshy fruit of 3, 8, 12
 - geographic origin and distribution of 9–10, 12, 18–19, 28–29, 41
 - leaf and trichome anatomy of 13–14, 19–21, 33, 39–41
 - ndhF* analysis of 3–4, 7–9, 15
 - phylogeny of 27–43
 - pitcairnioid ancestor of 18–19
 - pollination of 12–13
 - trnL*, *trnL-F*, and *trnT-F* analysis of, combined 34
 - trnL* analysis of 28–29, 33–35
 - trnL* and *trnL-F* analysis of, combined 33–36
 - trnL-F* analysis of 29, 33–35
 - trnT-L* analysis of 29, 33–35
- Bromuniola*
 - in Centothecoideae + Panicoideae phylogeny 489, 494–498, 500
 - geographic distribution of 489
 - morphological analysis of 494, 498
 - ndhF* analysis of 494–496
 - rpl16* analysis of 493–494, 496
 - spikelets of 489
- Bromus*
 - allopolyploids of 468–471
 - in Bromeae phylogeny 450
 - chloroplast DNA analysis of 366, 450–467
 - clades of 452

- DMCI* analysis of 363, 367, 369–370
EF-G analysis of 365, 367, 369–370
 future directions in study of 463
 hybridization of 450
 incongruity among data partitions 458–460
 mitochondrial genome of 366
 morphological analysis of 365, 450–451
ndhF analysis of 450–467
 as outgroup in *Elymus* study 374
 as outgroup in Triticeae study 363, 365–370
 phylogeny of 450–467
 in Poaceae phylogeny 242–244, 450
 in Pooideae phylogeny 343, 345, 450
 taxonomy and classification of 450–452
 total evidence (combined) analysis of 368–369
trnL analysis of 450–467
Bromus sect. *Boissiera* 452–453, 463
Bromus sect. *Bromopsis* 451–454, 456–459, 461–463
Bromus sect. *Bromus*
 in *Bromus* phylogeny 451–453, 455–461, 463
 relationship with *Bromus* sect. *Genea* 468–471
Bromus sect. *Ceratochloa* 451–453, 455–459, 462–463
Bromus sect. *Genea*
 in *Bromus* phylogeny 451–453, 455–461, 463
 isozyme analysis of 468–471
 morphological analysis of 468
Bromus sect. *Neobromus* 452–453, 455–459, 462–463
Bromus sect. *Nevskiella* 452–453, 463
Bromus sect. *Pnigma* 462
Bromus sect. *Triniusia* 452–453, 455–461, 463
 Bruhl, Jeremy J. 72–83, 149–164
 Bruhl, Jeremy J. and Karen L. Wilson—Towards a comprehensive survey of C₃ and C₄ photosynthetic pathways in Cyperaceae 99–148
 Bruneau, Anne 84–98
 Bud(s), of Arthrostylideae 308
Buergersiochloa
 in Bambuseae phylogeny 326–328
 in Bambusoideae phylogeny 303
 biogeography of 310–311
 fimbriae of 311
 lemma of 311
 morphological analysis of 317, 324, 326
 in Olyreae phylogeny 303, 310–311
 in Poaceae phylogeny 242–245
 relationship with *Ekmanochloa* 310
 rpl16 analysis of 327
 rpl16 and morphological analysis of, combined 328
 stamen number of 317
 unisexuality 311
Buergersiochloinae 311
Bulbostylis
 in Abildgaardieae phylogeny 149–164
 in Cyperaceae phylogeny 79, 81
 5.8S rDNA insertion in 87, 94
 embryo type of 150
 geographic distribution of 149
 ITS analysis of 155–156, 158–162
 morphology of 150
 photosynthetic pathway in 102, 112
 rbcL analysis of 79, 81, 151
 relationship with *Abildgaardia* 150, 161
 relationship with *Crosslandia* 161
 relationship with *Fimbristylis* 150, 161
 trnL–F analysis of 81, 156–158, 161–162
 trnL–F and ITS analysis of, combined 159–162
 C₃ photosynthesis
 in Abildgaardieae 161
 in Arthrostylideae 161
 in Chloridoideae 565
 in Cyperaceae 99–148
 in Paniceae 262
 in Panicoideae 504, 511–512, 514–529
 C₄ photosynthesis
 in Abildgaardieae 161
 in Chloridoideae 565, 580
 in Cyperaceae 99–148
 in Paniceae 262
 in Panicoideae 504, 511–512, 514–529
 in *Paspalum* 545
Calamagrostis, in Pooideae phylogeny 343, 346
Calamovilfa
 biogeography of 589, 594
 in Chloridoideae phylogeny 566, 570–571, 573–576, 581, 583–584
 ITS analysis of 566, 571, 576
 ITS and *trnL*–F analysis of, combined 573
 morphological analysis of 575–576, 583–584
 New World 581, 583–584, 589, 594
 relationship with *Eragrostis* 600
 trnL–F analysis of 566, 570
Calderonella
 biogeography of 509–510, 516
 in Centothecoideae + Panicoideae phylogeny 489, 494–500
 geographic distribution of 489
 morphological analysis of 494, 498–499
 ndhF analysis of 494–496
 New World 505–506, 509–510, 516
 in Panicoideae phylogeny 505–506, 516
 photosynthetic pathways in 516
 rpl16 analysis of 494, 496
 spikelets of 489, 498–499
 California, *Aegilops triuncialis* invasion in 295–300
Calotheca
 in Pooideae phylogeny 343, 346
 tribal/subtribal placement of 423
Calyptrocarya, photosynthetic pathway in 102
 CAM photosynthesis, among Bromeliaceae 4–5, 11–12, 16, 19–20, 28, 41
 Campbell, Lisa M. and Dennis Wm. Stevenson—Inflores-

- cence architecture and floral morphology of
Aratitiopea lopezii (Xyridaceae) 227–233
- Canada, Panicoideae in 509–510
- Canastra*
biogeography of 509, 511, 522
New World 505, 508–509, 511, 522
in Panicoideae phylogeny 505, 508, 522
photosynthetic pathway of 522
- Canistropsis*, in Bromeliaceae phylogeny 29
- Canistrum*
in Bromeliaceae phylogeny 8–9, 29, 34
CAM photosynthesis by 11
epiphytism of 12
geographic origin and distribution of 10
leaf and trichome anatomy of 14
ndhF analysis of 8–9
pollination of 13
trnL, *trnL-F*, and *trnT-F* analysis of, combined 34
trnL and *trnL-F* analysis of, combined 36
- Cape Floristic Region
ecological parameters in 215–216
Rhodocoma from, phylogeny and historical ecology of
213–226
- Capeobolus*
in Cyperaceae phylogeny 72
photosynthetic pathway in 102, 112
- Capillipedium*
in Andropogoneae phylogeny 532, 534, 537–542
ITS analysis of 534, 538, 541
ITS and *trnL-F* analysis of, combined 537, 540–542
morphological analysis of 541
ndhF analysis of 541
PHYB analysis of 541
tb1 analysis of 541
trnL-F analysis of 534, 539, 541
- Capitularina*
5.8S rDNA insertion in 88
photosynthetic pathway in 102, 112
- Carbon dioxide (CO₂) compensation point, in Cyperaceae
99–100
- Carbon isotope ratio ($\delta^{13}\text{C}$ value), in Cyperaceae
99–100
- Carex*
agmatoploidy of 63–64, 193
BAC studies of 68
in Cariceae phylogeny 165–192
chromosome distribution patterns in 65
chromosome evolution in 63–68, 193–203
chromosome numbers of 63, 65–68
chromosome races in, C-value measurement of 63
in Cyperaceae phylogeny 58, 72, 79, 84
in Cyperales phylogeny 68
5.8S rDNA insertion in 85, 87, 92, 94
economic uses of 84
ETS-1f analysis of 179–180
genome size of 66–68
geographic distribution of 165
global versus regional treatment of 165
habitats of 165
holocentric chromosomes of 63–64, 193
ITS, ETS-1f, and *trnL-F* analysis of, combined 165–
192
in New Zealand 418
photosynthesis in 147
photosynthetic pathway in 102, 105, 113–114
polyploidy in 193–194
primitive species of 166
rbcL analysis of 79
relationship with other genera of Cariceae 166
rps16 analysis of 57–58
speciation in 193–194
trnL-F analysis of 57–58, 180
vegetative anatomy of 99
- Carex bicknellii* group 199–202
- Carex brevior* group 199–202
- Carex lupulina*, BAC library of 68
- Carex* subgen. *Carex*
in Cariceae phylogeny 176–179, 183–189
ITS, ETS-1f, and *trnL-F* analysis of, combined 166,
174–179
primitive wetland versus advanced dry land 188
relationship with *Carex* subgen. *Vignea* 188
relationship with *Carex* subgen. *Vigneastra* 181
relationships among sects. *Ceratocystis*, *Spirostachyae*,
Rhynchocystis, and *Rostrales* 187
relationships among sects. *Clandestinae*, *Acrocystis*,
Pictae, and *Lamprochlaenae* 188
relationships among sects. *Laxiflorae*, *Panicaceae*, *Gri-
seae*, and *Careyanae* 187–188
relationships among sects. *Racemosae*, *Scitae*, *Limosae*,
and *Phacocystis* 188
variability between species 174–175
- Carex* subgen. *Carex* sect. *Abditispicae* 177, 179
- Carex* subgen. *Carex* sect. *Acrocystis* 166, 177, 188
- Carex* subgen. *Carex* sect. *Anomalae* 177
- Carex* subgen. *Carex* sect. *Aulocystis* 177, 179, 182
- Carex* subgen. *Carex* sect. *Carex* 177, 179, 186–187
- Carex* subgen. *Carex* sect. *Careyanae* 177, 179, 182, 187–
188
- Carex* subgen. *Carex* sect. *Ceratocystis* 177, 179, 182, 187
- Carex* subgen. *Carex* sect. *Clandestinae* 177, 179, 188
- Carex* subgen. *Carex* sect. *Collinsiae* 177, 179, 182, 186–
187
- Carex* subgen. *Carex* sect. *Depauperatae* 177, 179
- Carex* subgen. *Carex* sect. *Gracillimae* 185
- Carex* subgen. *Carex* sect. *Griseae* 177, 179, 182, 187–188
- Carex* subgen. *Carex* sect. *Hallerianae* 177, 179, 185–186
- Carex* subgen. *Carex* sect. *Hirtae* 186
- Carex* subgen. *Carex* sect. *Hirtifoliae* 177, 179, 185–186
- Carex* subgen. *Carex* sect. *Hymenochlaenae* 166, 177, 179,
182, 185–186
- Carex* subgen. *Carex* sect. *Lamprochlaenae* 177, 179, 188

- Carex* subgen. *Carex* sect. *Laxiflorae* 177, 179, 182, 187–188
- Carex* subgen. *Carex* sect. *Limosae* 177, 179, 188
- Carex* subgen. *Carex* sect. *Longicaules* 177, 179, 185–186
- Carex* subgen. *Carex* sect. *Longirostres* 185
- Carex* subgen. *Carex* sect. *Lupulinae* 68, 177, 179, 186–187
- Carex* subgen. *Carex* sect. *Paludosae* 177, 179, 186–187
- Carex* subgen. *Carex* sect. *Panicaceae* 177, 179, 182, 187–188
- Carex* subgen. *Carex* sect. *Phacocystis* 177, 179, 182, 187–188
- Carex* subgen. *Carex* sect. *Phyllostachyae* 168
- Carex* subgen. *Carex* sect. *Physocarpae* 186
- Carex* subgen. *Carex* sect. *Pictae* 188
- Carex* subgen. *Carex* sect. *Porocystis* 177, 179, 185–186
- Carex* subgen. *Carex* sect. *Pseudocypereae* 186
- Carex* subgen. *Carex* sect. *Racemosae* 177, 179, 187–188
- Carex* subgen. *Carex* sect. *Rhynchocystis* 177, 179, 186–187
- Carex* subgen. *Carex* sect. *Rostrales* 177, 179, 182, 186–187
- Carex* subgen. *Carex* sect. *Scirpinae* 177
- Carex* subgen. *Carex* sect. *Scitae* 188
- Carex* subgen. *Carex* sect. *Shortianae* 177, 179
- Carex* subgen. *Carex* sect. *Spirostachyae* 177, 179, 187
- Carex* subgen. *Carex* sect. *Squarrosae* 177, 179, 186–187
- Carex* subgen. *Carex* sect. *Sylvatica* 185
- Carex* subgen. *Carex* sect. *Triquetra* 185
- Carex* subgen. *Carex* sect. *Tumidae* 186
- Carex* subgen. *Carex* sect. *Vesicariae* 177, 179, 186–187
- Carex* subgen. *Carex* sect. *Virescentes* 185–186
- Carex* subgen. *Carex* sect. *Viridiflorae* 185
- Carex* subgen. *Kreczetoviczia* 188
- Carex* subgen. *Kuekenhalia* 186–187
- Carex* subgen. *Primocarex* 166
in Cariceae phylogeny 184, 186
ITS, ETS-1f, and *trnL*-F analysis of, combined 166
- Carex* subgen. *Psyllophora* 166
in Cariceae phylogeny 183
- Carex* subgen. *Psyllophora* sect. *Aciculares* 169
- Carex* subgen. *Psyllophora* sect. *Dornera* 169
- Carex* subgen. *Psyllophora* sect. *Leptocephalae* 170
- Carex* subgen. *Psyllophora* sect. *Leucoglochin* 187
- Carex* subgen. *Psyllophora* sect. *Obtusatae* 170
- Carex* subgen. *Psyllophora* sect. *Psyllophora* 170
- Carex* subgen. *Psyllophora* sect. *Rupestres* 170
- Carex* subgen. *Vignea*
androgynous versus gynecandrous 176, 184
in Cariceae phylogeny 176, 180–181, 183–184, 188–189
compound inflorescences in 184
ITS, ETS-1f, *trnL*-F, and *trnE*-D analysis of, combined 166–174, 176, 180–181, 189
as possible progenitor 188
relationship with *Carex* subgen. *Carex* 188
variability between species 174
- Carex* subgen. *Vignea* sect. *Chordorrhizae* 181, 184
- Carex* subgen. *Vignea* sect. *Curvulae* 170
- Carex* subgen. *Vignea* sect. *Deweyanae* 177, 180, 184
- Carex* subgen. *Vignea* sect. *Dispermae* 177
- Carex* subgen. *Vignea* sect. *Divisae* 184
- Carex* subgen. *Vignea* sect. *Foetidae* 181
- Carex* subgen. *Vignea* sect. *Glareosae* 176–177, 180, 184
- Carex* subgen. *Vignea* sect. *Heleoglochin* 180, 184
- Carex* subgen. *Vignea* sect. *Macrocephalae* 184
- Carex* subgen. *Vignea* sect. *Multiflorae* 180, 184
- Carex* subgen. *Vignea* sect. *Ovales*
AFLP analysis of 193–203
bicknellii group of 199–202
brevior group of 199–202
in Cariceae phylogeny 176–177, 180, 184
chromosome evolution in 193–203
chromosome patterns in 199–200, 202
eastern North American clades of 194, 199–202
ETS analysis of 194
geographic distribution of 194
ITS analysis of 194
tenera group of 199–201
tribuloides group of 199–201
- Carex* subgen. *Vignea* sect. *Phaestoglochin* 176–177, 180–181, 184
- Carex* subgen. *Vignea* sect. *Physoglochin* 176–177, 180, 183
- Carex* subgen. *Vignea* sect. *Stellulatae* 176–177, 180, 184, 194
- Carex* subgen. *Vignea* sect. *Vulpinae* 177, 180, 184
- Carex* subgen. *Vigneastra*
in Cariceae phylogeny 176–179, 183–185, 188–189
ITS, ETS-1f, and *trnL*-F analysis of, combined 166–173, 176–179
as possible progenitor 185
relationship with *Carex* subgen. *Carex* 181
- Carex* subgen. *Vigneastra* sect. *Baccantes* 177, 179
- Carex* subgen. *Vigneastra* sect. *Indicae* 177, 179, 187
- Carex* subgen. *Vigneastra* sect. *Polystachyae* 177, 179
- Carex tenera* group 199–201
- Carex tribuloides* group 199–201
- Caribbean
Bromeliaceae in 5, 7, 10–12
Chloridoideae in 585–588
Panicoideae in 509–511
- Cariceae
androgynous versus gynecandrous 176, 184, 188–189
“backbone” tree of 189
“bladder sedges” 179–180, 186–187
clade A (Core Unspicate Clade) of 175, 178, 183, 188
clade B (*Schoenoxiphium* Clade) of 175–176, 178, 183, 188
clade C (*Vignea* Clade) of 176, 178, 183–184, 188
clade D (Core *Carex* Clade) of 176–179, 184–189
in Cyperaceae phylogeny 73, 77, 82, 84
in Cyperales phylogeny 68
5.8S rDNA insertion in 85, 87, 94, 96
ETS-1f analysis of 166, 179–180
flowers of 81
geographic patterns in 165, 189

- ITS analysis of 166, 179
 as model system 189
 monophyly of 73
 nested ITS, ETS-1f, and *trnL* analysis of 165–192
 phylogeny of 165–192
rbcL analysis of 73, 77
 relationship with Dulichieae 78–81
 relationship with Scirpeae 77–81
 relationships among major clades in 175–176
trnT-L analysis of 166, 180
 utricle of 81
- Caricoideae
 in Cyperaceae phylogeny 72–73, 79, 82
 5.8S rDNA insertion in 87
 monophyly of 73
rbcL analysis of 72–73, 79
- Caris, Pieter 204–209
- Carnivory, among Bromeliaceae 21
- Carpha*
 in Cyperaceae phylogeny 77–78
 in Cyperales phylogeny 68
 photosynthetic pathway in 102, 114
rbcL analysis of 77–78
- Castellia*
 in Loliinae phylogeny 380–405
 molecular analysis of 389–392
 morphological analysis of 387, 394–395, 399
 morphological-molecular analysis of, combined 397
- Catabrosa*
 in Pooideae phylogeny 343, 346
 relationship with *Poa* 421–422, 427
 tribal/subtribal placement of 423
- Catabrosella*, tribal/subtribal placement of 423
- Catalán, Pilar et al.—A systematic approach to subtribe Loliinae (Poaceae: Pooideae) based on phylogenetic evidence 380–405
- Catapodium*
 morphological analysis of 387, 394
 morphological-molecular analysis of, combined 397
 in Poeae phylogeny 381, 403
 relationship with Loliinae 382, 387, 394, 397
- Cathariostachys*
 in Bambuseae phylogeny 323, 326–328, 330
 morphological analysis of 324, 326, 330
rpl16 analysis of 327
rpl16 and morphological analysis of, combined 328
- Catopsis*
 in Bromeliaceae phylogeny 7–9, 21, 34, 35
 CAM photosynthesis by 11
 geographic origin and distribution of 10–11
 leaf and trichome anatomy of 14
ndhF analysis of 7–9, 21
trnL, *trnL-F*, and *trnT-F* analysis of, combined 34
trnL and *trnL-F* analysis of, combined 36
- Caustis*
 in Cyperaceae phylogeny 78
- 5.8S rDNA insertion in 88, 94
 photosynthetic pathway in 102, 114
rbcL analysis of 78
- Cayouette, Jacques 450–467
- Celtica*, in Stipeae phylogeny 349
- Cenchrinae
 biogeography of 517
 morphological analysis of 507
ndhF analysis of 507
 New World 504–507, 517
 in Panicoideae phylogeny 504–505, 507, 517
 photosynthetic pathway of 517
- Cenchrus*
 biogeography of 510, 517
 New World 505, 507, 510, 517
 in Panicoideae phylogeny 505, 507, 517
 photosynthetic pathway of 517
- Centotheca*
 in Centothecoideae + Panicoideae phylogeny 489, 493–500
 geographic distribution of 489
 morphological analysis of 494
ndhF analysis of 494–496, 498
 in Poaceae phylogeny 251, 254
rpl16 analysis of 493–494, 496
 spikelets of 489
- Centothecoaceae
 in Bambusoideae phylogeny 503
 biogeography of 510, 516
 in Centothecoideae + Panicoideae phylogeny 488, 491–500
ndhF analysis of 490–500
 New World 504–506, 510, 516
 in Panicoideae phylogeny 504–506, 516
 photosynthetic pathways in 516
 in Poaceae phylogeny 252–253
rpl16 analysis of 490–500
 spikelets of 506
- Centothecoideae
 geographic distribution of 488–489
 leaf anatomy of 488
 morphological analysis of 488, 491–492, 494
ndhF analysis of 344, 487–502
 phylogeny of 487–491, 499
 in Poaceae phylogeny 241, 248, 250–254, 260, 335, 341, 344, 487
 relationship with Arundinoideae 488
 relationship with Panicoideae 487–504
rpl16 analysis of 487–502
 spikelets of 488–489, 491–492
- Centothecoideae + Panicoideae clade 487–502
- Central America
 Bromeliaceae origin and distribution in 5, 7, 10–12, 18–19
 Chloridoideae in 585–588
 Panicoideae of 510–511

- Centrochloa*
 biogeography of 509, 511, 522
 New World 505, 508–509, 511, 522
 in Panicoideae phylogeny 505, 508, 522
 photosynthetic pathway of 522
- Centromere(s)
 characterization of 64
 of Cyperales 62–64
 diffuse, in agmatoploidy 63–64
- Centropodia*, in Poaceae phylogeny 242–244
- Cephalocarpus*, photosynthetic pathway in 102, 114
- Cephalostachyum*
 in Bambuseae phylogeny 326–328
 morphological analysis of 324, 326
rpl16 analysis of 327
rpl16 and morphological analysis of, combined 328
- Cephalostemon* 6
- Ceratophyllum*, as outgroup in Bromeliaceae study 4
- Cerros-Tlatilpa, Rosa 530–544, 565–579
- Chaboissaea*
 biogeography of 587, 593
 in Chloridoideae phylogeny 566, 570–571, 573, 575, 577, 581, 584
 ITS analysis of 566, 571
 ITS and *trnL*-F analysis of, combined 573
 morphological analysis of 575, 584–585
 New World 581, 584, 587, 593
trnL-F analysis of 566, 570
- Chacoan Province, Panicoideae of 511–512
- Chaetium*
 biogeography of 509, 518
 New World 505, 507, 509, 518
 in Panicoideae phylogeny 505, 507, 518
 photosynthetic pathway of 518
- Chascolytrum*
 in Pooideae phylogeny 343, 346
 tribal/subtribal placement of 423
- Chase, Mark W. 55–61, 72–83, 248–258
- Chasmanthium*
 biogeography of 509, 516
 in Centothecoideae + Panicoideae phylogeny 489, 494–500
 geographic distribution of 489
 ITS analysis of 567, 571
 ITS and *trnL*-F analysis of, combined 573
matK analysis of 342
 morphological analysis of 494, 498
ndhF analysis of 342, 344, 494–498
ndhH analysis of 342
 New World 505–506, 509, 516
 as outgroup in Chloridoideae study 567
 in Panicoideae phylogeny 505–506, 516
 photosynthetic pathway of 516
 in Poaceae phylogeny 241–244, 342, 344
rbcL analysis of 342
rpl16 analysis of 494, 496, 498
 spikelets of 489
trnL-F analysis of 567, 570
- Chasmanthium* clade 497–498
- Chasmogamy
 versus cleistogamy 287
 cleistogamy ratio to 287–293
 cost-benefit analysis of 287
 density and 288
 in grasses (Poaceae) 286–294
 light and 289–290
 phenology of 290–291
 soil fertility and 288–289
 soil moisture and 289–290
- Cheplick, Gregory P.—Plasticity of chasmogamous and cleistogamous reproductive allocation in grasses 286–294
- Chevaliera*
 in Bromeliaceae phylogeny 34, 36, 41
trnL, *trnL*-F, and *trnT*-F analysis of, combined 34
trnL and *trnL*-F analysis of, combined 36
- Chevalierella*
 in Centothecoideae + Panicoideae phylogeny 497
 geographic distribution of 489
 spikelets of 489
- Chile, Panicoideae of 510
- Chionachne*
 in Andropogoneae phylogeny 262, 533–534, 538, 543
 ITS analysis of 534, 538, 543
ndhF analysis of 543
- Chionachninae
 in Andropogoneae phylogeny 262, 533, 538–540, 543
 ITS analysis of 538
 ITS and *trnL*-F analysis of, combined 540
 morphological analysis of 543
trnL-F analysis of 539
- Chionochoa*, relationship with *Festuca* 416
- Chionographis*, holocentric chromosomes in 63
- Chlorideae, in Chloridoideae phylogeny 565, 581
- Chloridinae
 biogeography of 586–587, 593
 in Chloridoideae phylogeny 576, 581–584
 morphological analysis of 584
 New World 581–584, 586–587, 593
- Chloridoideae
 amphitropical disjuncts in 586–588
 base chromosome number of 580
 biogeography of 565, 580–594
Bouteloua dicliny 605–614
 Caribbean species of 588
 chloroplast DNA analysis of 565–579
 clade A of 575–576
 clade B of 576
 clade C of 576–578
Eragrostis monophyly and infrageneric classification 595–604
 evolutionary history of 582, 589–590

- fossil record of 589–590
 fruits of 580–581
GBSSI analysis of 572–574, 576
 habitat preferences of 580
 inflorescence of 565, 581
 ITS analysis of 565–579, 582
 ITS and *trnL*-*F* analysis of, combined 572–573, 582
 lodicules of 580–581
matK analysis of 342, 565, 572–577, 582
 morphological analysis of 494, 497, 575, 580–581, 583–585
ndhF analysis of 342, 490, 494–497, 574
ndhF and *rbcL* analysis of, combined 574
ndhH analysis of 342
 New World 565–567, 580–594
 North American center of 586–588
 northwestern South American center of 586–588
 photosynthetic pathways in 565, 580
 phylogeny of 565–594
 in Poaceae phylogeny 237, 242–243, 248, 250–254, 260, 335, 342, 565
rbcL analysis of 342, 574
rp16 analysis of 490, 494, 496–497, 572–574
rpoC2 analysis of 574
rps16 analysis of 572–574, 576
 southern South American center of 586–588
 spikelets of 565, 581, 583–585
 synapomorphies of 580
trnL analysis of 565–579
trnL-*F* analysis of 565–579, 582
 unplaced species of 588
 world centers of 585
- Chloris*
 biogeography of 585, 587–588, 593
 breeding system of 608–610
 in Chloridoideae phylogeny 566, 570–573, 575, 577, 580–581, 583–584
 ITS analysis of 566, 571–572, 575, 577
 ITS and *trnL*-*F* analysis of, combined 572–573, 577
 morphological analysis of 575, 577, 584
 New World 581, 583–585, 587–588, 593
 as outgroup in Triodieae study 263
trnL-*F* analysis of 566, 570, 572, 577
- Chlorocharis*, photosynthetic pathway of 104–105
 “Chlorocyperacean” 104
 Chlorophyll fluorescence, of Tillandsioideae 45
 Chloroplast DNA analysis. *See also specific genes and taxa*
 of Abildgaardieae 151–162
 of Andropogoneae 530–544
 of Bromeliaceae 3–4
 of *Bromus* 450–467
 of Cariceae 165–192
 of Chloridoideae 565–579
 of *Elymus* 373–377
 of *Poa* 420–433
 of Triticeae 363, 365–367
- Chorizandra*
 in Cyperaceae phylogeny 78
 5.8S rDNA insertion in 88
 photosynthetic pathway in 102, 114
rbcL analysis of 78
rps16 analysis of 57–58
trnL-*F* analysis of 57–58
- Chromosome evolution
 in Cyperales 62–71
 and diversification 62
 Heilborn’s hypothesis on 194, 198, 202
 and reproductive isolation 62
 in *Rhynchospora* 193
 and speciation 62
- Chromosome fission/fusion 62–63
 in *Carex* 193
 in Cyperales 66–67
- Chromosome number
 of *Carex* sect. *Ovales* 193–203
 of Centothecoideae + Panicoideae clade 491–492
 of Chloridoideae 580
 of Cyperales 63, 65–68
 in Heilborn’s hypothesis 194
 of Isachneae 506
 of Paniceae 262, 507
 of Panicoideae 503–504
 of Thysanolaeneae 509
- Chrysitricheae
 in Cyperaceae phylogeny 73–78, 80
 5.8S rDNA insertion in, absence/presence of 88, 94, 96
rbcL analysis of 73–78, 80
- Chrysitrix*
 in Cyperaceae phylogeny 78
 5.8S rDNA insertion in 88, 94
 photosynthetic pathway in 102, 114
rbcL analysis of 78
- Chrysopogon*
 in Andropogoneae phylogeny 532, 534, 537–542
 biogeography of 515
 ITS analysis of 534, 538, 541–542
 ITS and *trnL*-*F* analysis of, combined 537, 540–542
 morphological analysis of 542
ndhF analysis of 542
 New World 504–505, 515
 in Panicoideae phylogeny 504–505, 515
 photosynthetic pathway of 515
PHYB analysis of 542
trnL-*F* analysis of 534, 539, 541–542
- Chusquea*
 altitudinal range of 307
 in Andes 311
 in Bambuseae phylogeny 307–308, 316–317, 319–323, 326–329
 base chromosome number of 307, 329
 biogeography of 307, 311
 buds/branching of 317, 325, 329

- culms of 307, 316, 329
 in eastern Brazil 311
 latitudinal range of 307
 lemma of 329
matK analysis of 342
 monophyly of 329
 morphological analysis of 317, 324–326, 329, 493–494
 morphology of 316
ndhF analysis of 342, 493–496
ndhH analysis of 342
 papillae of 307
 in Poaceae phylogeny 239, 242–245, 342
rbcL analysis of 342
 relationship with *Neurolepis* 307–308
rpl16 analysis of 319–322, 327, 493–494, 496
rpl16 and morphological analysis of, combined 328
 spikelets of 307, 316
- Chusquea* subgen. *Chusquea* 307–308, 329
 morphology of 316
- Chusquea* subgen. *Rettbergia* 307–308, 323, 325, 329
 morphology of 316
- Chusquea* subgen. *Swallenochloa* 307–308, 329
- Chusqueinae
 in Andes 311
 in Bambuseae phylogeny 306–308, 315–316, 323, 326–330
 biogeography of 307–308, 311
 buds/branching of 316
 in eastern Brazil 311
 glumes of 316
 lemmas of 316
 morphological analysis of 323, 326, 329
 morphology of 316
ndhF analysis of 316
 papillae of 307
 phylogeny of 307–308
 plastid genome analysis of 316
 relationship with Arthrostylidiinae 316
 relationship with Guaduinae 316
rpl16 analysis of 316, 327, 329
rpl16 and morphological analysis of, combined 328–329
 spikelets of 307, 316
 synflorescences of 316
trnL–F analysis of 316
- Cinninae, in Poaceae (Pooideae) phylogeny 380
- Cladium*
 in Cyperaceae phylogeny 77–78
 5.8S rDNA insertion in 88, 94
 genome size of 66
 photosynthetic pathway in 102, 114
rbcL analysis of 77–78
rps16 analysis of 57
trnL–F analysis of 57
- Cladoraphis*
 in Chloridoideae phylogeny 581, 583
- morphological analysis of 583
 New World 581, 583
 relationship with *Eragrostis* 595–597, 599–600, 602
- Clark, Lynn G. 237–247, 303–314, 487–502
- Clark, Lynn G. et al.—Phylogenetic relationships among the one-flowered, determinate genera of Bambusoideae (Poaceae: Bambusoideae) 315–332
- Clayton, Derek 267–270
- Cleistachne*, in Andropogoneae phylogeny 542
- Cleistogamy
 bet-hedging strategy in 287
 versus chasmogamy 287
 chasmogamy ratio to 287–293
 cost-benefit analysis of 287
 definition of 286
 density and 288
 environmental factors in 287
 in *Eremitis* 310
 in grasses (Poaceae) 286–294
 light and 289–290
 multiple strategies in 286
 phenology of 290–291
 reproductive assurance in 287
 reproductive dimorphism in 286
 soil fertility and 288–289
 soil moisture and 289–290
 theoretical considerations in 287
- Coatinga Province, Panicoideae of 511
- Coelachne*, in Panicoideae phylogeny 503
- Coelachyrum*, in Chloridoideae phylogeny 572–574
- Coelorachis*
 in Andropogoneae phylogeny 532, 534, 538–540, 542
 ITS analysis of 534, 538, 542
 ITS and *trnL–F* analysis of, combined 540
trnL–F analysis of 534, 539
- Coicinae
 in Andropogoneae phylogeny 262, 533, 538–540, 543
 ITS analysis of 538
 ITS and *trnL–F* analysis of, combined 540
 morphological analysis of 543
 New World 504–506
 in Panicoideae phylogeny 504–505
trnL–F analysis of 539
- Coix*
 in Andropogoneae phylogeny 530, 533–534, 538–540, 542–543
 ITS analysis of 534, 538
 ITS and *trnL–F* analysis of, combined 540
 New World 504–505
 in Panicoideae phylogeny 504–505
 spikelets of 530
trnL–F analysis of 534, 539
- Colantheria*
 branching of 308
 buds of 308
 in Poaceae phylogeny 251

- Coleanthinae, in Poaceae (Pooideae) phylogeny 380
Coleanthus, tribal/subtribal placement of 423
Coleochloa
 in Cyperaceae phylogeny 78
 5.8S rDNA insertion in 94
 photosynthetic pathway in 102, 114
 rbcL analysis of 78
Colpodium, tribal/subtribal placement of 423
 Columbus, J. Travis 530–544, 580–594, 605–614
 Columbus, J. Travis et al.—Phylogenetics of Chloridoideae (Gramineae): A preliminary study based on nuclear ribosomal internal transcribed spacer and chloroplast *trnL*–F sequences 565–579
 Concerted convergence
 in Bromeliaceae 14, 16, 22
 in Liliales 22
Connellia
 in Bromeliaceae phylogeny 4, 16, 39–40
 ndhF analysis of 4, 16
 Cope, Tom 267–270
Cornucopia, tribal/subtribal placement of 423
Costularia, photosynthetic pathway in 102, 115
Cottea
 amphitropical disjunct in 583
 biogeography of 588, 594
 in Chloridoideae phylogeny 566, 570–571, 573–575, 581, 583
 ITS analysis of 566, 571
 ITS and *trnL*–F analysis of, combined 573
 morphological analysis of 583
 New World 581, 583, 588, 594
 in Poaceae phylogeny 251
 trnL–F analysis of 566, 570
 Cotteinae
 biogeography of 588, 594
 in Chloridoideae phylogeny 575–576, 581–583
 geographic origin of 583
 morphological analysis of 583
 New World 581–583, 588, 594
Cottendorfia
 in Bromeliaceae phylogeny 4, 7–9, 14–18, 28, 39–40
 CAM photosynthesis by 11
 epiphytism of 12
 geographic origin and distribution of 10–11, 18
 leaf and trichome anatomy of 14
 matK-rps16 analysis of, combined 28
 ndhF analysis of 4, 7–9, 14–18
 pollination of 13
Courtoisina
 in Cyperaceae phylogeny 80
 photosynthetic pathway in 102, 115
 rbcL analysis of 80
coxII, in Triticeae analysis 363–366, 369
Craspedorhachis, in Poaceae phylogeny 251, 254
Critesion, in Triticeae phylogeny 369–370
Crithopsis
 chloroplast DNA analysis of 366
 DMC1 analysis of 367
 EF-G analysis of 367
 mitochondrial genome of 366
 morphological analysis of 365–366
 total evidence (combined) analysis of 367–369
 in Triticeae phylogeny 365–369
Crosslandia
 in Abildgaardieae phylogeny 149–164
 embryo type of 150
 geographic distribution of 149
 ITS analysis of 155–156, 158–160
 morphology of 150
 photosynthetic pathway in 102, 115
 relationship with *Abildgaardia* 161
 relationship with *Bulbostylis* 161
 relationship with *Fimbristylis* 161–162
 trnL–F analysis of 155–158
 trnL–F and ITS analysis of, combined 159–162
Crypsis
 in Chloridoideae phylogeny 566, 570–571, 573, 575–576, 581, 584
 ITS analysis of 566, 571, 576
 ITS and *trnL*–F analysis of, combined 573
 morphological analysis of 575–576, 584
 New World 581, 584
 trnL–F analysis of 566, 570
 Cryptangieae
 in Cyperaceae phylogeny 73, 77
 5.8S rDNA insertion in 88
 monophyly of 73
 rbcL analysis of 73, 77–78
Cryptanthus
 in Bromeliaceae phylogeny 4, 29, 36
 ndhF analysis of 4
 trnL and *trnL*–F analysis of, combined 36
Cryptochloa
 florets of 311
 in Olyrinae (Olyreae) phylogeny 311
 synflorescence of 310
Ctenium
 biogeography of 587, 593
 in Chloridoideae phylogeny 566, 570–571, 573, 580–581
 ITS analysis of 566, 571
 ITS and *trnL*–F analysis of, combined 573
 New World 581, 587, 593
 trnL–F analysis of 566, 570
Ctenopsis
 in Loliinae phylogeny 380–405
 molecular analysis of 386, 390–392, 414–415
 morphological analysis of 389, 394–395
 morphological-molecular analysis of, combined 397
 relationship with *Festuca* 406–407
 Cuba, Panicoideae in 509–510

Culm(s)

- of Arthrostylidiinae 308–309
- of Bambuseae 306, 315, 320, 325
- of Centothecoideae + Panicoideae clade 491–492
- of *Diandrollyra* 311
- of Olyreae 310, 315
- of Poaceae 261, 286

Cutandia

- in Pooideae phylogeny 343, 346, 380–381
- relationship with *Elymus* 377
- relationship with *Festuca* 406–407
- tribal/subtribal placement of 423

Cyathochaeta, photosynthetic pathway in 102, 115*Cyathocoma*, photosynthetic pathway in 102, 115*Cymbopogon*

- in Andropogoneae phylogeny 530, 532, 534, 537–540, 542
- ITS analysis of 534, 538
- ITS and *trnL*-F analysis of, combined 537, 540, 542
- morphological analysis of 542
- New World 504–505
- in Panicoideae phylogeny 504–505
- trnL*-F analysis of 534, 539

Cymophyllus

- in Cariceae phylogeny 175, 177–179, 183
- ITS, ETS-1f, and *trnL*-F analysis of, combined 166, 175, 177–178
- ITS analysis of 179

Cynodon

- biogeography of 587, 589, 593
- breeding system of 608–610
- in Chloridoideae phylogeny 566, 568, 570–573, 577, 580–581, 584
- ITS analysis of 566, 568, 571–572
- ITS and *trnL*-F analysis of, combined 573
- morphological analysis of 584
- New World 581, 584, 587, 589, 593
- trnL*-F analysis of 566, 570, 572

Cynodonteae

- biogeography of 590, 593
- in Chloridoideae phylogeny 565, 575–578, 581–585
- morphological analysis of 584–585
- New World 581–585, 590, 593
- in Poaceae phylogeny 250, 252–255

Cynosureae 401

Cynosurinae

- molecular analysis of 389–392
- morphological analysis of 387
- in Poeae phylogeny 381, 403, 421
- in Pooideae phylogeny 346
- relationship with Loliinae 381–382, 387, 389–392, 401

Cynosurus

- molecular analysis of 390–392
- morphological analysis of 387, 393–394
- morphological-molecular analysis of, combined 397
- in Poeae phylogeny 380, 403

in Pooideae phylogeny 343, 346

relationship with Loliinae 382, 387, 389–394, 397, 401

Cyperaceae

- carbon isotope ratio ($\delta^{13}\text{C}$ value) in 99–100
- chromosome evolution in *Carex* sect. *Ovales* 193–203
- CO₂ compensation point in 99–100
- CO₂-use efficiency in 105
- in Cyperales phylogeny 55
- ecological importance of 84
- economic importance of 72, 84
- ethnobotanical importance of 72
- 5.8S rDNA insertion in 84–98
- flowers of 56
- genomic resources in 65
- geographic distribution of 84
- helophytic habit in 105
- holocentric chromosomes in 63
- ITS analysis of 81
- Kranz anatomy of 99
- maximum cells distant count in 99–104
- NADP-ME versus NAD-ME type 103
- nuclear ribosomal DNA analysis of 81, 84–98
- one cell distant criterion in 100–105
- Oxychloe* placement in 56
- photosynthesis in 104, 147–148
- photosynthetic carbon reduction (PCR) cells in 100
- photosynthetic pathways (C₃ and C₄) in 99–148
- phylogeny of 55–61, 72–83
- phylogeny of Abildgaardieae 149–164
- phylogeny of Cariceae 165–192
- plastid genome analysis of 55–61, 72–83
- in Poales phylogeny 55
- pollination of 77
- primary carbon assimilation (PCA) cells in 100–103
- pseudomonads of 56, 77
- radiate versus non-radiate chlorenchyma in 99, 101
- rbcL* analysis of 72–85
- rbcL* and *trnL*-F analysis of, combined 81
- relationship with Bromeliaceae 27–28
- relationship with Juncaceae 55–61, 72, 85
- relationship with Thurniaceae 56
- rps16* analysis of 56–60, 81
- Schoenus* inflorescence unit (spikelet) 204–209
- silica bodies of 56
- subfamily delimitation in 82
- trnL*-F analysis of 56–60, 81
- 26S rRNA analysis of 93
- vegetative anatomy of 99, 104
- water-use efficiency in 105

Cyperales

- agmatoploidy of 63–64, 66–67
- BAC libraries in 64, 68
- centromeres and telomeres of 62–64
- chromosome evolution in 62–71
- chromosome fission/fusion in 66–67
- chromosome numbers of 63, 65–68

- genome size of 65–68
 genomic resources on 64, 68
 graphical mapping of genome size versus chromosome number among 66–67
 holocentric chromosomes of 63–64
 Juncaceae placement in 55
 pairwise comparisons of chromosome number/genome size among 66–68
 phylogeny of 68
 polyploidy in 67
 relationship with Poales 64
rps16 analysis of 56–60
trnL-F analysis of 56–60
- Cypereae**
 absence of 5.8S rDNA insertion in 87, 94, 96
 C₄ photosynthesis in 99
 in Cyperaceae phylogeny 73, 80
rbcL analysis of 73, 80
- Cyperochloa**
 in Centothecoideae + Panicoideae phylogeny 489, 494–500
 morphological analysis of 494
ndhF analysis of 494–496, 498
 in Poaceae phylogeny 254
rpl16 analysis of 493–494, 496
- Cyeroideae**
 C₄ photosynthesis in 99
 in Cyperaceae phylogeny 72–73, 78–80, 82
 5.8S rDNA insertion in 87
 inflorescence unit of 204, 208
rbcL analysis of 72–73, 78–80
- Cyperus**
 BAC studies of 68
 in Cyperaceae phylogeny 58, 73, 80–81, 84
 in Cyperales phylogeny 68
 5.8S rDNA insertion in, absence/presence of 86, 92, 94
 economic uses of 84
 genome size of 66
 kinetochore binding sites of 63
 photosynthesis in 147–148
 photosynthetic pathways in (C₃ and C₄) 101–102, 115–125
rbcL analysis of 73, 80–81
rps16 analysis of 58
trnL-F analysis of 58
 vegetative anatomy of 99
- Cyperus rotundus**
 genomic resources in 64–65
 as starch/food source 64–65
- Cyphochlaena**, in Panicoideae phylogeny 503
- Cypringlea**
 in Cyperaceae phylogeny 72, 81
 photosynthetic pathway in 102, 125
rbcL analysis of 81
- Cyrtococcum**
 New World 505, 508
- in Panicoideae phylogeny 505, 508
- Dactylidinae**
 molecular analysis of 389–392
 morphological analysis of 387
 in Poeae phylogeny 380, 403, 421
 in Pooideae phylogeny 346
 relationship with Loliinae 380, 382, 387, 401
 relationship with *Poa* 421–422
- Dactylis**
 molecular analysis of 390–392, 414–415
 morphological analysis of 387, 393–394
 morphological-molecular analysis of, combined 397
 in Poeae phylogeny 403
 in Pooideae phylogeny 343
 relationship with *Festuca* 406–408
 relationship with Loliinae 382, 387, 390–394, 397, 401
 relationship with *Poa* 421–422, 424, 427
 tribal/subtribal placement of 423
- Dactyloctenium**
 in Chloridoideae phylogeny 566, 570–571, 573, 580–581
 ITS analysis of 566, 571
 ITS and *trnL-F* analysis of, combined 573
 New World 581
 relationship with *Eragrostis* 600
trnL-F analysis of 566, 570
- Danthonia**
 cleistogamy in 286–287
matK analysis of 342
 morphological analysis of 494
ndhF analysis of 342, 494–496
ndhH analysis of 342
 in Poaceae phylogeny 242–244, 342
rbcL analysis of 342
rpl16 analysis of 493–494, 496
- Danthonidium**, in Poaceae phylogeny 251
- Danthonieae**, in Poaceae phylogeny 252–253
- Danthonioideae** 261–262
 African species of 261
 Australasian species of 259, 261
 haustorial synergids of 261
matK analysis of 342
 morphological analysis of 494, 497
ndhF analysis of 342, 490, 494–497
ndhH analysis of 342
 phylogeny of 259, 261–262
 in Poaceae phylogeny 237, 242–243, 250–254, 260–262, 335, 342
rbcL analysis of 342
rpl16 analysis of 490, 494, 496–497
 synapomorphy of 261
- Danthoniopsis**
 in Centothecoideae + Panicoideae phylogeny 487, 489, 494–500
 morphological analysis of 494
ndhF analysis of 494–498

- in Poaceae phylogeny 237, 241–244
rbcL analysis of 240, 242, 245
 relationship with Andropogoneae 537–541
rpl16 analysis of 494, 496–498
- Danthoniopsis* clade 497–499
- Dasyochloa*
 biogeography of 593
 in Chloridoideae phylogeny 566, 570–571, 573–574, 577, 581, 585
 ITS analysis of 566, 571
 ITS and *trnL*-F analysis of, combined 573
 morphological analysis of 585
 New World 581, 585, 593
trnL-F analysis of 566, 570
- Dasypoa*, tribal/subtribal placement of 423
- Dasypyrum*
 chloroplast DNA analysis of 366, 368
DMCI analysis of 367, 370
EF-G analysis of 367, 370
 genome transfer to *Elymus* 374–378
 mitochondrial genome of 366
 morphological analysis of 365
 total evidence (combined) analysis of 367–369
 in Triticeae phylogeny 365–370
- Davidse, Gerrit 503–529
- Davis, Jerrold I 237–247
- Davis, Jerrold I and Robert J. Soreng—A preliminary phylogenetic analysis of the grass subfamily Pooideae (Poaceae), with attention to structural features in the plastid and nuclear genomes, including an intron loss in GBSSI 335–348
- Decaryochloa*
 in Bambuseae phylogeny 323, 326–328, 330
 morphological analysis of 324, 326, 330
rpl16 analysis of 327
rpl16 and morphological analysis of, combined 328
- Decumbentes group, of *Paspalum* 545–562
- Deinacanthon*
 in Bromeliaceae phylogeny 34–36
trnL, *trnL*-F, and *trnT*-F analysis of, combined 34
trnL and *trnL*-F analysis of, combined 35–36
- DELTA system, of grasses (Poaceae) 267–268
- Dendrocalamus*, in Bambuseae phylogeny 329
- Denham, Silvia S. and Fernando O. Zuloaga—Phylogenetic relationships of the Decumbentes group of *Paspalum*, *Thrasya*, and *Thrasyopsis* (Poaceae: Panicoideae: Paniceae) 545–562
- Deschampsia*
 in Pooideae phylogeny 343, 346–347
 relationship with *Poa* 421–422, 424, 427
- Desmazeria*
 in Pooideae phylogeny 343, 346, 381
 tribal/subtribal placement of 423
- Desmoschoenus*
 in Cyperaceae phylogeny 73, 80–82
 photosynthetic pathway in 102, 125
- rbcL* analysis of 80
- Deuterocohnia*
 in Bromeliaceae phylogeny 4, 7–9, 14–17, 28, 34–36
 CAM photosynthesis by 11
 epiphytism of 12
 geographic origin and distribution of 11–12
 leaf and trichome anatomy of 13–14
matK-rps16 analysis of, combined 28
 narrow endemism in 21
ndhF analysis of 4, 7–9, 14–17
 pollination of 13, 21
 species diversity of 21
trnL, *trnL*-F, and *trnT*-F analysis of, combined 34
trnL and *trnL*-F analysis of, combined 36
- Dhooge, Sandra 72–83
- Diandrochloa*
 in Chloridoideae phylogeny 576
 relationship with *Eragrostis* 595–596, 599, 602
- Diandrolyra*
 biogeography of 310
 culm of 311
 in Olyrinae (Olyreae) phylogeny 311
 stamen number of 311
 synflorescence of 311
- Diarrhena*
 ITS analysis of 351–356
matK analysis of 342
ndhF analysis of 342
ndhH analysis of 342
 in Poaceae phylogeny 242–244, 342
rbcL analysis of 342
 in Stipeae phylogeny 350–356
- Diarrheneae
 in Poaceae phylogeny 252–253
 in Pooideae phylogeny 345–346
- Dichantherium*
 biogeography of 511, 519
 cleistogamy and chasmogamy in 287–293
 New World 505, 508, 511, 519
 in Panicoideae phylogeny 505, 508, 519
 photosynthetic pathway of 511, 519
- Dichanthium*
 in Andropogoneae phylogeny 532, 534, 537–542
 ITS analysis of 534, 538, 541
 ITS and *trnL*-F analysis of, combined 537, 540–542
 morphological analysis of 541
ndhF analysis of 541
 New World 504–505
 in Panicoideae phylogeny 504–505
PHYB analysis of 541
trnL-F analysis of 534, 539, 541
- Dicliny
 in *Bouteloua* 605–614
 definition of 605
 partial 605
- Didymiandrum*, photosynthetic pathway in 102, 125

- Didymogonyx*, internode lengths of 308
- Diectomis*
 in Andropogoneae phylogeny 532, 535, 537–541
 ITS analysis of 535, 538, 541
 ITS and *trnL*–*F* analysis of, combined 537, 540–542
 morphological analysis of 541
trnL–*F* analysis of 535, 539, 541
- Dielsiochloa*, in Pooideae phylogeny 343, 346
- Digitaria*
 amphitropical disjuncts in 511
 biogeography of 511, 517–518
 ITS analysis of 536, 538
 New World 504–507, 511, 517–518
 as outgroup in Andropogoneae study 533, 536
 in Panicoideae phylogeny 504–505, 507, 517–518
 photosynthetic pathway of 517–518
 in Poaceae phylogeny 251, 256
 relationship with Andropogoneae 538–540
trnL–*F* analysis of 536
- Digitariinae
 biogeography of 517–518
 New World 504–507, 517–518
 in Panicoideae phylogeny 504–505, 507, 517–518
 photosynthetic pathway of 517–518
- Dimeria*
 in Andropogoneae phylogeny 530
 spikelets of 530
- Dimerieae, in Andropogoneae phylogeny 532
- Dimeriinae
 in Andropogoneae phylogeny 262, 532
 in Panicoideae phylogeny 503
- Dimorphostachys*, relationship with *Paspalum* 546
- Dinebra*
 in Chloridoideae phylogeny 566, 570–571, 573, 577, 581
 ITS analysis of 566, 571
 ITS and *trnL*–*F* analysis of, combined 573
 New World 581
trnL–*F* analysis of 566, 570
- Dioecy, evolution of 605–614
- Diplacrum*
 in Cyperaceae phylogeny 78
 5.8S rDNA insertion in, absence/presence of 88, 94
 photosynthetic pathway in 102, 125
rbcL analysis of 78
- Diplasia*, photosynthetic pathway in 102, 125
- Dissanthelium*
 in *Poa* phylogeny 421, 429–432
 tribal/subtribal placement of 423
- Distichia*
 5.8S rDNA insertion in 89
 flower of 55, 60
 geographic distribution of 55
 growth form of 55
 inflorescence of 55
 in Juncaceae phylogeny 55, 58–60
 relationship with *Oxychloe* 90
rps16 analysis of 57, 59
trnL–*F* analysis of 57, 59
- Distichlis*
 biogeography of 589, 593
 breeding system of 608
 in Chloridoideae phylogeny 566, 570–571, 573–574, 577, 581, 584
 ITS analysis of 566, 571, 577
 ITS and *trnL*–*F* analysis of, combined 573
 morphological analysis of 584
ndhF and *rbcL* analysis of, combined 574
 New World 581, 584, 589, 593
 in Poaceae phylogeny 242–244
trnL–*F* analysis of 566, 570, 577
- Diversification. *See also specific taxa*
 chromosome evolution and 62
- DMC1*, in Triticeae analysis 363, 365–367, 369–370
- 5.8S rDNA insertion, in Cyperaceae 84–98
- 18S rDNA
 in Bromeliaceae analysis 19
 in Poaceae analysis 252
- 26S rDNA
 in Cyperaceae analysis 93
 in *Oryza* analysis 93
- Dormancy, seed
 in *Aegilops triuncialis* 296–298
 invasive species 296
- Doyle, Jeff J. 595–604
- Draffin, Suzanne J. 406–419
- Dransfield, Soejatmi 315–332
- Drought tolerance, among Bromeliaceae 5
- Drymochloa*, in Loliinae phylogeny 381, 400
- Dulichieae
 in Cyperaceae phylogeny 73, 82
 monophyly of 73
rbcL analysis of 73
 relationship with Cariceae 78–81
- Dulichium*
 in Cyperaceae phylogeny 79
 genome size of 66
 photosynthetic pathway in 102, 125
rbcL analysis of 79
- Dupontia*
 in Poeae phylogeny 421
 in Pooideae phylogeny 343
 relationship with *Poa* 421, 427, 431, 433
 tribal/subtribal placement of 423
- Duthiea*
matK analysis of 342
ndhF analysis of 342
ndhH analysis of 342
 in Poaceae phylogeny 342
 in Pooideae phylogeny 345–346
rbcL analysis of 342

- Duvall, Melvin R. et al.—Phylogeny of the grasses (Poaceae) revisited 237–247
- Dyckia*
 in Bromeliaceae phylogeny 3–4, 7–9, 14–17, 35–36
 CAM photosynthesis by 11
 epiphytism of 12
 geographic origin and distribution of 10–12, 18–19
 leaf and trichome anatomy of 13–14
matK-rps16 analysis of, combined 28
ndhF analysis of 3–4, 7–9, 14–17
 pollination of 13
trnL and *trnL-F* analysis of, combined 36
- Dyckieae 16, 35
- Dyer, Andrew R.—Phenotypic plasticity may facilitate invasion by *Aegilops triuncialis* 295–300
- Ecdeiocolea*
matK analysis of 342
ndhF analysis of 341–342
ndhH analysis of 341–342
 as outgroup in Poaceae study 238–246
rbcL analysis of 342
 relationship with Poaceae 238
- Ecdeiocoleaceae
ndhH analysis of 341–344
 plastid genome analysis of 336–337
 relationship with Joinvilleaceae 237–238
 relationship with Poaceae 237–238
- Echinochloa*
 biogeography of 519–520
 New World 505, 507, 519–520
 in Panicoideae phylogeny 505, 507, 519–520
 photosynthetic pathway of 519–520
- Echinolaena*
 biogeography of 509, 522
 New World 505, 508–509, 522
 in Panicoideae phylogeny 505, 508, 522
 photosynthetic pathway of 522
- Echinopogon*, in Pooideae phylogeny 343, 346
- Ectrosia*
 in Chloridoideae phylogeny 566, 570–571, 573, 576
 ITS analysis of 566, 571
 ITS and *trnL-F* analysis of, combined 573
 relationship with *Eragrostis* 600–601
trnL-F analysis of 566, 570
- Ecuador, Chloridoideae in 586–587
- Edmuntoa*
 in Bromeliaceae phylogeny 29, 36
trnL and *trnL-F* analysis of, combined 36
- EF-G*, in Triticeae analysis 363–365, 369–370
- Eglaria*, photosynthetic pathway in 102, 125
- Ehrharta*
matK analysis of 342
ndhF analysis of 342, 344
ndhH analysis of 342
 in Poaceae phylogeny 241–244, 342, 344
rbcL analysis of 342
- Ehrharteae
 biogeography of 306
 in Ehrhartoideae phylogeny 303, 306
 in Poaceae phylogeny 252–253
 relationship with Oryzeae 306
- Ehrhartoideae
matK analysis of 342
ndhF analysis of 242, 248–249, 342, 490
ndhH analysis of 342
 New World, classification and biogeography of 305–306, 311
 phylogeny of 303, 305–306
 in Poaceae phylogeny 237, 242–243, 245, 250–253, 255, 260, 303–305, 335, 342, 345
rbcL analysis of 342
 relationship with Bambusoideae 305, 307
 relationship with Pooideae 305
rpl16 analysis of 490
- Ekmanochloa*
 biogeography of 310
 leaf blades of 310
 lemmas of 310
 in Olyrinae (Olyreae) phylogeny 310
 relationship with *Buergersiochloa* 310
- Eleocharideae
 in Cyperaceae phylogeny 73, 81
 5.8S rDNA insertion in, absence/presence of 88, 94
 ITS analysis of 81, 156
 monophyly of 73
 as outgroup in Abildgaardieae study 151
rbcL analysis of 73, 81
rbcL and *trnL-F* analysis of, combined 81
- Eleocharis*
 BAC studies of 68
 in Cyperaceae phylogeny 79, 81, 84
 in Cyperales phylogeny 68
 5.8S rDNA insertion in, absence/presence of 86, 88–90, 92, 94–96
 economic uses of 64–65, 84
 genome size of 66
 ITS analysis of 95, 156
 in New Zealand 418
 nuclear ribosomal DNA analysis of 95
 one cell distant criterion in 101–105
 photosynthesis in, conflicting data on 148
 photosynthetic pathways in (C₃ and C₄) 101–102, 104–105, 125–128, 161
 photosynthetic pathways in (NAD-ME) 103, 105
 phylogeny of 95
rbcL analysis of 79
rbcL and *trnL-F* analysis of, combined 81
trnL-F analysis of 157–158
trnL-F and ITS analysis of, combined 159
- Eleusine*
 biogeography of 588, 593

- in Chloridoideae phylogeny 566, 570–571, 573, 575, 577, 580–581, 584
 ITS analysis of 566, 571
 ITS and *trnL*-F analysis of, combined 573
 morphological analysis of 575, 584
 New World 581, 584, 588, 593
 relationship with *Eragrostis* 600
trnL-F analysis of 566, 570
- Eleusininae
 biogeography of 588, 593
 in Chloridoideae phylogeny 575, 581–582, 584
 morphological analysis of 584
 New World 581–582, 584, 588, 593
- Elionurus*
 in Andropogoneae phylogeny 262, 532, 535, 538–540, 542–543
 ITS analysis of 535, 538
 ITS and *trnL*-F analysis of, combined 540, 542
ndhF analysis of 542
 New World 505
 in Panicoideae phylogeny 505
PHYB analysis of 542
tb1 analysis of 542
trnL-F analysis of 535, 539, 542
- Elymandra*
 biogeography of 515
 New World 505, 515
 in Panicoideae phylogeny 505, 515
 photosynthetic pathway of 515
- Elymus*
 allopolyploids of 372–379
 chloroplast DNA analysis of 368, 373–377
GBSSI analysis of 374, 376–378
 H genome of 372–378
 hexaploidy in 372–373, 375–376, 378
 meiotic behavior of 374
 octoploidy of 372, 378
 P genome of 372
 phylogeny of 372–379
 in Pooideae phylogeny 343, 377–378
 relationship with *Lolium* 380
 St genome of 372–378
 tetraploidy of 372–378
 in Triticeae phylogeny 372–378
 W genome of 372
 Y genome of 372, 374, 376–377
- Elytrostachys*
 in Arthrotyliidiinae (Bambuseae) phylogeny 308
 branching of 308
 buds of 308
 spikelets of 308
- Embryo(s)
 of Centothecoideae + Panicoideae clade 491–492
 of Panicoideae 503
 of Poaceae 261
- Encholirium*
 in Bromeliaceae phylogeny 3–4, 7–9, 14–17, 28, 35
 CAM photosynthesis by 11
 epiphytism of 12
 geographic origin and distribution of 10–12, 18–19
 leaf and trichome anatomy of 13–14
matK-rps16 analysis of, combined 28
ndhF analysis of 3–4, 7–9, 14–17
 pollination of 13
- Enneapogon*
 amphitropical disjunct in 583
 biogeography of 588, 594
 in Chloridoideae phylogeny 566, 570–571, 573–575, 581, 583
 ITS analysis of 566, 571
 ITS and *trnL*-F analysis of, combined 573
 morphological analysis of 583
 New World 581, 583, 588, 594
trnL-F analysis of 566, 570
- Enteropogon*
 biogeography of 587, 593
 in Chloridoideae phylogeny 566, 570–571, 573, 577, 581, 584
 ITS analysis of 566, 571
 ITS and *trnL*-F analysis of, combined 573
 morphological analysis of 584
 New World 581, 584, 587, 593
trnL-F analysis of 566, 570
- Entoplocamia*, in Chloridoideae phylogeny 576
- Epidryos*, habitat of 20
- Epiphytism
 of Bromeliaceae 4–5, 12, 19–21, 28
 effect on dispersion and range size 21
 evolution of, relationship with carnivory 21
 of Tillandsioideae 14, 19–21, 50
- Epischoenus*, photosynthetic pathway in 102, 128
- Eragrostideae
 biogeography of 590, 594
 in Chloridoideae phylogeny 565, 574–577, 581–583
 morphological analysis of 583
 New World 581–583, 590, 594
 in Poaceae phylogeny 252–255
- Eragrostidinae
 biogeography of 586, 594
 in Chloridoideae phylogeny 576, 581–583
 morphological analysis of 583
 New World 581–583, 586, 594
- Eragrostiella*
 in Chloridoideae phylogeny 576
 relationship with *Eragrostis* 595–597
- Eragrostis*
 allopolyploidy in 596, 602
 biogeography of 585, 587–588, 594–595
 in Chloridoideae phylogeny 566, 570–576, 581, 583, 595
 economic importance of 595

- GBSSI* analysis of 572–574, 597–602
 ITS analysis of 566, 571
 ITS and *trnL*-F analysis of, combined 573
matK analysis of 596–597, 600
 monophyly and infrageneric classification of 595–604
 morphological analysis of 494, 575–576, 583, 595–596
ndhF analysis of 494–496, 574
ndhF and *rbcL* analysis of, combined 574
 New World 581, 583, 585, 587–588, 594
 as outgroup in Triodieae study 263
 photosynthetic pathway of 565, 595
 in Poaceae phylogeny 242–244, 256, 595
 relationship with *Pappophorum* 572
 relationship with *Poa* 420
rpl16 analysis of 494, 496, 572–574
rps16 analysis of 597–602
 spikelets of 595–596
trnL-F analysis of 566, 570
- Eragrostis* sect. *Eragrostis* 601–602
Eragrostis sect. *Lappula* 601–602
Eragrostis sect. *Platystachya* 601–602
Eragrostis sect. *Psilantha* 601–602
Eragrostis subgen. *Caesia* 602
Eragrostis subgen. *Eragrostis* 602
 Eragrostoideae 580
- Eremitis*
 in Bambusoideae phylogeny 303
 biogeography of 310
 cleistogamy in 310
matK analysis of 342
ndhF analysis of 342
ndhH analysis of 342
 in Olyreae phylogeny 303, 310
 in Parianinae phylogeny 310
 in Poaceae phylogeny 242–245, 342
 pollination of 310
rbcL analysis of 342
 synflorescence of 310
- Eremocaulon*
 biogeography of 309
 in Guaduinae (Bambuseae) phylogeny 309
- Eremochloa*
 New World 504–505
 in Panicoideae phylogeny 504–505
- Eremopoa*
 in *Poa* phylogeny 420–422, 424, 427, 429
 tribal/subtribal placement of 423
- Eremopyrum*
 chloroplast DNA analysis of 366, 368
DMCI analysis of 366–367, 369–370
EF-G analysis of 366–367, 369–370
 mitochondrial genome of 366
 morphological analysis of 365
 relationship with *Elymus* 374, 377
 total evidence (combined) analysis of 367–369
 in Triticeae phylogeny 362, 365–370
- Eriachne*
 geographic distribution of 497
 in Poaceae phylogeny 237–238, 241–245, 260–261, 497, 500, 504
rbcL analysis of 497
rpl16 analysis of 493
- Eriachneae
 in Arundinoideae phylogeny 503
 in Centothecoideae + Panicoideae phylogeny 499
 in Panicoideae phylogeny 503
 in Poaceae phylogeny 252–255
- Erianthus*
 in Andropogoneae phylogeny 532, 535, 538–540
 ITS analysis of 535, 538
 ITS and *trnL*-F analysis of, combined 540
trnL-F analysis of 535, 539
- Eriocaulaceae, relationship with Xyridaceae 227
- Eriochloa*
 biogeography of 518
 New World 505, 507, 518
 in Panicoideae phylogeny 505, 507, 518
 photosynthetic pathway of 511, 518
- Eriochrysis*
 New World 505
 in Panicoideae phylogeny 505
- Erioneuron*
 biogeography of 588, 593
 in Chloridoideae phylogeny 566, 570–571, 573–574, 577, 581, 585
 ITS analysis of 566, 571
 ITS and *trnL*-F analysis of, combined 573
 morphological analysis of 585
 New World 581, 585, 588, 593
trnL-F analysis of 566, 570
- Eriophorum*
 BAC studies of 68
 in Cyperaceae phylogeny 58, 79
 5.8S rDNA insertion in 88, 94, 96
 genome size of 66
 as outgroup in Cariceae study 172–173
 photosynthetic pathway in 102, 128–129
rbcL analysis of 79
rps16 analysis of 57–58
trnL-F analysis of 57–58
- Euandropogoneae, in Andropogoneae phylogeny 532
- Euchlaena*, in Andropogoneae phylogeny 533
- Euclasta*
 in Andropogoneae phylogeny 530, 532, 535, 538–541
 ITS analysis of 535, 538, 541
 morphological analysis of 541
 New World 504–505
 in Panicoideae phylogeny 504–505
 spikelets of 530
- Eulalia*
 in Andropogoneae phylogeny 530, 532, 535, 538–540
 ITS analysis of 535, 538

- ITS and *trnL*-F analysis of, combined 540
 spikelets of 530
trnL-F analysis of 535, 539
- Eustachys*
 biogeography of 587–589, 593
 in Chloridoideae phylogeny 566, 570–574, 577, 580–581, 584
 ITS analysis of 566, 571–572
 ITS and *trnL*-F analysis of, combined 573
 morphological analysis of 584
ndhF analysis of 574
 New World 581, 584, 587–589, 593
trnL-F analysis of 566, 570, 572
- Evandra*
 in Cyperaceae phylogeny 78
 photosynthetic pathway in 102, 129
rbcL analysis of 78
- Everardia*, photosynthetic pathway in 102, 129
- Everett, Joy 349–361
- Exocarya*
 in Cyperaceae phylogeny 58
 photosynthetic pathway in 102, 129
rps16 analysis of 58
trnL-F analysis of 58
- Exochogyne*, photosynthetic pathway in 102, 129
- External transcribed spacer (ETS)
 in *Carex* sect. *Ovales* analysis 194
 in Cariceae analysis 165–192
- Farrago*, in Poaceae phylogeny 251, 254
- Fascicularia*
 AFLP analysis of 36–37
 in Bromeliaceae (Bromelioideae) phylogeny 29, 34–37, 41
 geographic origin and distribution of 41
 leaf blade anatomy of 41
 morphology of 41
trnL, *trnL*-F, and *trnT*-F analysis of, combined 34
trnL and *trnL*-F analysis of, combined 35–36
- Fernseea*
 in Bromeliaceae (Bromelioideae) phylogeny 29, 35–36
trnL and *trnL*-F analysis of, combined 35–36
- Festuca*
 base chromosome number of 413
 broad- versus fine-leaved 389–393, 396–401, 406
 chloroplast DNA analysis of 381
 dispersal of 418
 DNA content of 413, 416
 ecology of 416–418
 FEVRE lineages of 389–392, 396–399, 408, 413–415
 genome size of 413, 416
 geographic distribution of 380
 geographic origins of 406–407, 418
 habitat range of 406
 hilum of 380–381
 ITS analysis of 381, 386, 389–390, 392, 397–398, 406–419
 lemma of 380–381
 in Loliinae phylogeny 380–405
 molecular analysis of 386, 389–392, 406–419
 morphological analysis of 382–385, 387–399
 morphological-molecular analysis of, combined 397–398
 morphology of 380–381, 406
 New Zealand, phylogeny and biogeography of 406–419
 Northern Hemisphere taxa of 406
 phylogeny of 380–382, 400–403, 406–419
 ploidy levels in 413, 418
 in Poaceae phylogeny 256
 in Poaeae phylogeny 380, 424
 in Pooideae phylogeny 343, 346, 380
 relationship with *Austrofestuca* 407
 relationship with *Dactylis* 406–407
 relationship with *Leucopoa* 381
 relationship with *Lolium* 380–381, 406–413
 relationship with *Poa* 407–408, 421–422
 relationship with *Schedonorus* 381, 406–413
 relationship with *Vulpia* 381, 406–407, 413
 southern endemic clade of 414–416, 418
 spikelets of 380–381
 subgeneric classifications of 381
trnL-F analysis of 381, 386, 389, 391, 397–398, 406–419
Festuca subgen. *Drymanthele* 382–383, 389–393, 401–402, 410
Festuca subgen. *Drymanthele* sect. *Phaeochloa* 402
Festuca subgen. *Drymanthele* sect. *Pseudoscariosa* 402
Festuca subgen. *Drymanthele* sect. *Scariosa* 402
Festuca subgen. *Festuca* 382–383, 400–403, 409
Festuca subgen. *Festuca* sect. *Amphigenes* 383, 389–393, 400–403, 410
Festuca subgen. *Festuca* sect. *Aulaxyper* 381, 383, 386, 389–393, 399–402, 406, 409, 413–416
Festuca subgen. *Festuca* sect. *Eskia* 383, 390–393, 401–402, 406, 410
Festuca subgen. *Festuca* sect. *Festuca* 383, 390–393, 400–402, 406, 408–409, 413
Festuca subgen. *Festuca* sect. *Pseudatropis* 383, 410
Festuca subgen. *Festuca* sect. *Pseudoscariosae* 383, 389–392
Festuca subgen. *Festuca* sect. *Scariosae* 383, 389–392, 410
Festuca subgen. *Festuca* sect. *Subbulbosae* 383, 389–393, 400–401, 410
Festuca subgen. *Helleria* 410
Festuca subgen. *Leucopoa* 382–384, 389–393, 400, 402, 410–412
Festuca subgen. *Leucopoa* sect. *Breviaristatae* 383, 403, 411
Festuca subgen. *Leucopoa* sect. *Leucopoa* 383, 410–411
Festuca subgen. *Schedonorus* 380, 382, 384, 389–393, 399, 401–402, 411
Festuca subgen. *Schedonorus* sect. *Plantynia* 384, 402
Festuca subgen. *Schedonorus* sect. *Schedonorus* 384, 402
Festuca subgen. *Schedonorus* sect. *Subulatae* 384

- Festuca* subgen. *Subbulbosae* 402
Festuca subgen. *Subulatae* 382, 389, 393, 401–402, 411
 Festucaceae 194
 Festuceae 401
 relationship with Centothecoideae 488
Festucella, tribal/subtribal placement of 423
 Festucinae, in Poeae (Pooideae) phylogeny 380
Festucopsis
 chloroplast DNA analysis of 366
 DMCI analysis of 366–367
 EF-G analysis of 363, 366–367
 mitochondrial genome of 366
 morphological analysis of 363, 365
 total evidence (combined) analysis of 367–369
 in Triticeae phylogeny 362–363, 365–369
Ficinia
 absence of 5.8S rDNA insertion in 87, 94
 in Cyperaceae phylogeny 58, 73, 80–82
 photosynthetic pathway in 102, 129
 rbcL analysis of 73, 80–82
 rps16 analysis of 58
 trnL-F analysis of 58
Filgueirasia
 in Arthrostylidiinae (Bambuseae) phylogeny 308
 biogeography of 308
 Fimbristylideae 149
 C₄ photosynthesis in 99
Fimbristylis
 in Abildgaardieae phylogeny 149–164
 in Cyperaceae phylogeny 81
 5.8S rDNA insertion in 86–87, 92, 94, 96
 embryo type of 150
 geographic distribution of 149
 ITS analysis of 151, 155–156, 158–160
 monophyly and subgroups of 151
 morphology of 150
 one cell distant criterion in 103
 photosynthetic pathways in (C₃ and C₄) 101–102, 129–131, 161
 relationship with *Abildgaardia* 104, 150, 161–162
 relationship with *Bulbostylis* 150, 161
 relationship with *Crosslandia* 161–162
 sections of, assignment of species to 161–162
 trnL-F analysis of 81, 154–158
 trnL-F and ITS analysis of, combined 159–162
 Tylocarya inclusion in 150
Fimbristylis sect. *Abildgaardia* 161
Fimbristylis sect. *Actinoschoenus* 161
Fimbristylis sect. *Cymosae* 161
Fimbristylis sect. *Dichelostylis* 161
Fimbristylis sect. *Fimbristylis* 161
Fimbristylis sect. *Fuscae* 161
Fimbristylis sect. *Helocharoides* 161
Fimbristylis sect. *Leptocladae* 161
Fimbristylis sect. *Miliaceae* 161
Fimbristylis sect. *Mischospora* 161
Fimbristylis sect. *Neodichelostylis* 161
Fimbristylis sect. *Nutantes* 161–162
Fimbristylis sect. *Pogonostylis* 161
Fimbristylis sect. *Signatae* 161
Fimbristylis sect. *Tenerae* 161–162
Fimbristylis sect. *Trichelostylis* 161
Fingerhuthia
 in Chloridoideae phylogeny 566, 570–571, 573–574, 576, 581, 583
 ITS analysis of 566, 571
 ITS and *trnL-F* analysis of, combined 573
 morphological analysis of 583
 New World 581, 583
 trnL-F analysis of 566, 570
 Fission/fusion of chromosomes 62–63
 in *Carex* 193
 in Cyperales 66–67
Flagellaria, as outgroup in Bromeliaceae study 4
 Flagellariaceae
 relationship with Bromeliaceae 4, 8, 27–28
 relationship with Poaceae 255
 Fleshy fruits
 of *Alvimia* 308
 of Bromelioideae 3, 8, 12, 21
 Flower(s). *See also specific taxa*
 of *Aegilops triuncialis* 296
 of Andropogoneae 530
 of Bambuseae 315
 of *Bouteloua* 606
 of *Carex* 184–186
 of Centothecoideae + Panicoideae clade 491–492
 of Chloridoideae 565
 of Cyperaceae 56
 of Isachneae 506
 of Juncaceae 55
 of Paniceae 508
 of Panicoideae 503
 of Parianinae 310
 of *Paspalum* 545–546, 552, 555–556
 of Poaceae 259
 of Poaceae, chasmogamous and cleistogamous 286–294
 of Prionaceae 55–56
 of *Schoenus* 204–209
 of *Thrasya* 546, 552, 555–556
 of *Thrasypopsis* 546, 552, 556
 of Thurniaceae 55–56
 unisexual, evolution of 605–614
 Fossil record, of Bromeliaceae 5
Fosterella
 in Bromeliaceae phylogeny 3–4, 7–9, 14–17, 28, 34–36, 40–41
 CAM photosynthesis by 11, 41
 epiphytism of 12
 geographic origin and distribution of 10–11, 18, 35
 leaf and trichome anatomy of 13–14, 39–40
 matK-rps16 analysis of, combined 28

- morphology of 41
ndhF analysis of 3–4, 7–9, 14–17
 pollination of 13
 relationship with Bromelioideae 35, 40–41
trnL, *trnL-F*, and *trnT-F* analysis of, combined 34
trnL analysis of 28
trnL and *trnL-F* analysis of, combined 36
- Friar, Elizabeth A. 605–614
- Froesiochloa*, biogeography of 310
- Fruit(s)
 of Arthrotyliidiinae 308
 of Bromeliaceae 3, 8, 12
 of Chloridoideae 580–581
 of Poaceae 261
- Fruits, fleshy
 of *Alvimia* 308
 of Bromelioideae 3, 8, 12
- Fuirena*
 in Cyperaceae phylogeny 79, 81
 5.8S rDNA insertion in 88, 94
 genome size of 66
 photosynthetic pathway in 102, 131
rbcL analysis of 79, 81
- Fuireneae
 in Cyperaceae phylogeny 79
 5.8S rDNA insertion in, absence/presence of 88, 94
 ITS analysis of 156
 as outgroup in Abildgaardieae study 151
rbcL analysis of 79
- Furness, Carol A. 72–83
- Gahnia*
 in Cyperaceae phylogeny 78
 photosynthetic pathway in 102, 131
rbcL analysis of 78
- Galapageian Province, Panicoideae of 510–511
- Gastridium*, in Pooideae phylogeny 343, 346
- Gaudinia*
ndhF analysis of 341
 in Poaceae phylogeny 341
 in Pooideae phylogeny 343, 346
- GBSSI**
 in Bambuseae analysis 330
 in Chloridoideae analysis 572–574, 576
 in *Elymus* analysis 374, 376–378
 in *Eragrostis* analysis 572–574, 597–602
 in Poaceae analysis 248–249, 252, 260, 337, 343–344
 in Pooideae analysis 343
 in Triticeae analysis 369
- Genome size
 of Cyperales 65–68
 of *Festuca* 413, 416
- Germainiinae
 in Andropogoneae phylogeny 262, 532, 538–540
 biogeography of 515
 ITS analysis of 538
 ITS and *trnL-F* analysis of, combined 540
- New World 504–506, 515
 in Panicoideae phylogeny 504–505, 515
 photosynthetic pathways in 515
trnL-F analysis of 539
- Germination speed, and invasion 295–296
- Gerritea*
 biogeography of 509–510, 522
 New World 505, 508–510, 522
 in Panicoideae phylogeny 505, 508, 522
 photosynthetic pathway of 522
- Ghamkhar, Kioumars 72–83
- Ghamkhar, Kioumars et al.—Phylogeny of Abildgaardieae (Cyperaceae) inferred from ITS and *trnL-F* data 149–164
- Gillespie, Lynn J. et al.—Phylogeny of *Poa* (Poaceae) based on *trnT-trnF* sequence data: major clades and basal relationships 420–434
- Givnish, Thomas J. et al.—Phylogeny, adaptive radiation, and historical biogeography of Bromeliaceae inferred from *ndhF* sequence data 3–26
- Glaziophyton*
 in Arthrotyliidiinae phylogeny 308
 in Bambuseae phylogeny 308, 326–328
 biogeography of 308
 culms of 309
 internode lengths of 308
 morphological analysis of 324–326
rpl16 analysis of 322, 327
rpl16 and morphological analysis of, combined 328
 spikelets of 308
- Glomeropitcairnia*
 in Bromeliaceae phylogeny 7–9, 16, 21, 34–36
 CAM photosynthesis by 11
 epiphytism of 12
 geographic origin and distribution of 10–11
 leaf and trichome anatomy of 14
ndhF analysis of 7–9, 16, 21
 pollination of 13
trnL, *trnL-F*, and *trnT-F* analysis of, combined 34
trnL and *trnL-F* analysis of, combined 36
- Glyceria*
matK analysis of 342
 morphological analysis of 494
ndhF analysis of 342, 494–496
ndhH analysis of 342
 in Poaceae phylogeny 242–244, 342
rbcL analysis of 342
rpl16 analysis of 494, 496
- Goetghebeur, Paul 72–83, 204–209
- Goldman, Douglas H. 237–247
- Gouinia*
 biogeography of 587–588, 593
 in Chloridoideae phylogeny 566, 570–571, 573, 577, 581, 584
 ITS analysis of 566, 571
 ITS and *trnL-F* analysis of, combined 573

- morphological analysis of 584
 New World 581, 584, 587–588, 593
trnL-F analysis of 566, 570
- Gouiniinae** 592
 biogeography of 593
 in Chloridoideae phylogeny 581–582, 584
 morphological analysis of 584, 592
 New World 581–582, 584, 593
- Gouldochloa*
 in Centothecoideae + Panicoideae phylogeny 489, 493–498, 500
 geographic distribution of 489
 morphological analysis of 494, 498
ndhF analysis of 494–496
rpl16 analysis of 493–494, 496
 spikelets of 489
- Graham, Sean W. 450–467
- Gramineae. *See also* Poaceae
 Chloridoideae phylogeny 565–579
 Stipeae molecular analysis 349–361
- Grass(es). *See also* Poaceae
 diversification of 248–258
 electronic world database of 267–270
 morphological versus molecular classification of 259–266
 phylogeny of 237–266
 socioeconomic and ecological importance of 248
- Grasslands, Mediterranean climate, invasion in 295–296
- Gravel, Geneviève 84–98
- Greigia*
 AFLP analysis of 37
 in Bromeliaceae (Bromelioideae) phylogeny 29, 34, 35, 36–37, 41
 CAM photosynthesis by 41
 geographic origin and distribution of 35, 41
trnL, *trnL*-F, and *trnT*-F analysis of, combined 34
trnL and *trnL*-F analysis of, combined 35–36
- Greslania*
 in Bambuseae phylogeny 315, 323, 326–328, 330
 culms of 316
 glumes of 316
 morphological analysis of 324, 326
 morphology of 316
 in Poaceae phylogeny 251
rpl16 analysis of 327
rpl16 and morphological analysis of, combined 328
- Griffith, M. Patrick 565–579
- Guacamaya* 7
- Guadua*
 in Amazonia 311
 in Bambuseae phylogeny 309, 317, 326–328
 biogeography of 309, 311
 in Guaduinae phylogeny 309
matK analysis of 342
 morphological analysis of 324, 326
ndhF analysis of 342
ndhH analysis of 342
 in Poaceae phylogeny 341–342
rbcL analysis of 342
rpl16 analysis of 327
rpl16 and morphological analysis of, combined 328
- Guaduelia*
matK analysis of 342
ndhF analysis of 342
ndhH analysis of 342
 in Poaceae phylogeny 242–244, 342
 in Puelioideae phylogeny 303
rbcL analysis of 342
- Guaduelleae, in Poaceae phylogeny 252–253
- Guaduinae
 in Bambuseae phylogeny 306, 309, 315–316, 323–330
 biogeography of 309, 311
 in eastern Brazil 311
 elevation range of 309
 in Guayana Highlands, Venezuela 311
 morphological analysis of 326, 329
ndhF analysis of 329
 papillae of 309
 phylogeny of 309
 relationship with Arthrostylidiinae 308–309
 relationship with Chusqueinae 316
rpl16 analysis of 327, 329
rpl16 and morphological analysis of, combined 328
 stomata of 309
- Guayana Highlands, Venezuela, Bambusoideae in 311
- Guayana Shield
 Bromeliaceae origin and distribution in 4–5, 7, 9–12, 18–19
 Bromelioideae origin and distribution in 29
- Guiana Highlands Region, Panicoideae of 510
- Guzmania*
 in Bromeliaceae phylogeny 7–9, 28, 34–36
 CAM photosynthesis by 11
 epiphytism of 12
 geographic origin and distribution of 10–11, 18
 leaf and trichome anatomy of 14
ndhF analysis of 7–9
 pollination of 13, 22
trnL, *trnL*-F, and *trnT*-F analysis of, combined 34
trnL and *trnL*-F analysis of, combined 36
- Gymnachne*, tribal/subtribal placement of 423
- Gymnopogon*
 biogeography of 587–588, 593
 in Chloridoideae phylogeny 566, 570–571, 573, 580–581
 ITS analysis of 566, 571
 ITS and *trnL*-F analysis of, combined 573
 New World 581, 587–588, 593
trnL-F analysis of 566, 570
- Gymnoschoenus*
 in Cyperaceae phylogeny 78
 photosynthetic pathway in 102, 131

- rbcL* analysis of 78
- Gynerieae
 in Centothecoideae + Panicoideae phylogeny 499
 morphological analysis of 506
ndhF analysis of 344
 New World 504–506
 in Panicoideae phylogeny 504–506
 in Poaceae phylogeny 245, 344
- Gynerium*
 biogeography of 509
 in Centothecoideae + Panicoideae phylogeny 489, 494–497, 499
 culms of 325
matK analysis of 342
 morphological analysis of 494
ndhF analysis of 342, 344, 494–496
ndhH analysis of 341–342
 New World 505–506, 509
 in Panicoideae phylogeny 504–506
 in Poaceae phylogeny 237, 241–245, 254, 260, 341–342, 344
rbcL analysis of 342
 relationship with *Steyermarkochloa* 508
rpl16 analysis of 494, 496
- Gynodioecy 605–614
- Hackelochloa*
 in Andropogoneae phylogeny 535, 538–540
 ITS analysis of 535, 538, 567, 571
 ITS and *trnL*-F analysis of, combined 540, 573
 as outgroup in Chloridoideae study 567
trnL-F analysis of 535, 539, 567, 570
- Hainardia*
 molecular analysis of 390–392
 morphological analysis of 387, 394, 396–399
 morphological-molecular analysis of, combined 397
 in Poaceae phylogeny 403
 relationship with Loliinae 382, 387, 390–394, 396–399, 401
- Hainardieae 402
 morphology of 380
 in Poaceae phylogeny 252–253, 255–256
 in Pooideae phylogeny 335, 345–346, 380
- Hakonechloa*, in Poaceae phylogeny 251, 254
- HAMBADD clade, of *Poa* 421, 429–430, 432
- Hardy, Christopher R. and H. Peter Linder—Phylogeny and historical ecology of *Rhodocoma* (Restionaceae) from the Cape Floristic Region 213–226
- Harpochloa*, in Chloridoideae phylogeny 580
- Hechtia*
 in Bromeliaceae phylogeny 3–4, 7–9, 14–16, 28, 34–35, 40
 CAM photosynthesis by 11–12
 concerted convergence in 14, 16, 22
 epiphytism of 12
 geographic origin and distribution of 10, 18–19
 leaf and trichome anatomy of 13–14, 16, 20, 22
matK-rps16 analysis of, combined 28
 morphology of 35
ndhF analysis of 3–4, 7–9, 14–16
 pollination of 13
trnL, *trnL*-F, and *trnT*-F analysis of, combined 34
trnL analysis of 28
- Hechtioideae** 16
 in Bromeliaceae phylogeny 9, 16
 CAM photosynthesis by 11
 epiphytism of 12
 geographic origin and distribution of 10
 leaf and trichome anatomy of 14
 morphology of 16
ndhF analysis of 9, 16
 pollination of 13
 Heilborn's hypothesis on chromosome evolution 194, 198, 202
- Helictotrichon*
ndhF analysis of 341
 as outgroup in *Poa* study 422
 in Poaceae phylogeny 341
 in Pooideae phylogeny 343, 346
- Hellerochloa*
 in Loliinae phylogeny 381–382, 387, 390–393, 397–398
 molecular analysis of 390–392, 414–415
 morphological analysis of 387, 393
 morphological-molecular analysis of, combined 397–398
 relationship with *Festuca* 406–407, 413
- Hellmuthia*
 absence of 5.8S rDNA insertion in 88, 92, 94, 96
 in Cyperaceae phylogeny 73, 80–81
 geographic distribution of 73
 photosynthetic pathway in 102, 131
rbcL analysis of 73, 80–81
- Hemarthria*
 in Andropogoneae phylogeny 532, 535, 538–540
 ITS analysis of 535, 538
 ITS and *trnL*-F analysis of, combined 540
 New World 505
 in Panicoideae phylogeny 505
trnL-F analysis of 535, 539
- Henrardia*
 chloroplast DNA analysis of 366, 368
DMC1 analysis of 366–367, 370
EF-G analysis of 366–367, 370
 mitochondrial genome of 366
 morphological analysis of 365–366
 in Poaceae phylogeny 251
 total evidence (combined) analysis of 367–369
 in Triticeae phylogeny 365–370
- Herbaceous bamboos. *See* Olyreae
- Hesperostipa*
 ITS analysis of 356–357, 359
matK analysis of 342

- ndhF* analysis of 342
ndhH analysis of 342
 in Poaceae phylogeny 342
rbcL analysis of 342
 relationship with *Austrostipa* 358
 in Stipeae phylogeny 350–351, 356–357, 359–360
- Heterachne*, in Chloridoideae phylogeny 576
- Heterantherium*
 chloroplast DNA analysis of 366
DMCI analysis of 367, 370
EF-G analysis of 367, 370
 mitochondrial genome of 366
 morphological analysis of 365
 total evidence (combined) analysis of 367–369
 in Triticeae phylogeny 365–370
- Heteranthoecia*, in Panicoideae phylogeny 503
- Heteropogon*
 in Andropogoneae phylogeny 530, 532, 535, 537–540, 542
 ITS analysis of 535, 538
 ITS and *trnL*-F analysis of, combined 537, 540, 542
 New World 505
 in Panicoideae phylogeny 505
 spikelets of 530
trnL-F analysis of 535, 539
- Hickelia*
 in Bambuseae phylogeny 315, 326–328
 morphological analysis of 324, 326
rpl16 analysis of 327
rpl16 and morphological analysis of, combined 328
- Hickelinae
 in Bambuseae phylogeny 315–316, 323, 326–328, 330
 buds/branching of 316
 glumes of 316
 morphological analysis of 326, 330
 morphology of 316
ndhF analysis of 316
 plastid genome analysis of 316
 relationship with Bambusinae 316
 relationship with Melocanninae 316
 rhizomes of 330
rpl16 analysis of 316, 327
rpl16 and morphological analysis of, combined 328
 spikelets of 316
 synflorescences of 316
- Hierochloë*. See *Anthoxanthum*
- Hilaria*
 biogeography of 593
 in Chloridoideae phylogeny 566, 570–571, 573, 581, 584
 ITS analysis of 566, 571
 ITS and *trnL*-F analysis of, combined 573
 morphological analysis of 584
 New World 581, 584, 593
trnL-F analysis of 566, 570
- Hilariinae** 592
 biogeography of 587, 589–590, 593
 in Chloridoideae phylogeny 581–582, 584
 morphological analysis of 584, 592
 New World 581–582, 584, 587, 589–590, 593
- Hipp, Andrew L. et al.—Chromosome number changes associated with speciation in sedges: a phylogenetic study in *Carex* section *Ovales* (Cyperaceae) using AFLP data 193–203
- Hodkinson, Trevor R. 55–61, 72–83
- Hodkinson, Trevor R. et al.—Large trees, supertrees, and diversification of the grass family 248–258
- Hohenbergia*
 in Bromeliaceae phylogeny 34
trnL, *trnL*-F, and *trnT*-F analysis of, combined 34
trnL and *trnL*-F analysis of, combined 36
- Hohenbergiopsis*
 in Bromeliaceae phylogeny 36
trnL and *trnL*-F analysis of, combined 36
- Holcinae, in Pooideae phylogeny 346–347
- Holcus*, in Pooideae phylogeny 343, 346–347
- Holocentric chromosomes
 of *Carex* 193
 of Cyperales 63–64
- Homolepis*
 biogeography of 509, 522
 New World 505, 508–509, 522
 in Panicoideae phylogeny 505, 508, 522
 photosynthetic pathway of 522
- Homologous series in variation, law of 261
- Homoplasy, in Poaceae 260–261
- Hookerchloa*, tribal/subtribal placement of 423
- Hordeum*
 BAC studies of 68
 chloroplast DNA analysis of 366, 368
DMCI analysis of 363, 366–367, 369–370
EF-G analysis of 363, 366–367, 369–370
 genome transfer to *Elymus* 372–378
 mitochondrial genome of 366
 morphological analysis of 363, 365–366
ndhF analysis of 341
 in Poaceae phylogeny 341
 in Pooideae phylogeny 343, 345
 relationship with *Critesion* 369–370
 total evidence (combined) analysis of 367–369
 in Triticeae phylogeny 362–363, 365–370
- Horres, Ralf et al.—Systematics of Bromelioideae (Bromeliaceae)—evidence from molecular and anatomical studies 27–43
- Hubbardia*, in Centothecoideae + Panicoideae phylogeny 497
- Hubbardieae
 in Centothecoideae + Panicoideae phylogeny 488, 499
 in Isachneae phylogeny 503
 in Panicoideae phylogeny 503
- Hummingbird pollination, of Bromeliaceae 11–13, 21–22

- Hunter, Angela M. 406–419
- Hyalopoa*, tribal/subtribal placement of 423
- Hymenachne*
 biogeography of 522
 New World 505, 508, 522
 in Panicoideae phylogeny 505, 508, 522
 photosynthetic pathway of 522
- Hyparrhenia*
 in Andropogoneae phylogeny 530, 532, 535, 537–542
 biogeography of 541
 ITS analysis of 535, 538, 541
 ITS and *trnL*–*F* analysis of, combined 537, 540–542
 morphological analysis of 541
 New World 505
 in Panicoideae phylogeny 505
PHYB analysis of 541
trnL–*F* analysis of 535, 539, 541
- Hyperthelia*
 in Andropogoneae phylogeny 532, 535, 537–541
 ITS analysis of 535, 538, 541
 ITS and *trnL*–*F* analysis of, combined 537, 540–541
 morphological analysis of 541
 New World 504–505
 in Panicoideae phylogeny 504–505
trnL–*F* analysis of 535, 539, 541
- Hypolytreae
 in Cyperaceae phylogeny 73–78
 5.8S rDNA insertion in 88, 94, 96
 pollen of 77
rbcL analysis of 73–78
- Hypolytrum*
 in Cyperaceae phylogeny 72–73, 78
 5.8S rDNA insertion in 86, 88
 in Juncaceae phylogeny 58
 photosynthetic pathway in 102, 131–132
rbcL analysis of 73, 78
rps16 analysis of 57–58
trnL–*F* analysis of 57–58
- Ichnanthus*
 biogeography of 511, 522–523
 New World 505, 508, 511, 522–523
 in Panicoideae phylogeny 505, 508, 522–523
 photosynthetic pathway of 522–523
- Imperata*
 in Andropogoneae phylogeny 262, 530, 532, 535, 538–540
 biogeography of 515
 ITS analysis of 535, 538
 ITS and *trnL*–*F* analysis of, combined 540
 New World 505, 515
 in Panicoideae phylogeny 505, 515
 photosynthetic pathway of 515
trnL–*F* analysis of 535, 539
- Inflorescence. *See also specific taxa*
 of *Aegilops triuncialis* 296
 of Andropogoneae 530
 of *Bouteloua* 606
 of *Carex* 184–186
 of Chloridoideae 565
 of Juncaceae 55
 of Paniceae 508
 of Parianinae 310
 of *Paspalum* 545–546, 552, 555–556
 of Poaceae 259
 of *Schoenus* 204–209
 of *Thrasya* 546, 552, 555–556
 of *Thrasypsis* 546, 552, 556
- Ingram, Amanda L. and Jeff J. Doyle—*Eragrostis* (Poaceae): monophyly and infrageneric classification 595–604
- Insect pollination
 of Bromeliaceae 12–13
 of *Pariana* 310
- Internal transcribed spacer (ITS)
 in Abildgaardieae analysis 151–156, 158–162
 in Andropogoneae analysis 262, 530–544
 in *Bromus* analysis 450–467
 in *Carex* sect. *Ovales* analysis 194
 in Cariceae analysis 165–192
 in Chloridoideae analysis 565–579, 582
 in Cyperaceae analysis 81, 84–98
 in *Eleocharis* analysis 95
 in *Festuca* analysis 381, 386, 389–390, 392, 397–398, 406–419
 in Loliinae analysis 382, 385–399
 in Poaceae analysis 248–249, 251–254, 259–260
 in Stipeae analysis 349–360
 in Triodieae analysis 263
- Invasion
 of *Aegilops triuncialis* 295–300
 bet-hedging strategy in 296, 298
 germination speed and 295–296
 lag phase of 295
 phenotypic plasticity and 295–300
 seed dormancy in 296
- Isachne*
 biogeography of 497, 510–511, 516
ndhF analysis of 241–242, 261
 New World 505–506, 510–511, 516
 in Panicoideae phylogeny 503–506, 516
 photosynthetic pathway of 516
 in Poaceae phylogeny 237, 241–244, 261, 497, 500
 synapomorphy of 261
- Isachneae
 base chromosome number of 506
 biogeography of 516
 in Centothecoideae + Panicoideae phylogeny 488, 491–500
 Hubbardieae placement in 503
ndhF analysis of 490–500
 New World 504–506, 516
 in Panicoideae phylogeny 503–506, 516

- photosynthetic pathways in 516
 in Poaceae phylogeny 252–253
rpl16 analysis of 490–500
 spikelets of 506
- Ischaemeae, in Andropogoneae phylogeny 532
- Ischaeminae
 in Andropogoneae phylogeny 262, 532, 538–540
 biogeography of 515
 ITS analysis of 538
 ITS and *trnL*-F analysis of, combined 540
 New World 504–506, 515
 in Panicoideae phylogeny 503–505, 515
 photosynthetic pathways in 515
trnL-F analysis of 539
- Ischaemum*
 in Andropogoneae phylogeny 530
 biogeography of 515
 New World 505, 515
 in Panicoideae phylogeny 505, 515
 photosynthetic pathways in 515
- Iseilema*
 in Andropogoneae phylogeny 532, 535, 537–540
 ITS analysis of 535, 538
 ITS and *trnL*-F analysis of, combined 537, 540, 542
trnL-F analysis of 535, 539
- Isolepis*
 absence of 5.8S rDNA insertion in 87, 92, 94, 96
 in Cyperaceae phylogeny 72–73, 79–82, 84
 in Cyperales phylogeny 68
 photosynthetic pathway in 102, 132
rbcL analysis of 73, 79–82
- Isolepis humillima*
 in Cyperaceae phylogeny 81
 relationship with *Schoenoplectus* 81
- Ixophorus*
 biogeography of 509–510
 New World 505, 508–510
 in Panicoideae phylogeny 505, 508
- Jacobs, Surrey et al.—Systematics of the tribe Stipeae (Gramineae) using molecular data 349–361
- Jarava*
 geographic distribution of 350
 ITS analysis of 356–357, 359
 morphology of 350
 relationship with *Achnatherum* 349–350, 355, 359
 in Stipeae phylogeny 349–350, 356–357, 359
 wind dispersal of 350
- Joinvillea*
matK analysis of 342
ndhF analysis of 341–342
ndhH analysis of 341–342
 as outgroup in Bromeliaceae study 4
 as outgroup in Poaceae study 238–246
 plastid genome analysis of 340–342
rbcL analysis of 342
 relationship with Poaceae 238
- Joinvilleaceae
ndhH analysis of 341–344
 plastid genome analysis of 336–337
 relationship with Bromeliaceae 4, 8
 relationship with Poaceae 237–238, 255, 260
- Jones, Eleanor et al.—The Juncaceae–Cyperaceae interface: a combined plastid sequence analysis 55–61
- Jouvea*
 biogeography of 587, 593
 in Chloridoideae phylogeny 566, 570–571, 573, 581
 ITS analysis of 566, 571
 ITS and *trnL*-F analysis of, combined 573
 New World 581, 587, 593
trnL-F analysis of 566, 570
- Joycea*, in Danthonioideae phylogeny 261
- Judziewicz, Emmet J. and Lynn G. Clark—Classification and biogeography of New World grasses: Anomochlooideae, Pharoideae, Ehrhartoideae, and Bambusoideae 303–314
- Juncaceae
 bracts of 55, 60
 in Cyperales phylogeny 55, 68
 5.8S rDNA insertion in, absence/presence of 86, 89, 93–95
 flowers of 55–56, 60
 genomic resources in 64–65
 geographic distribution of 55
 growth form of 55
 holocentric chromosomes in 63
 inflorescences of 55
 leaf variation among 55, 60
 phylogeny of 55–61
 plastid genome analysis of 55–61
 in Poales phylogeny 55
 relationship with Bromeliaceae 27–28
 relationship with Cyperaceae 55–61, 72, 85
 relationship with Prioniaceae 55–56
 relationship with Thurniaceae 55–56
 rhizomes of 55
rps16 analysis of 55–61
trnL analysis of 55–61
trnL-F analysis of 55–61
- Juncus*
 BAC studies of 68
 bracts of 55, 60
 clades of 60
 in Cyperales phylogeny 68
 5.8S rDNA insertion in, absence/presence of 86, 89, 94
 flowers of 55, 60
 genome size of 66
 inflorescence of 55
 in Juncaceae phylogeny 55–56, 58–60
 leaf variation in 55, 60
 in New Zealand 418
 paraphyly of 60
rps16 analysis of 57, 59

- trnL*-F analysis of 57, 59
- Juncus* subgen. *Agathyron* 60
- Juncus* subgen. *Juncus* 60
- Juncus* subgen. *Septati* 60
- Kaokochloa*, in Chloridoideae phylogeny 583
- Karatophyllum*, fossil record of 5
- Karinia*, photosynthetic pathway in 102, 132
- Karroochloa*
- morphological analysis of 494
 - ndhF* analysis of 494–496
 - in Poaceae phylogeny 242–244
 - rpl16* analysis of 494, 496
- Keane, Ryan M. 450–467
- Kengia*, in Chloridoideae phylogeny 572
- Khaosokia*
- in Cyperaceae phylogeny 72–73, 79, 81
 - photosynthetic pathway in 102, 132
 - rbcL* analysis of 73, 79, 81
- Kinney, Michael S. 565–579
- Kinney, Michael S. et al.—Dicliny in *Bouteloua* (Poaceae: Chloridoideae): implications for the evolution of dioecy 605–614
- Kobresia*
- in Cariceae phylogeny 175, 177–179, 183, 188–189
 - in Cyperaceae phylogeny 79
 - 5.8S rDNA insertion in 85
 - ITS, ETS-1f, and *trnL*-F analysis of, combined 166, 175, 177–178
 - ITS analysis of 179
 - photosynthetic pathway in 103, 132
 - rbcL* analysis of 79
- Kobresia* subgen. *Compositae* 171
- Kobresia* subgen. *Kobresia* 172
- Kobresia* subgen. *Kobresia* sect. *Hemicarex* 172
- Kobresia* subgen. *Kobresia* sect. *Kobresia* 172
- Koyamaea*, photosynthetic pathway in 103, 132
- Kranz syndrome
- in Arthropogoninae 507
 - in Chloridoideae 580
 - in Melinidinae 507
 - in Paniceae 508
 - in Panicoideae 504, 511–512, 514–529
 - in Paspalinae 508
 - in Setariinae 508
- Kunhardtia* 7
- Kyllinga*
- absence of 5.8S rDNA insertion in 87, 94
 - in Cyperaceae phylogeny 80
 - photosynthetic pathway in 103, 132–133
 - rbcL* analysis of 80
- Kyllingiella*
- absence of 5.8S rDNA insertion in 87, 92, 94
 - in Cyperaceae phylogeny 80–81
 - photosynthetic pathway in 103, 133–134
 - rbcL* analysis of 80
- Lagenocarpus*
- in Cyperaceae phylogeny 77–78
 - 5.8S rDNA insertion in 88
 - photosynthetic pathway in 101, 103, 134
 - rbcL* analysis of 77–78
- Lamarckia*
- molecular analysis of 390–392
 - morphological analysis of 387, 393–394
 - morphological-molecular analysis of, combined 397
 - in Poeae phylogeny 380, 403
 - relationship with Loliinae 382, 387, 390–394, 397, 401
- Lasiacis*
- biogeography of 509, 520
 - New World 505, 508–509, 520
 - in Panicoideae phylogeny 505, 508, 520
 - photosynthetic pathway of 520
 - in Poaceae phylogeny 254
- Leaf anatomy
- of *Anthoxanthum odoratum* versus *A. alpinum* 473–480
 - of Bambusoideae 303
 - of Centothecoideae + Panicoideae clade 488, 491–492
 - of Chloridoideae 580
 - of *Paspalum* 545, 552, 555
 - of Poaceae 259, 261
 - of *Thrasya* 546, 552, 555
 - of Triodieae 263
- Lecomtella*, in Panicoideae phylogeny 503
- Lecomtelleae, in Panicoideae phylogeny 503
- Lecomtellinae, in Panicoideae phylogeny 503
- Lee, William G. 406–419
- Leersia*
- biogeography of 306
 - in Oryzae (Ehrhartoideae) phylogeny 306
 - in Poaceae phylogeny 242–244
- Lepidosperma*
- in Cyperaceae phylogeny 78
 - photosynthetic pathway in 103, 134–135
 - rbcL* analysis of 78
- Lepironia*
- in Cyperaceae phylogeny 78
 - 5.8S rDNA insertion in 88, 94
 - photosynthetic pathway in 103, 135
 - rbcL* analysis of 78
- Leptochloa*
- biogeography of 587–588, 593
 - in Chloridoideae phylogeny 566, 570–573, 575, 577, 580–581
 - in Danthonioideae phylogeny 259
 - ITS analysis of 566, 571–572, 575
 - ITS and *trnL*-F analysis of, combined 572–573
 - morphological analysis of 575, 577
 - New World 581, 587–588, 593
 - trnL*-F analysis of 566, 570
- Leptothrium*
- biogeography of 588, 593
 - in Chloridoideae phylogeny 566, 570–571, 573, 581

- ITS analysis of 566, 571
 ITS and *trnL*-F analysis of, combined 573
 New World 581, 588, 593
trnL-F analysis of 566, 570
- Leptureae
 in Chloridoideae phylogeny 574–575
 in Poaceae phylogeny 252–253, 255
- Lepturidium*
 biogeography of 589, 593
 in Chloridoideae phylogeny 565–567, 570–571, 573
 New World 565–567, 589, 593
- Leucopoa*
 in Loliinae phylogeny 381, 400
 in Pooideae phylogeny 343, 346
- Libyella*, tribal/subtribal placement of 423
- Liliales, concerted convergence in 22
- Limnas*, tribal/subtribal placement of 423
- Limnopoa*, in Panicoideae phylogeny 503
- Lindbergella*, tribal/subtribal placement of 423
- Linder, H. Peter 213–226
- Lindmania*
 in Bromeliaceae phylogeny 4, 7–9, 14–16, 28
 CAM photosynthesis by 11
 epiphytism of 12
 geographic origin and distribution of 9–10
 leaf and trichome anatomy of 14
matK-rps16 analysis of, combined 28
ndhF analysis of 4, 7–9, 14–16
 pollination of 13
- Lindmanioidae** 15–16
 in Bromeliaceae phylogeny 9, 15–16
 CAM photosynthesis by 11
 epiphytism of 12
 geographic origin and distribution of 10
 leaf and trichome anatomy of 14
 morphology of 15–16
ndhF analysis of 9, 15–16
 pollination of 13, 21
- Lipocarpha*
 absence of 5.8S rDNA insertion in 87, 94
 in Cyperaceae phylogeny 80
 photosynthesis in, conflicting data on 104, 148
 photosynthetic pathway in 103–104, 135
rbcL analysis of 80
- Lithachne*
 florets of 311
 ITS analysis of 351, 355–356
 leaf blades of 311
matK analysis of 342
ndhF analysis of 342
ndhH analysis of 342
 in Olyrinae (Olyreae) phylogeny 311
 in Poaceae phylogeny 241–245, 342
rbcL analysis of 342
 relationship with Stipeae 350–351, 355–356
 sleep movements of 311
- Little bluestem. *See Schizachyrium scoparium*
- Littledalea*
 in Pooideae phylogeny 343, 345
 relationship with *Bromus* 450, 463
- Lloyd, Kelvin M. et al.—Phylogeny and biogeography of endemic *Festuca* (Poaceae) from New Zealand based on nuclear (ITS) and chloroplast (*trnL*-*trnF*) nucleotide sequences 406–419
- Lolieae 402
- Loliinae
 broad-leaved group of 389–393, 396–401
 cladistic analysis of 385, 399
 evolutionary rate in 381
 fine-leaved group of 389–393, 396–401
 ITS analysis of 382, 385–399
 molecular analysis of 382, 385–399
 morphological analysis of 381–393
 morphological-molecular analysis of, combined 382, 393–399
 phylogeny of 380–405
 in Poeae phylogeny 402–403, 421
 in Pooideae phylogeny 346
 relationship with *Poa* 421–422
 reticulate history of 399–400
trnL-F analysis of 382, 385–399
- Lolium*
 inflorescence traits of 380
 in Loliinae phylogeny 380–405
 molecular analysis of 389–392, 414–415
 morphological analysis of 382–385, 387, 393–399
 morphological-molecular analysis of, combined 397–398
 in Pooideae phylogeny 343, 346, 380
 relationship with *Elymus* 380
 relationship with *Festuca* 380–381, 406–413
- López-Rodríguez, José A. 380–405
- Lophatherum*
 in Centothecoideae + Panicoideae phylogeny 494–498
 geographic distribution of 489
 morphological analysis of 494
ndhF analysis of 494–496
rpl16 analysis of 494, 496
 spikelets of 489
- Lopholepis*, in Poaceae phylogeny 251, 254
- Lophopyrum*
 chloroplast DNA analysis of 366, 368
DMCI analysis of 367
EF-G analysis of 367
 mitochondrial genome of 366
 morphological analysis of 365
 total evidence (combined) analysis of 367–369
 in Triticeae phylogeny 362, 365–369
- Loudetia*
 biogeography of 516
 in Centothecoideae + Panicoideae phylogeny 494–498
 morphological analysis of 494

- ndhF* analysis of 494–498
 New World 504–505, 516
 in Panicoideae phylogeny 504–505, 516
 photosynthetic pathway of 516
rpl16 analysis of 494, 496–498
- Loudetiopsis*
 New World 504–505
 in Panicoideae phylogeny 504–505
- Luziola*
 biogeography of 306
 in Luziolinae (Oryzeae) phylogeny 306
 in Poaceae phylogeny 251
- Luziolinae
 biogeography of 306
 in Oryzeae (Ehrhartoideae) phylogeny 306
- Luzula*
 base chromosome number of 66–67
 5.8S rDNA insertion in, absence/presence of 89, 92, 94
 genome size of 66–67
 inflorescence of 55
 in Juncaceae phylogeny 55–56, 58
 kinetochore binding sites of 63
rps16 analysis of 57, 59
trnL–*F* analysis of 57, 59
- Lycurus*
 amphitropical disjunct in 584
 biogeography of 588, 593
 in Chloridoideae phylogeny 566, 570–571, 573, 577, 581, 584
 ITS analysis of 566, 571
 ITS and *trnL*–*F* analysis of, combined 573
 morphological analysis of 584–585
 New World 581, 584, 588, 593
trnL–*F* analysis of 566, 570
- Lygeae
 in Poaceae phylogeny 252–253
 in Pooideae phylogeny 345
- Lygeum*
matK analysis of 342
ndhF analysis of 342
ndhH analysis of 342
 as outgroup in Stipeae study 351
 in Poaceae phylogeny 242–244, 342
rbcL analysis of 342
- Lymania*
 in Bromeliaceae phylogeny 36
trnL and *trnL*–*F* analysis of, combined 36
- Machaerina*, photosynthetic pathway in 103, 135
- Maclurolyra*
 biogeography of 310
 in Olyrinae (Olyreae) phylogeny 310–311
 synflorescence of 311
- Macrochloa*, in Stipeae phylogeny 349
- Madrean Region, Panicoideae of 510
- Mapania*
 in Cyperaceae phylogeny 73, 78
 5.8S rDNA insertion in 88, 94
 in Juncaceae phylogeny 58
 photosynthetic pathway in 103, 135–136
 pollen of 77
 pseudomonads of 56, 77
rbcL analysis of 73, 78
rps16 analysis of 57–58
trnL–*F* analysis of 57–58
- Mapanioideae
 in Cyperaceae phylogeny 72–78, 82
 5.8S rDNA insertion in, absence/presence of 88
 morphology of 82
 one- or two-tribe composition of 73–77
rbcL analysis of 72–78
- Marchant, Adam D. 72–83, 149–164
- Marsippospermum*
 5.8S rDNA insertion in 89
 flower of 55, 60
 inflorescence of 55
 in Juncaceae phylogeny 55, 58–60
 relationship with *Oxychloe* 90
rps16 analysis of 57, 59
trnL–*F* analysis of 57, 59
- Maschalocephalus*, geographic origin and distribution of 19
- Mason-Gamer, Roberta J.—Allopolyploids of the genus *Elymus* (Triticeae, Poaceae): a phylogenetic perspective 372–379
- matK*
 in Bromeliaceae analysis 3–4, 15, 28
 in Chloridoideae analysis 342, 565, 572–577, 582
 in *Eragrostis* analysis 596–597, 600
 in Poaceae analysis 241, 248, 252, 254, 259–260, 337–343
 in *Rhodocoma* analysis 213–214, 218
- Mayaca* 6
- Mayacaceae, relationship with Bromeliaceae 4, 8, 27–28
- Maydeae
 in Andropogoneae phylogeny 531–533, 543
 in Panicoideae phylogeny 503
 spikelets of 543
- McCubbin, Andrew G. 62–71
- Mediterranean climate grasslands, invasion in 295–296
- Megalachne*, relationship with *Bromus* 450, 463
- Megastachya*
 in Centothecoideae + Panicoideae phylogeny 489, 494–498
 geographic distribution of 489
 morphological analysis of 494
ndhF analysis of 494–496
rpl16 analysis of 494, 496
 spikelets of 489
- Megathyrsus*
 New World 505, 507
 in Panicoideae phylogeny 505, 507
- Melanacra clade
 in Bromeliaceae phylogeny 7

- ndhF* analysis of 7
- Melanocenchris*
 in Chloridoideae phylogeny 566, 570–571, 573
 ITS analysis of 566, 571
 ITS and *trnL*-F analysis of, combined 573
trnL-F analysis of 566, 570
- Melica*
matK analysis of 342
 morphological analysis of 494
ndhF analysis of 342, 494–496
ndhH analysis of 342
 in Poaceae phylogeny 242–244, 342
rbcL analysis of 342
rpl16 analysis of 494, 496
- Meliceae
 in Poaceae phylogeny 252–253
 in Pooideae phylogeny 345–346
- Melinideae
 in Panicoideae phylogeny 503
 photosynthetic pathway of 507
- Melinidinae
 biogeography of 510, 518–519
 New World 504–507, 510, 518–519
 in Panicoideae phylogeny 503–505, 507, 518–519
 photosynthetic pathway of 518–519
- Melinis*
 New World 505, 507
 in Panicoideae phylogeny 503, 505, 507
- Melocalamus*, in Poaceae phylogeny 251
- Melocanninae
 in Bambuseae phylogeny 306, 316, 323, 326–330
 morphological analysis of 323, 326, 330
 relationship with Hickelinae 316
rpl16 analysis of 327
rpl16 and morphological analysis of, combined 328
- Merostachys*
 in Arthrostylidiinae (Bambuseae) phylogeny 308–309
 biogeography of 308–309
 branching of 308
 buds of 308
 in Poaceae phylogeny 251
- Merxmuellera*
 in Danthonioideae phylogeny 261
matK analysis of 342
 morphological analysis of 494
ndhF analysis of 342, 494–496
ndhH analysis of 342
 in Poaceae phylogeny 242–244, 342
rbcL analysis of 342
rpl16 analysis of 494, 496, 498
- Mesoamerica
 Bambusoideae in mountains of 311
 Panicoideae in 509–510
- Mesomelaena*
 in Cyperaceae phylogeny 78
 photosynthetic pathway in 103
- rbcL* analysis of 78
- Mesosetum*
 biogeography of 509, 523
 New World 505, 508–509, 523
 in Panicoideae phylogeny 505, 508, 523
 photosynthetic pathway of 511, 523
- Metcalfia*
 in Pooideae phylogeny 346
 relationship with *Bromus* 450, 463
- Mexico
 Chloridoideae in 583, 585–588
 Panicoideae in 509–510, 512
- Mezobromelia*
 in Bromeliaceae phylogeny 4
ndhF analysis of 4
- Mibora*, in Pooideae phylogeny 343, 346
- Miborinae, in Pooideae phylogeny 346
- Micraira*
 geographic distribution of 497
ndhF analysis of 241–242, 261, 497
 in Poaceae phylogeny 237–238, 241–244, 260–261,
 497, 500, 504, 506
rpl16 analysis of 493
 synapomorphy of 261
- Micraira* clade 497
- Micraireae
 in Centothecoideae + Panicoideae phylogeny 499
 in Poaceae phylogeny 252–253
- Micrairoideae, in Poaceae phylogeny 241–245
- Microbriza*, tribal/subtribal placement of 423
- Microchloa*
 biogeography of 593
 breeding system of 608–610
 in Chloridoideae phylogeny 566, 570–573, 577, 580–
 581, 584
 ITS analysis of 566, 571–572
 ITS and *trnL*-F analysis of, combined 573
 morphological analysis of 584
 New World 581, 584, 593
trnL-F analysis of 566, 570
- Microdracoides*
 in Cyperaceae phylogeny 78
 photosynthetic pathway in 103, 136
rbcL analysis of 78
- Microlaena*, in Poaceae phylogeny 251
- Micropyropsis*
 in Loliinae phylogeny 380–405
 molecular analysis of 389–392
 morphological analysis of 387, 393–399
 morphological-molecular analysis of, combined 397–
 398
- Micropyrum*
 in Loliinae phylogeny 380–405
 molecular analysis of 386, 414–415
 morphological analysis of 389, 394–395
 morphological-molecular analysis of, combined 397

- relationship with *Festuca* 406–407
- Microstegium*
- in Andropogoneae phylogeny 542
 - cleistogamy and chasmogamy in 287–293
 - invasion of eastern U. S. by 288
 - New World 504–505
 - in Panicoideae phylogeny 504–505
 - spikelets of 288
- Miliinae
- placement of genera in 423–424
 - in Poeae phylogeny 421
 - in Pooideae phylogeny 346
 - relationship with *Poa* 421–427, 431–433
- Milium*
- in Poeae phylogeny 421
 - in Pooideae phylogeny 343, 345–346
 - relationship with *Poa* 421, 427, 433
 - tribal/subtribal placement of 423
- Millam, Kendra C. 3–26
- Miscanthus*
- in Andropogoneae phylogeny 530, 532, 535, 538–540, 542–543
 - ITS analysis of 535, 538
 - ITS and *trnL*–F analysis of, combined 540, 542–543
 - ndhF* analysis of 542
 - New World 504–505
 - in Panicoideae phylogeny 504–505
 - in Poaceae phylogeny 238
 - spikelets of 530
 - trnL*–F analysis of 535, 539
- Mitochondrial genome
- of Poaceae 248
 - of Triticeae 363, 365, 369
- Mnesithea*
- biogeography of 515
 - New World 505, 515
 - in Panicoideae phylogeny 505, 515
 - photosynthetic pathways in 515
- Mniochloa*, biogeography of 310
- Mnium*, 5.8S rDNA insertion in 86
- Molecular clock analysis, of Bromeliaceae 5, 9–12
- Molineriella*, in Pooideae phylogeny 343, 346–347
- Molinia*
- matK* analysis of 342
 - ndhF* analysis of 342
 - ndhH* analysis of 341–342, 344
 - in Poaceae phylogeny 242–244, 341–342
 - rbcL* analysis of 342
- Monanthochloë*
- biogeography of 593
 - in Chloridoideae phylogeny 566, 570–571, 573, 575, 577, 581, 584
 - ITS analysis of 566, 571
 - ITS and *trnL*–F analysis of, combined 573
 - morphological analysis of 575, 584
 - New World 581, 584, 593
- trnL*–F analysis of 566, 570
- Monanthochloinae
- biogeography of 590, 593
 - in Chloridoideae phylogeny 575, 578, 581–582, 584
 - morphological analysis of 584
 - New World 581–582, 584, 590, 593
- Monermeae, in Pooideae phylogeny 380, 401
- Monocliny 605–614
- Monodia*
- “soft” anatomy of 263
 - spikelets of 263
 - in Triodieae phylogeny 262–263
- Monoecy 605–614
- Monotrema* 7
- Moorochloa*
- New World 505, 507
 - in Panicoideae phylogeny 505, 507
- Morelotia*, photosynthetic pathway in 103, 136
- Morrone, Osvaldo 503–529
- Moth pollination, of Bromeliaceae 13
- Muasya, A. Muthama 72–83, 84–98
- Muhlenbergia*
- amphitropical disjuncts in 584
 - biogeography of 585, 587–588, 590, 593
 - in Chloridoideae phylogeny 566, 570–575, 577, 581, 583–584
 - ITS analysis of 566, 571–572, 577
 - ITS and *trnL*–F analysis of, combined 572–573, 577
 - morphological analysis of 575, 577, 584–585
 - New World 581, 583–585, 587–588, 590, 593
 - as outgroup in Triodieae study 263
 - trnL*–F analysis of 566, 570, 577
- Muhlenbergiinae
- biogeography of 587–590, 593
 - in Chloridoideae phylogeny 581–585
 - morphological analysis of 584–585
 - New World 581–585, 587–590, 593
- Müller, Jochen 380–405
- Munroa*
- biogeography of 587, 589, 593
 - in Chloridoideae phylogeny 566, 570–571, 573–574, 577, 581
 - ITS analysis of 566, 571
 - ITS and *trnL*–F analysis of, combined 573
 - New World 581, 587, 589, 593
 - trnL*–F analysis of 566, 570
- Munroinae
- biogeography of 590, 593
 - in Chloridoideae phylogeny 581–582, 585
 - morphological analysis of 585
 - New World 581–582, 585, 590, 593
- Myriocladus*
- in Arthrostylidiinae (Bambuseae) phylogeny 308–309
 - biogeography of 308–309
 - internode lengths of 308

- NAD-ME photosynthesis
 in Chloridoideae 565, 580
 in *Eragrostis* 595
 in Paniceae 262, 508
 in Panicoideae 504
 in *Panicum* 507
- NADP-ME photosynthesis
 in Andropogoneae 262
 in Arthropogoninae 507
 in Paniceae 262, 508
 in Panicoideae 504
 in Paspalinae 508
 in *Paspalum* 545
 in Poaceae 261
 in Setariinae 508
- Nardeae
 in Poaceae phylogeny 252–253
 in Pooideae phylogeny 345
- Nardurooides*
 in Loliinae phylogeny 380–405
 molecular analysis of 390–392, 414–415
 morphological analysis of 389, 394–395
 morphological-molecular analysis of, combined 397
 relationship with *Festuca* 406–407
- Nardurus*, in Pooideae phylogeny 380
- Nardus*
 ITS analysis of 351, 355–357
matK analysis of 342
ndhF analysis of 341–342, 344
ndhH analysis of 342
 as outgroup in Stipeae study 350–351, 355–357
 in Poaceae phylogeny 242–244, 341–342, 344
rbcL analysis of 342
- Narenga*
 in Andropogoneae phylogeny 532, 535, 538–540
 ITS analysis of 535, 538
 ITS and *trnL*-F analysis of, combined 540
trnL-F analysis of 535, 539
- Nassella*
 ITS analysis of 351, 356–357, 359
matK analysis of 342
ndhF analysis of 342
ndhH analysis of 342
 in Poaceae phylogeny 242–244, 342
rbcL analysis of 342
 in Stipeae phylogeny 350–351, 356–357, 359
- Nastinae. *See* Hickelinae
- Nastus*
 in Bambuseae phylogeny 323, 326–328, 330
 glumes of 330
 hooded ovary of 330
 morphological analysis of 324, 326, 330
 morphology of 316
 rachilla extension of 330
rpl16 analysis of 322, 327
rpl16 and morphological analysis of, combined 328
- Navia*
Aratitiopea lopezii inclusion in 227
 in Bromeliaceae phylogeny 3–4, 14–18, 28
 CAM photosynthesis by 11
 epiphytism of 12
 geographic origin and distribution of 10–11, 18
 leaf and trichome anatomy of 14
matK-rps16 analysis of, combined 28
ndhF analysis of 3–4, 14–18
 pollination of 13
- Navioideae 16
 in Bromeliaceae phylogeny 9, 16–18
 CAM photosynthesis by 11
 epiphytism of 12
 geographic origin and distribution of 10
 leaf and trichome anatomy of 14
 morphology of 16
ndhF analysis of 9, 16–18
 pollination of 13
- ndhF*
 in Andropogoneae analysis 262, 530–544
 in Arthrotyliidiinae analysis 308, 329
 in *Arundoclaytonia* analysis 493–498, 508
 in Bambuseae analysis 315–316, 323–325, 329–330
 in Bromeliaceae analysis 3–26, 28
 in *Bromus* analysis 450–467
 in Cenchrinae analysis 507
 in Centothecoideae + Panicoideae clade analysis 487–502
 in Chloridoideae analysis 342, 490, 494–497, 574
 in Guaduinae analysis 329
 in *Micraira* analysis 497
 in *Paspalum-Thrasya* analysis 546–547
 in Poaceae analysis 238, 240, 242, 248–249, 251–252, 254–255, 259–260, 335–348
 versus *rbcL* 4
- ndhH*
 in Ecdeiocoleaceae analysis 341–344
 in Joinvilleaceae analysis 341–344
 in Poaceae analysis 336–343
- Neeragrostis*
 in Chloridoideae phylogeny 567, 570–571, 573–574, 576
 ITS analysis of 567, 571
 ITS and *trnL*-F analysis of, combined 573
 relationship with *Eragrostis* 595–597, 599–600, 602
trnL-F analysis of 567, 570
- Neesenbeckia*
 in Cyperaceae phylogeny 78
 photosynthetic pathway in 103, 136
rbcL analysis of 78
- Neesiochloa*
 biogeography of 587, 593
 in Chloridoideae phylogeny 581
 New World 581, 587, 593

Nelmesia

- in Abildgaardieae phylogeny 149–150, 160–161
- embryo type of 150
- geographic distribution of 149
- morphology of 150
- photosynthetic pathway in 103, 136

Nemum

- in Abildgaardieae phylogeny 149–151, 160–161
- in Cyperaceae phylogeny 79, 81
- 5.8S rDNA insertion in 87, 94
- embryo type of 150
- geographic distribution of 149
- morphology of 150
- photosynthetic pathway in 103, 136
- rbcL* analysis of 79, 81, 151

Neobouteloua

- biogeography of 587, 593
- in Chloridoideae phylogeny 567, 570–571, 573, 581
- ITS analysis of 567, 571
- ITS and *trnL*-F analysis of, combined 573
- New World 581, 587, 593
- trnL*-F analysis of 567, 570

Neoglaziovia

- in Bromeliaceae phylogeny 36
- trnL* and *trnL*-F analysis of, combined 36

Neoregelia

- in Bromeliaceae phylogeny 8–9, 29, 34–36
- CAM photosynthesis by 11
- epiphytism of 12
- geographic origin and distribution of 10
- leaf and trichome anatomy of 14
- ndhF* analysis of 8–9
- pollination of 13
- trnL*, *trnL*-F, and *trnT*-F analysis of, combined 34
- trnL* and *trnL*-F analysis of, combined 36

Neostapfia

- biogeography of 593
- in Chloridoideae phylogeny 567, 570–571, 573, 577, 581, 585
- ITS analysis of 567, 571
- ITS and *trnL*-F analysis of, combined 573
- morphological analysis of 585
- New World 581, 585, 593
- trnL*-F analysis of 567, 570

Nephelochloa, tribal/subtribal placement of 424

Neurachneae, in Panicoideae phylogeny 503

Neurolepis

- in Bambuseae phylogeny 307–308, 319, 323, 326–329
- biogeography of 307–308
- elevational range of 307
- leaf sizes of 307–308
- morphological analysis of 324, 326
- morphology of 307–308, 316
- papillae of 307
- in Poaceae phylogeny 251, 254
- relationship with *Chusquea* 307–308

rpl16 analysis of 319, 322, 327

rpl16 and morphological analysis of, combined 328

spikelets of 307

Neuropoa, tribal/subtribal placement of 424

New Zealand, *Festuca* of 406–419

Neyraudia

in Chloridoideae phylogeny 581

New World 581

Niche conservatism 224

Nicoraella. See *Anatherostipa*

Nidularium

in Bromeliaceae phylogeny 8–9, 29, 36

CAM photosynthesis by 11

epiphytism of 12

geographic origin and distribution of 10

leaf and trichome anatomy of 14

ndhF analysis of 8–9

pollination of 13

trnL and *trnL*-F analysis of, combined 36

Nidularoid clade 29

Nieuwborg, Roland 72–83

Noll, Jeffrey D. 237–247

North America

Carex subgen. *Vignea* sect. *Ovales* in 194, 199–202

Chloridoideae in 585–590

Panicoideae in 509–511

Poa in 420

Notodanthonia, in Danthonioideae phylogeny 261

Nuclear ribosomal genes. See also *specific genes and taxa*

in Abildgaardieae analysis 151–156, 158–162

in Andropogoneae analysis 262, 530–544

in *Bromus* analysis 450–467

in *Carex* sect. *Ovales* analysis 194

in Cariceae analysis 165–192

in Chloridoideae analysis 565–579, 582

in Cyperaceae analysis 81, 84–98

in *Eleocharis* analysis 95

in Poaceae analysis 248–249, 259–260

in Triodieae analysis 263

Obee, Elizabeth M. and James A. Quinn—Phenotypic plasticity of reproduction in *Schizachyrium scoparium* (Poaceae) populations in relation to ecological history 273–285

Ochagavia

AFLP analysis of 36–37

in Bromeliaceae (Bromelioideae) phylogeny 29, 34–37, 41

geographic origin and distribution of 41

morphology of 41

trnL, *trnL*-F, and *trnT*-F analysis of, combined 34

trnL and *trnL*-F analysis of, combined 35–36

Oja, Tatjana—Preliminary isozyme evidence on the hybrid origin and diploid progenitors of *Bromus pectinatus* (Poaceae) 468–471

Olmea

biogeography of 309

- in Guaduinae (Bambuseae) phylogeny 309
- Olonova, Marina V.—Variability of bluegrass (*Poa*) in Siberia 435–449
- Olyra*
- biogeography of 310–311
 - chromosome numbers of 311
 - dispersal mode of 310
 - matK* analysis of 342
 - ndhF* analysis of 342
 - ndhH* analysis of 342
 - in Olyrinae (Olyreae) phylogeny 310–311
 - phylogeny of 310–311
 - in Poaceae phylogeny 242–245, 342
 - rbcL* analysis of 342
 - synflorescences of 311
- Olyreae
- in Bambusoideae phylogeny 303, 306, 310–311, 315
 - biogeography of 311, 315
 - branch patterns of 310, 315
 - culms of 310, 315
 - dispersal modes of 310
 - in eastern Brazil 311
 - flowers of 315
 - glumes of 310
 - height of 310
 - latitudinal range of 310
 - monophyly of 306, 315
 - New World, classification and biogeography of 310–311
 - phylogeny of 310–311, 323
 - in Poaceae phylogeny 245, 252–253
 - rpl16* analysis of 322
 - spikelets of 310, 315
 - synflorescences of 310
- Olyrinae
- biogeography of 310–311
 - in Olyreae phylogeny 310–311
 - phylogeny of 310–311
 - relationship with *Buergersiochloa* 311
- Ophiochloa*
- biogeography of 509, 511, 523
 - New World 505, 508–509, 511, 523
 - in Panicoideae phylogeny 505, 508, 523
 - photosynthetic pathway of 523
- Oplismenopsis*
- biogeography of 509, 511, 523
 - New World 505, 508–509, 511, 523
 - in Panicoideae phylogeny 505, 508, 523
 - photosynthetic pathway of 523
- Oplismenus*
- New World 505, 508
 - in Panicoideae phylogeny 505, 508
 - in Poaceae phylogeny 251
- Orcuttia*
- biogeography of 593
 - in Chloridoideae phylogeny 567, 570–571, 573, 577, 581, 585
 - ITS analysis of 567, 571
 - ITS and *trnL*-F analysis of, combined 573
 - morphological analysis of 585
 - New World 581, 585, 593
 - trnL*-F analysis of 567, 570
- Orcuttieae
- in Chloridoideae phylogeny 565, 574–575, 577
 - in Poaceae phylogeny 252–253
- Orcuttiinae** 592
- biogeography of 587, 589–590, 593
 - in Chloridoideae phylogeny 577, 581–582, 585
 - morphological analysis of 585, 592
 - New World 581–582, 585, 587, 589–590, 593
- Orectanthe*
- androecium of 230
 - gynoecium of 231
 - inflorescence of 228–233
 - perianth of 228–230
- Oreobambos*, in Poaceae phylogeny 251
- Oreobolopsis*
- in Cyperaceae phylogeny 72, 79
 - photosynthetic pathway in 103, 136
 - rbcL* analysis of 79
- Oreobolus*
- in Cyperaceae phylogeny 78
 - 5.8S rDNA insertion in 86, 88, 94
 - photosynthetic pathway in 103, 136
 - rbcL* analysis of 78
 - relationship with *Oxychloe* 85
- Orlovich, David A. 406–419
- Ornithophily. *See* Bird pollination
- Orthoclada*
- in Centothecoideae + Panicoideae phylogeny 489, 494–498
 - geographic distribution of 489
 - morphological analysis of 494, 498
 - ndhF* analysis of 494–496
 - New World 505–506
 - in Panicoideae phylogeny 505–506
 - rpl16* analysis of 494, 496
 - spikelets of 489
- Orthophytum*
- in Bromeliaceae phylogeny 36
 - trnL* and *trnL*-F analysis of, combined 36
- Oryza*
- BAC studies of 68
 - biogeography of 306
 - 5.8S rDNA insertion in 86, 93
 - in Ehrhartoideae phylogeny 306, 344–345
 - ITS analysis of 351, 355–356
 - matK* analysis of 342
 - morphological analysis of 324, 494
 - ndhF* analysis of 341–342, 344–345, 494–496
 - ndhH* analysis of 342, 344–345

- in *Oryzae* phylogeny 306
- as outgroup in Bambuseae study 319, 322–323, 326–328
- as outgroup in Stipeae study 350–351, 355–356
- plastid genome analysis of 336–337
- in Poaceae phylogeny 242–244, 341–342, 344–345
- polyploidy in 372
- rbcL* analysis of 342
- rpl16* analysis of 322, 494, 496
- 26S rRNA analysis of 93
- Oryzae*
 - biogeography of 306
 - in Ehrhartoideae phylogeny 303, 306
 - phylogeny of 306
 - in Poaceae phylogeny 252–254
 - relationship with Ehrharteae 306
- Oryzinae*
 - biogeography of 306
 - in *Oryzae* (Ehrhartoideae) phylogeny 306
- Oryzoideae*. *See* Ehrhartoideae
- Oryzopsis*
 - ITS analysis of 350–351, 355–356, 359–360
 - matK* analysis of 342
 - morphology of 350
 - ndhF* analysis of 342
 - ndhH* analysis of 342
 - in Poaceae phylogeny 342
 - rbcL* analysis of 342
 - relationship with *Achnatherum* 350
 - relationship with *Piptatherum* 350, 359–360
 - in Stipeae phylogeny 350–351, 355–356, 359–360
- Otachyrium*
 - biogeography of 509, 511, 523
 - New World 505, 508–509, 511, 523
 - in Panicoideae phylogeny 505, 508, 523
 - photosynthetic pathway of 523
- Oatea*
 - in Bambuseae phylogeny 309, 319, 322, 326–328
 - biogeography of 309
 - in Guaduinae phylogeny 309
 - morphological analysis of 324, 326
 - rpl16* analysis of 319, 322, 327
 - rpl16* and morphological analysis of, combined 328
- Oxycaryum*
 - absence of 5.8S rDNA insertion in 87, 94, 96
 - in Cyperaceae phylogeny 80
 - photosynthetic pathway in 103
 - rbcL* analysis of 80
- Oxychloe*
 - in Cyperaceae phylogeny 85, 89–92
 - 5.8S rDNA insertion in 89
 - flower of 55, 60
 - geographic distribution of 55
 - growth form of 55
 - inflorescence of 55
 - in Juncaceae phylogeny 55–56, 58–60, 84, 90–92
 - rbcL* analysis of 85, 89–92
 - rbcL* chimera in 89–92
 - rbcL* contaminant in 89–92
 - rps16* analysis of 59
 - rps16* and *trnL-F* analysis of, combined 90
 - trnL-F* analysis of 57, 59
- Oxytenanthera*
 - in Bambuseae phylogeny 326–328
 - morphological analysis of 324, 326
 - rpl16* analysis of 322, 327
 - rpl16* and morphological analysis of, combined 328
- PACC clade, of Poaceae 248, 335
- PACCAD clade, of Poaceae 237, 239, 241, 251, 254–255, 260–261, 263–264, 303–304, 335, 341–342, 344–345, 487, 495, 565
- PACMAD clade, of Poaceae 241–245
- PAM clade, of Poaceae 424–427
- Panama, Panicoideae in 509–510
- Paniceae
 - base chromosome number of 262, 507
 - biogeography of 509, 511, 516–529
 - in Centothecoideae + Panicoideae phylogeny 487–488, 491–500
 - inflorescences of 508
 - ndhF* analysis of 490–500
 - New World 504–507, 509, 511, 516–529
 - in Panicoideae phylogeny 262, 503–505, 507, 516–529
 - Paspalum-Thrasya-Thrasypsis* relationship 545–562
 - photosynthetic pathways in 262, 504, 516–529
 - in Poaceae phylogeny 245, 250, 252–254
 - relationship with Andropogoneae 262
 - rpl16* analysis of 490–500
 - spikelets of 262, 507–508
 - Thysanolaena* placement in 509
- Panicinae
 - biogeography of 510, 519–521
 - New World 504–508, 519–521
 - in Panicoideae phylogeny 503–505, 507–508, 519–521
 - photosynthetic pathways in 508, 519–521
- Panicoideae
 - Amazonian Region 510–511
 - amphitropical disjuncts in 511
 - Andean Region 510–512
 - Andropogoneae phylogeny 530–544
 - Atlantic Province 511
 - base chromosome number of 503–504
 - Brazilian Region 511
 - Caribbean Region 510–511
 - Central American Province 510–511
 - Chacoan Province 511–512
 - Coatinga Province 511
 - distribution in relation to biogeographical areas 509–511
 - distribution in relation to continents and political units 509
 - economic and ecological importance of 545

- embryos of 503
Galapageian Province 510–511
geographic distribution of 503
Guiana Highlands Region 510
Holarctic Kingdom 510
Kranz syndrome 504, 511–512, 514–529
Madrean Region 510
matK analysis of 342
Mesoamerican 509–510
morphological analysis of 491–492, 494, 503
NADP-ME anatomy of 261
ndhF analysis of 342, 344, 487–502
ndhH analysis of 342
Neotropical Kingdom 510–511
New World, classification and biogeography of 503–529
North American 509–510
North American Atlantic Region 510–511
Paraná Province 511–512
Paspalum-Thrasya-Thrasypsis relationship 545–562
photosynthetic pathways in 504, 511–512, 514–529
phylogeny of 262, 487–491, 499, 503–504
in Poaceae phylogeny 237, 241–243, 248, 250–254, 260–261, 335, 341–342, 344, 487, 503
rbcL analysis of 342
relationship with Centothecoideae 487–504
rpl16 analysis of 487–502
South American 509–511
synapomorphy of 261, 503
Uplands of Central Brazil Province 511–512
West Indian Province 510–511
- Panicum*
amphitropical disjuncts in 511
biogeography of 510–511, 520–521, 523–528
economic and ecological importance of 545
ITS analysis of 536, 538, 567, 571
ITS and *trnL*-F analysis of, combined 540, 573
New World 504–508, 510–511, 520–521, 523–528
as outgroup in Andropogoneae study 533, 536
as outgroup in Chloridoideae study 567
in Panicoideae phylogeny 504–505, 507–508, 520–521, 523–528
photosynthetic pathway of 507, 511–512, 520–521, 523–528
in Poaceae phylogeny 242–245, 256
relationship with Andropogoneae 538–540
relationship with *Paspalum*, *Thrasya*, and *Thrasypsis* 545–548, 550–551
reproductive plasticity of 282–284
trnL-F analysis of 536, 539, 567, 570
- Panicum* sect. *Agrostoides* 508
Panicum sect. *Cordovensia* 508
Panicum sect. *Laxa* 508
Panicum sect. *Lorea* 508
Panicum sect. *Megista* 508
Panicum sect. *Monticola* 508
Panicum sect. *Obtusa* 508
Panicum sect. *Parvifolia* 508
Panicum sect. *Parviglumia* 508
Panicum sect. *Prionitia* 508
Panicum sect. *Stolonifera* 508
Panicum sect. *Tenera* 508
Panicum sect. *Tuerckheimiana* 508
Panicum sect. *Valida* 508
Panicum sect. *Verrucosa* 508
- Pappophoreae
in Chloridoideae phylogeny 565, 575, 583
in Poaceae phylogeny 252–253
- Pappophorum*
biogeography of 587–588, 593
in Chloridoideae phylogeny 567, 570–577, 581
ITS analysis of 567, 571–572
ITS and *trnL*-F analysis of, combined 573, 575–576
matK analysis of 572, 574
morphological analysis of 576
New World 581, 587–588, 593
rbcL analysis of 574
relationship with *Eragrostis* 596–597
rpoC2 analysis of 574
trnL-F analysis of 567, 570, 572
“*Pappostipa*”, in Stipeae phylogeny 350, 360
- Pappophoroideae 580
- Paracolpodium*, tribal/subtribal placement of 424
- Parafestuca*
molecular analysis of 389–392
morphological analysis of 387, 394
morphological-molecular analysis of, combined 397
as outgroup in Loliinae study 382, 387, 389–392, 394, 397
tribal/subtribal placement of 424
- Paramapania*, photosynthetic pathway in 103, 136
- Paraná Province, Panicoideae of 511–512
- Parapholiinae
molecular analysis of 389–392
morphological analysis of 387, 396–399
in Poeae phylogeny 381, 403, 421
in Pooideae phylogeny 346
relationship with Loliinae 380, 382, 387, 389–393, 396–399, 401
- Parapholis*
morphological analysis of 387, 394
morphological-molecular analysis of, combined 397
in Poeae phylogeny 403
in Pooideae phylogeny 343, 346
relationship with Loliinae 382, 387, 394, 397, 401
- Paratheria*
New World 505, 507
in Panicoideae phylogeny 505, 507
- Pariana*
in Bambuseae phylogeny 326–328
in Bambusoideae phylogeny 303
biogeography of 310

- matK* analysis of 342
 morphological analysis of 324, 326, 494
ndhF analysis of 342, 494–496
ndhH analysis of 342
 in Olyreae phylogeny 303, 310
 in Parianinae phylogeny 310
 in Poaceae phylogeny 242–245, 342
 pollination of 310
rbcL analysis of 342
rpl16 analysis of 327, 494, 496
rpl16 and morphological analysis of, combined 328
 Parianeae, in Poaceae phylogeny 252–253, 310
 Parianinae
 biogeography of 310
 inflorescence of 310
 in Olyreae phylogeny 310
 relationship with *Buergersiochloa* 311
 spikelets of 310
 stamen number of 310
 taxonomic problems in 310
 Parnell, John A. N. 55–61
Parodilyra
 biogeography of 311
 in Olyrinae (Olyreae) phylogeny 310–311
 spikelets of 311
Parodiochloa, in *Poa* phylogeny 421, 429, 431
 Partial dicliny 605
Pascopyrum, St genome of 372
Paspalidium
 New World 505, 507–508
 in Panicoideae phylogeny 505, 507–508
 photosynthetic pathway of 528
 Paspalinae
 biogeography of 510, 521–528
 New World 504–508, 510, 521–528
 in Panicoideae phylogeny 504–505, 507–508, 521–528
 photosynthetic pathways in 508, 521–528
 spikelets of 508
Paspalum
 Alma group of 550–552
 amphitropical disjuncts in 511
 axillary penduncles of 553
 biogeography of 509–511, 545
 Bonplandiana group of 556
 in Centothecoideae + Panicoideae phylogeny 494–496
 cladistic analysis of 547–552
 Decumbentes group of 545–562
 Dissecta group of 550–552, 556
 Disticha group of 550–552, 556
 inflorescence of 545–546, 552, 555–556
 leaf anatomy of 545, 552, 555
 lemmas of 545–546, 553–554
 lower glumes of 545–546, 552–554
 lower palea of 545–546, 553
 morphological analysis of 494, 545–562
 ndhF analysis of 494–496, 546–547
 New World 504–506, 508–511
 Notata group of 550–552
 in Paniceae phylogeny 545
 in Panicoideae phylogeny 504–505, 508
 papillae of 553, 555
 pedicels of 555
 photosynthetic pathway of 511, 545
 in Poaceae phylogeny 256
 racemes of 545–546, 552–553, 556
 Racemosa group of 550–552, 556
 rachis of 546, 555–556
 relationship with *Panicum* 545–546
 relationship with *Thrasya* 545–556
 relationship with *Thrasyopsis* 545–547, 549–556
 rpl16 analysis of 494, 496
 rpoC2 analysis of 547
 spikelets of 545–546, 552–555
 upper anthercium of 545, 555
 upper glumes of 546, 553–554
Paspalum subgen. *Ceresia* 546, 550–552, 556
Paspalum subgen. *Harpostachys* 550–556
Patosia
 geographic distribution of 55
 growth form of 55
 inflorescence of 55
 in Juncaceae phylogeny 55
 PCK photosynthesis
 in Chloridoideae 565, 580
 in *Eragrostis* 595
 in Paniceae 262
 in Panicoideae 504
 Pennington, Susan J. 503–529, 580–594
Pennisetum
 biogeography of 510, 517
 in Centothecoideae + Panicoideae phylogeny 494–496
 economic and ecological importance of 545
 morphological analysis of 494
 ndhF analysis of 494–496, 507
 New World 505, 507, 510, 517
 in Panicoideae phylogeny 505, 507, 517
 photosynthetic pathway of 517
 in Poaceae phylogeny 242–245
 rpl16 analysis of 494, 496
Pentarrhaphis, breeding system of 612
Pentaschistis, in Danthonioideae phylogeny 261
Pereilema
 biogeography of 593
 in Chloridoideae phylogeny 567, 570–571, 573, 581, 584–585
 ITS analysis of 567, 571
 ITS and *trnL*-F analysis of, combined 573
 morphological analysis of 584–585
 New World 581, 584–585, 593
 trnL-F analysis of 567, 570
 Perianth. *See also specific taxa*
 of *Aratitiopea lopezii* 228–230

- Peridictyon*
 chloroplast DNA analysis of 366, 368
DMCI analysis of 366–367
EF-G analysis of 366–367
 mitochondrial genome of 366
 morphological analysis of 365
 total evidence (combined) analysis of 367–369
 in Triticeae phylogeny 362, 365–369
- Perrierbambus*
 in Bambuseae phylogeny 326–328
 morphological analysis of 324, 326
rpl16 analysis of 327
rpl16 and morphological analysis of, combined 328
- Peru, Chloridoideae in 586–587
- Perulifera*, in Panicoideae phylogeny 503
- Petersen, Gitte 362–371
- Peterson, Paul M. 450–467
- Peterson, Paul M. et al.—Classification and biogeography of
 New World grasses: Chloridoideae 580–594
- Phaenosperma*
matK analysis of 342
ndhF analysis of 342
ndhH analysis of 342
 in Poaceae phylogeny 239, 242–244, 342
 in Pooideae phylogeny 345–346
rbcL analysis of 342
- Phaenosperrmateae, in Pooideae phylogeny 345–346
- Phaenosperrmatideae, in Poaceae phylogeny 252–253
- Phalaridinae
 as outgroup in *Poa* study 422
 in Pooideae phylogeny 346
- Phanopyrum*
 biogeography of 509–510, 528
 New World 505, 508–510, 528
 in Panicoideae phylogeny 505, 508, 528
 photosynthetic pathway of 528
- Phareae, in Poaceae phylogeny 252–253
- Pharoideae
matK analysis of 342
ndhF analysis of 342, 344
ndhH analysis of 342
 New World, classification and biogeography of 304–
 305, 311
 in Poaceae phylogeny 238–239, 250–255, 260, 303–
 304, 341–342, 344
rbcL analysis of 342
 spikelets of 261
- Pharus*
 biogeography of 305
matK analysis of 342
 morphology of 305
ndhF analysis of 241–242, 248–249, 341–342
ndhH analysis of 341–342
 in Pharoideae phylogeny 304–305
 in Poaceae phylogeny 242–244, 248–249, 341–342
rbcL analysis of 342
- Pheidochloa*, in Poaceae phylogeny 241–245, 251
- Phenotypic plasticity
 definition of 273
 in *Schizachyrium scoparium* 273–285
- Philydrum*, as outgroup in Cyperaceae study 73, 78
- Phippsia*
 relationship with *Poa* 421–422, 427
 tribal/subtribal placement of 424
- Phleum*
 in Poeae phylogeny 421
 in Pooideae phylogeny 343, 346
 relationship with *Poa* 421, 427, 433
 tribal/subtribal placement of 424
- Photosynthesis
 in Centothecoideae + Panicoideae clade 491–492
 in Panicoideae 504, 511–512, 514–529
 in Poaceae 259, 261
- Photosynthesis, C₃ and C₄
 in Abildgaardieae 161
 in Arthrostylideae 161
 in Chloridoideae 565, 580
 in Cyperaceae 99–148, 161
 in Paniceae 262
 in Panicoideae 504, 511–512, 514–529
 in *Paspalum* 545
- Photosynthesis, CAM, in Bromeliaceae 4–5, 11–12, 16, 19–
 20, 28, 41
- Photosynthesis, NAD-ME
 in Chloridoideae 565, 580
 in *Eragrostis* 595
 in Paniceae 262, 508
 in Panicoideae 504
 in *Panicum* 507
- Photosynthesis, NADP-ME
 in Andropogoneae 262
 in Arthropogoninae 507
 in Paniceae 262, 508
 in Panicoideae 504
 in Paspalinae 508
 in *Paspalum* 545
 in Poaceae 261
 in Setariinae 508
- Photosynthesis, PCK
 in Chloridoideae 565, 580
 in *Eragrostis* 595
 in Paniceae 262
 in Panicoideae 504
- Phragmites*
 culms of 325
 morphological analysis of 494
ndhF analysis of 494–496
 in Poaceae phylogeny 242–244
 relationship with *Steyermarkochloa* 508
rpl16 analysis of 494, 496
- PHYB*
 in Andropogoneae analysis 531, 537–543

- in Poaceae analysis 248–249, 252, 255, 487
- Phyllorachideae, in Poaceae phylogeny 252–253
- Phylloscirpus*
- in Cyperaceae phylogeny 79
 - photosynthetic pathway in 103, 136
 - rbcL* analysis of 79
- Phyllostachys*
- in Bambuseae phylogeny 326–328
 - hooded ovary of 330
 - morphological analysis of 324, 326
 - rpl16* analysis of 322, 327
 - rpl16* and morphological analysis of, combined 328
- Pierce, Simon—The jeweled armor of *Tillandsia*—multifaceted or elongated trichomes provide photoprotection 44–52
- Pimentel, Manuel and Elvira Sahuquillo—Relationships among some populations of *Anthoxanthum alpinum* and *A. odoratum* (Poaceae, Pooideae): a morphological/anatomical approach 472–484
- Piptatherum*
- ITS analysis of 351, 355–356, 359–360
 - matK* analysis of 342
 - ndhF* analysis of 342
 - ndhH* analysis of 342
 - in Poaceae phylogeny 242–244, 342
 - rbcL* analysis of 342
 - relationship with *Oryzopsis* 350, 359–360
 - in Stipeae phylogeny 350–351, 355–356, 359–360
- Piptochaetium*
- ITS analysis of 351, 356, 360
 - in Stipeae phylogeny 350–351, 356, 360
- Piresiella*, biogeography of 310
- Pitcairnia*
- in Bromeliaceae phylogeny 3–4, 7–9, 14–17, 28, 34–36, 41
 - CAM photosynthesis by 11–12
 - epiphytism of 12
 - geographic origin and distribution of 10–11, 18–19, 35
 - growth form of 22
 - leaf and trichome anatomy of 13–14
 - matK-rps16* analysis of, combined 28
 - ndhF* analysis of 3–4, 7–9, 14–17
 - pollination of 13, 22
 - trnL*, *trnL-F*, and *trnT-F* analysis of, combined 34
 - trnL* and *trnL-F* analysis of, combined 36
- Pitcairnia* subgen. *Pepinia* 15
- Pitcairnia* subgen. *Pitcairnia* 15
- Pitcairnioideae 16
- as ancestral Bromeliaceae 18–19
 - in Bromeliaceae phylogeny 3–4, 7–9, 15–17, 28, 34–35, 39–40
 - CAM photosynthesis by 11–12
 - epiphytism of 12
 - geographic origin and distribution of 10–12, 18–19
 - leaf and trichome anatomy of 5, 13–14
 - morphology of 16
 - ndhF* analysis of 3–4, 7–9, 15–17
 - paraphyly of 8, 15, 39
 - pollination of 12–13, 22
 - recircumscription of 15, 17
 - trnL* analysis of 28
 - winged seeds of 3
- Plagiantha*
- biogeography of 509, 511, 528
 - New World 505, 508–509, 511, 528
 - in Panicoideae phylogeny 505, 508, 528
 - photosynthetic pathway of 528
- Plasticity, phenotypic 273
- Plasticity, reproductive
- in *Aegilops triuncialis* invasion 295–300
 - chasmogamous and cleistogamous 286–294
 - of *Panicum* 282–284
 - of *Schizachyrium scoparium* 273–285
 - of *Sporobolus* 282–284
- Plastid genome analysis. *See also specific genes and taxa*
- of Andropogoneae 262
 - of Bambuseae 316
 - of *Bromus* 450–467
 - of Cyperaceae 55–61, 72–83
 - of Ecdeiocoleaceae 336–337
 - of Joinvilleaceae 336–337
 - of Juncaceae 55–61
 - of Poaceae 237–238, 240, 242, 245, 248–249, 254–255, 259–260, 335–348
 - of Poales 336
 - of Pooideae 335–348
- Plectrachne*
- spikelets of 262
 - in Triodieae phylogeny 262–263
- Pleuraphis*
- biogeography of 593
 - breeding system of 608–610
 - in Chloridoideae phylogeny 567, 570–571, 573, 580–581, 584
 - ITS analysis of 567, 571
 - ITS and *trnL-F* analysis of, combined 573
 - morphological analysis of 584
 - New World 581, 584, 593
 - trnL-F* analysis of 567, 570
- Pleuropogon*
- matK* analysis of 342
 - ndhF* analysis of 342
 - ndhH* analysis of 341–342, 344
 - in Poaceae phylogeny 341–342
 - rbcL* analysis of 342
- Pleurostachys*
- in Cyperaceae phylogeny 77–78
 - 5.8S rDNA insertion in 88, 94
 - photosynthetic pathway in 101, 103, 136
 - rbcL* analysis of 77–78

Poa

- Anthochloa* inclusion in 421, 429–431
- ArcSyl clade of 421, 428–429
- “Australopoa” group of 421, 430, 432
- BAPO clade of 421, 427–429, 432
- chloroplast DNA analysis of 420–433
- cladistic analysis of 421, 424–431
- Dissanthelium* inclusion in 421, 429–432
- Edmond classification of 420
- Eremopoa* inclusion in 420–422, 424, 427, 429
- genera allied with 421–424, 426–427, 431–432
- generic relationships and affinities of 432–433
- HAMBADD clade of 421, 429–430, 432
- height of 437, 443
- hybridization of 420–421
- infrageneric classifications of 432
- leaf sheaths of 438–439
- lemmas of 437–438, 442–447
- ligules of 437–438, 442–447
- molecular analysis of 390–392, 414–415
- morphological analysis of 387, 394, 421, 435–449
- morphological-molecular analysis of, combined 397, 421
- New World 420
- in New Zealand 418
- North American 420
- as outgroup in Loliinae study 382, 387, 390–394, 397
- panicles of 437–439
- phylogeny of 420–434
- Poa* (main) clade of 424
- in Poaceae phylogeny 256
- in Poeae phylogeny 380
- polyploidy in 420
- PoM clade of 421, 429–430, 432
- in Pooideae phylogeny 343, 380
- POSTA clade of 428
- “Punapoa” group of 421, 430, 432
- rachillas of 438, 442–447
- relationship with Alopecurinae 421–427, 431–433
- relationship with *Festuca* 407–408, 421–422
- relationship with Miliinae 421–427, 431–433
- in Siberia, variability of 435–449
- Soreng classification of 420
- spikelets of 438–440
- SPOSTA clade of 421, 428–430, 432
- tribal/subtribal placement of 424
- trnT*-F analysis of 420–434
- Tzvelev classification of 420
- Poa lepidula*** 431
- Poa* sect. *Bulbosae* 435
- Poa* sect. *Coenopoa* 435
- Poa* sect. *Malacanthae* 435, 437
- Poa* subgen. *Andinae* 420–421, 424–425, 427, 431, 433
- Poa* subgen. *Arctopoa* 420–421, 425, 428, 432, 435
- Poa* subgen. *Arctopoa* sect. *Aphydris* 432
- Poa* subgen. *Arctopoa* sect. *Arctopoa* 432, 435
- Poa* subgen. *Arctopoa* sect. *Sylvestres* 432
- Poa* subgen. *Dioicopoa* 420
- Poa* subgen. *Ochlopoa* 432
- Poa* subgen. *Ochlopoa* sect. *Alpinae* 432
- Poa* subgen. *Ochlopoa* sect. *Arenariae* 432
- Poa* subgen. *Ochlopoa* sect. *Micrantherae* 432
- Poa* subgen. *Ochlopoa* sect. *Parodiochloa* 432
- Poa* subgen. *Poa* 420–421, 425–426, 432, 435
- Poa* subgen. *Poa* sect. *Abbreviatae* 421, 428, 435
- Poa* subgen. *Poa* sect. *Acutifoliae* 421, 430, 432
- Poa* subgen. *Poa* sect. *Alpinae* 421, 428, 435
- Poa* subgen. *Poa* sect. ***Anthochloa*** 431–432
- Poa* subgen. *Poa* sect. *Austrofestuca* 421, 430, 432
- Poa* subgen. *Poa* sect. *Bolbophorum* 421, 428
- Poa* subgen. *Poa* sect. *Brizoides* 421, 430, 432
- Poa* subgen. *Poa* sect. *Dasypoa* 421, 430, 432
- Poa* subgen. *Poa* sect. *Dioicopoa* 420–421, 430, 432
- Poa* subgen. *Poa* sect. *Homalopoa* 421, 430, 432, 435
- Poa* subgen. *Poa* sect. *Leptophyllae* 430, 432
- Poa* subgen. *Poa* sect. *Macropoa* 421, 430, 432, 435–437
- Poa* subgen. *Poa* sect. *Madropoa* 421, 430, 432
- Poa* subgen. *Poa* sect. *Nivicolae* 430, 432, 435
- Poa* subgen. *Poa* sect. *Ochlopoa* 420–421, 428, 435
- Poa* subgen. *Poa* sect. *Oreinos* 421, 428–430, 435
- Poa* subgen. *Poa* sect. *Pandemos* 421
- Poa* subgen. *Poa* sect. *Parodiochloa* 421, 428
- Poa* subgen. *Poa* sect. *Poa* 421, 430, 432, 438–440
- Poa* subgen. *Poa* sect. *Secundae* 421, 428–430
- Poa* subgen. *Poa* sect. *Stenopoa* 421, 428–430, 435, 440–448
- Poa* subgen. *Poa* sect. *Sylvestres* 421, 428
- Poa* subgen. *Poa* sect. *Tichopoa* 421, 428
- Poa* subgen. *Poa* supersect. ***Homalopoa*** 432
- Poa* subgen. *Poa* supersect. *Poa* 432
- Poa* subgen. *Pseudopoa* 420, 432
- Poa* subgen. *Pseudopoa* sect. *Pseudopoa* 432
- Poa* subgen. *Psilantha* 420
- Poa* subgen. ***Stenopoa*** 432
- Poa* subgen. ***Stenopoa*** sect. *Abbreviatae* 432
- Poa* subgen. ***Stenopoa*** sect. *Oreinos* 432
- Poa* subgen. ***Stenopoa*** sect. *Pandemos* 432
- Poa* subgen. ***Stenopoa*** sect. *Secundae* 432
- Poa* subgen. ***Stenopoa*** sect. *Stenopoa* 432
- Poa* subgen. ***Stenopoa*** sect. *Tichopoa* 432
- Poaceae
 - adh1* analysis of 252
 - Andropogoneae phylogeny 530–544
 - Anisopogon/Phaenosperma* clade of 239
 - Anthoxanthum alpinum* and *A. odoratum* relationships 472–484
 - atpB* analysis of 248
 - Bambusa/Chusquea* clade of 239
 - Bambuseae one-flowered, determinate genera phylogeny 315–332
 - BEP clade of 237, 239, 241, 246, 248, 255, 260, 263–264, 303–304, 335, 341–342, 345, 487

- Bouteloua* dicliny 605–614
Bromus pectinatus origin and progenitors 468–471
Bromus phylogeny 450–467
Bromus position in 242–244, 450
 Centothecoideae + Panicoideae clade of 487–502
 chasmogamy/cleistogamy in 286–294
 Chloridoideae phylogeny 565–579
 chloroplast DNA analysis of 248
 cladistic analyses of 237, 239, 241–246, 248, 251, 254–255, 259–261, 263–264, 303–304, 335, 487–502
 conglomerate taxa of 238, 337
 culm of 261, 286
 DELTA system of 267–268
 descriptive database of 267–269
 diversification of 248–258
 “divide and conquer” strategy for 249, 254
 18S rDNA analysis of 252
 early diverging (basal) 238–239, 252–253, 255, 260, 303, 344
 economic and ecological importance of 545
 electronic world database of 267–270
 embryo of 261
Eragrostis monophyly and infrageneric classification 595–604
 European Molecular Biology Laboratory data on 249
 evolutionary history of 263, 589–590
 evolutionary success of 255
 fruit of 261
 future study directions for 256
GBSSI analysis of 248–249, 252, 260, 337, 343–344
 GPWG classification of 237, 246, 260–261
 homoplasmy in 260–261
 inflorescence of 259, 261
 ITS analysis of 248–249, 251–254, 259–260
 large trees of 248–258
 leaf anatomy of 259, 261
 Loliinae phylogeny 380–405
matK analysis of 241, 248, 252, 254, 259–260, 337–343
 mitochondrial genome of 248
 morphological versus molecular classification of 259–266
 multigene analysis of 249
ndhF analysis of 238, 240–242, 248–249, 251–252, 254–255, 259–260, 336–347
ndhH analysis of 336–343
 New Systematics on 259
 New World grasses, classification and biogeography of 303–314
 New World Chloridoideae, classification and biogeography of 580–594
 New World Panicoideae, classification and biogeography of 503–529
 New Zealand *Festuca*, phylogeny and biogeography of 406–419
 nomenclature change in, conservatism in 264
 nuclear ribosomal analysis of 248–249, 252–254, 259–260
 outgroup effects on 239, 245–246
 PACC clade of 248, 335
 PACCAD, B-E, P clade of 255
 PACCAD clade of 237, 239, 241, 251, 254–255, 260–261, 263–264, 303–304, 335, 341–342, 344–345, 487, 565
 PACCAD-P, B-E group of 251, 255
 PACCAD-P group of 251, 254
 PACMAD clade of 241–245
 paradigm shift on 259
Paspalum-Thrasya-Thrasypsis relationship 545–562
 phenetic classification of 259, 263
 photosynthetic pathways in 259, 261
PHYB analysis of 248–249, 252, 255, 487
 phylogeny of 237–266, 303–304, 315, 335–348
 phylogeny of, versus classification 263, 381
 plastid genome analysis of 237–238, 240–242, 245, 248–249, 254–255, 259–260, 335–348
Poa phylogeny 420–434
Poa variability in Siberia 435–449
 Pooideae phylogeny 335–348
rbcL analysis of 238, 240–242, 245, 248–249, 252, 255, 259–260, 337–343
 relationship with Bromeliaceae 4, 8, 27–28
 relationship with Ecdiaceae 237–238
 relationship with Joinvilleaceae 237–238, 260
 reproductive plasticity in *Aegilops triuncialis* invasion 295–300
 reproductive plasticity of *Schizachyrium scoparium* 273–285
 ribosomal RNA analysis of 259
rp116 analysis of 248, 259–260
rpoC2 analysis of 241–245, 248–249, 252, 254, 259–260
rps4 analysis of 248, 259–260
rps15 analysis of 336
 sample size of 263–264
 seedlings of 261
 self-fertilization in 286
 sheath fertilization in 286
 socioeconomic and ecological importance of 248
 species richness of 256
 spikelet clade of 252, 255
 spikelets of 259, 261, 286
 Stipeae molecular analysis 349–361
 subfamilies of 237, 250–253, 259–260, 263
 supertrees of 248–258
 synapomorphies of 260–261
 synonym database of 267–269
 tribes of 252–253
 Triticeae phylogeny 362–371
trnL analysis of 248, 252, 254
trnL-F analysis of 254

- trnT* inversion in 237–238
- Poales
- Bromeliaceae placement in 4, 19, 27–28
 - Juncaceae placement in 55
 - phylogeny of 227
 - plastid genome analysis of 336
 - relationship with Cyperales 64
- Poeae
- morphology of 380
 - PAM clade of 424–427
 - phylogeny of 380, 401–403, 421
 - in Poaceae phylogeny 251–253, 255–256
 - in Pooideae phylogeny 335, 345–347, 380–381
- Poeae s.l. 335, 343, 346–347
- Pogonarthria*
- in Chloridoideae phylogeny 567, 570–571, 573–574, 576, 581, 583
 - ITS analysis of 567, 571
 - ITS and *trnL*-F analysis of, combined 573
 - morphological analysis of 583
 - New World 581, 583
 - relationship with *Eragrostis* 595, 599–600, 602
 - trnL*-F analysis of 567, 570
- Pogonatherum*
- New World 504–505
 - in Panicoideae phylogeny 504–505
 - in Poaceae phylogeny 251
- Pohlidium*
- biogeography of 509–510, 516
 - in Centothecoideae + Panicoideae phylogeny 489, 494–498
 - geographic distribution of 489
 - morphological analysis of 494, 498–499
 - ndhF* analysis of 494–496
 - New World 505–506, 509–510, 516
 - in Panicoideae phylogeny 505–506, 516
 - photosynthetic pathway of 516
 - rpl16* analysis of 494, 496
 - spikelets of 489, 498–499
- Poidium*, tribal/subtribal placement of 424
- Poinae
- placement of genera in 423–424
 - as outgroup in Loliinae study 382
 - in Poeae (Pooideae) phylogeny 380, 421
 - in Pooideae phylogeny 346
 - relationship with Alopecurinae and Miliinae 421–427, 431–433
- Pollination
- of Bromeliaceae 5, 11–13, 21–22
 - of Cyperaceae 77
- Pollinia*, in Andropogoneae phylogeny 532
- Polyploidy, in *Elymus* 372–379
- Polypogon*, in Pooideae phylogeny 343, 346
- Polytrias*
- New World 504–505
 - in Panicoideae phylogeny 504–505
- PoM clade, of *Poa* 421, 429–430, 432
- Pommereullinae, in Chloridoideae phylogeny 574–575
- Pooideae
- Anthoxanthum alpinum* and *A. odoratum* relationships 472–484
 - Bromus pectinatus* origin and progenitors 468–471
 - Bromus* phylogeny 450–467
 - Bromus* position in 343, 345, 450
 - BT/P clade of 345–346
 - Elymus* allopolyploids 372–379
 - florets of 380
 - GBSSI* analysis of 343
 - glumes of 380
 - lemmas of 380
 - Loliinae phylogeny 380–405
 - matK* analysis of 342
 - monophyly of 335
 - morphology of 380
 - ndhF* analysis of 242, 248–249, 341–344, 346, 490
 - ndhH* analysis of 342
 - New Zealand *Festuca*, phylogeny and biogeography of 406–419
 - phylogeny of 335–348, 380
 - plastid genome analysis of 335–348
 - Poa* phylogeny 420–434
 - Poa* variability in Siberia 435–449
 - in Poaceae phylogeny 237, 242–243, 248–253, 255, 260, 335, 342, 345
 - rbcL* analysis of 342
 - relationship with Ehrhartoideae 305
 - rpl16* analysis of 490
 - spikelets of 380
 - Stipeae molecular analysis 349–361
- Portea*
- in Bromeliaceae (Bromelioideae) phylogeny 36, 41
 - leaf blade anatomy of 39
 - trnL* and *trnL*-F analysis of, combined 36
- POSTA clade, of *Poa* 428
- Potarophytum* 7
- Principina*, photosynthetic pathway in 103, 136
- Prioniaceae
- in Cyperales phylogeny 55
 - flowers of 55–56
 - phylogeny of 55
 - in Poales phylogeny 55
 - relationship with Juncaceae 55–56
- Prionium*
- absence of 5.8S rDNA insertion in 93–95
 - geographic distribution of 55
 - habitat of 55
 - as outgroup in Cyperaceae study 73, 78, 85
 - in Prioniaceae phylogeny 55–56
 - relationship with Cyperaceae 56, 58, 60
 - relationship with Juncaceae 55–56, 58, 60
 - rps16* analysis of 58
 - in Thurniaceae phylogeny 55–56, 58, 60

- trnL*-F analysis of 58
- Psathyrostachys*
chloroplast DNA analysis of 366, 368
DMC1 analysis of 366–367, 370
EF-G analysis of 366–367, 370
mitochondrial genome of 366
morphological analysis of 363, 365–366
Ns genome of 372
total evidence (combined) analysis of 367–369
in Triticeae phylogeny 362–363, 365–370, 374
- Pseudechinolaena*
New World 505, 508
in Panicoideae phylogeny 505, 508
- Pseudochaetochloa*, in Poaceae phylogeny 254
- Pseudodanthonia*, relationship with *Bromus* 450, 463
- Pseudoroegneria*
chloroplast DNA analysis of 366, 368–369
DMC1 analysis of 366–367, 370
EF-G analysis of 366–367, 370
genome transfer to *Elymus* 372, 374–378
mitochondrial genome of 366
morphological analysis of 365
total evidence (combined) analysis of 367–369
in Triticeae phylogeny 365–370
- Pseudosasa*
matK analysis of 342
ndhF analysis of 342
ndhH analysis of 342
Old World taxa, in New World 309
in Poaceae phylogeny 238, 342
rbcL analysis of 342
- Pseudoschoenus*, photosynthetic pathway in 103, 136
- Pseudosorghum*, in Andropogoneae phylogeny 541
- Pseudozoysia*, in Poaceae phylogeny 251, 254
- Psilureae 402
- Psilurinae, in Poeae (Pooideae) phylogeny 380
- Psilurus*
in Loliinae phylogeny 380–405
molecular analysis of 386, 389–392, 414–415
morphological analysis of 389, 394–399
morphological-molecular analysis of, combined 397–398
relationship with *Festuca* 406–407, 413
- Ptilostipa*, in Stipeae phylogeny 350
- Ptilothrix*, photosynthetic pathway in 103, 136
- Puccinellia*
molecular analysis of 390–392
morphological analysis of 387, 394
morphological-molecular analysis of, combined 397
in New Zealand 418
as outgroup in Loliinae study 382, 387, 390–392, 394, 397
in Poeae phylogeny 381, 421
in Pooideae phylogeny 343, 346
relationship with *Poa* 421–422, 433
tribal/subtribal placement of 424
- Puccinelliinae
placement of genera in 423–424
in Poeae phylogeny 421
in Pooideae phylogeny 346
relationship with *Poa* 421–422, 427
- Puelia*
matK analysis of 342
ndhF analysis of 342
ndhH analysis of 342
in Poaceae phylogeny 242–244, 251, 342
in Puelioideae phylogeny 303
rbcL analysis of 342
- Puelieae, in Poaceae phylogeny 252–253
- Puelioideae
matK analysis of 342
ndhF analysis of 342, 344
ndhH analysis of 342
phylogeny of 303
in Poaceae phylogeny 238–239, 242–243, 250–251, 254–255, 260, 303, 341–342, 344
rbcL analysis of 342
- “Punapoa” group, of *Poa* 421, 430, 432
- Puya*
in Bromeliaceae phylogeny 3, 7–9, 16, 28, 34–36, 40–41
CAM photosynthesis by 11, 19, 41
concerted convergence in 22
epiphytism of 12
flowers of 13
geographic origin and distribution of 10, 12, 18–19
growth form of 22
leaf and trichome anatomy of 13–14, 20, 22
morphology of 41
narrow endemism in 21
ndhF analysis of 3, 7–9, 16
pollination of 13, 22
relationship with Bromelioideae 40–41
species diversity of 21
trnL, *trnL*-F, and *trnT*-F analysis of, combined 34
trnL analysis of 28
trnL and *trnL*-F analysis of, combined 36
as ur-bromeliad 19
- Puya* subgen. *Puya* 12
- Puya* subgen. *Puyopsis* 12
- Puyeae, in Bromeliaceae phylogeny 35
- Puyoideae 16**
in Bromeliaceae phylogeny 9, 16–17
CAM photosynthesis by 11
epiphytism of 12
geographic origin and distribution of 10
leaf and trichome anatomy of 14
morphology of 16
ndhF analysis of 9, 16–17
pollination of 12–13
- Pycreus*
absence of 5.8S rDNA insertion in 87, 94

- in Cyperaceae phylogeny 80
- photosynthesis in, conflicting data on 148
- photosynthetic pathway in 103, 136–138
- rbcL* analysis of 80
- Queenslandiella*
 - in Cyperaceae phylogeny 80
 - photosynthetic pathway in 103, 138
 - rbcL* analysis of 80
- Quesnelia*
 - in Bromeliaceae phylogeny 36
 - trnL* and *trnL*–*F* analysis of, combined 36
- Quinn, James A. 273–285
- Racemobambos*, in Bambuseae phylogeny 329
- Racinaea*, in Bromeliaceae phylogeny 28
- Raddia*
 - biogeography of 311
 - glumes of 311
 - in Olyrinae (Olyreae) phylogeny 311
- Raddiella*, biogeography of 310
- Rames, of Andropogoneae 262
- Ramirea*
 - in Cyperaceae phylogeny 80
 - rbcL* analysis of 80
- Rapatea*
 - as outgroup in Bromeliaceae (Bromelioideae) study 33
 - trnL*, *trnL*–*F*, and *trnT*–*F* analysis of, combined 34
- Rapateaceae
 - historical biogeography of 19–20
 - relationship with Bromeliaceae 4, 8, 19, 27
- rbcL*
 - in Abildgaardieae analysis 73, 79, 81, 149–151
 - in Andropogoneae analysis 533
 - in Bambuseae analysis 330
 - in Bromeliaceae analysis 19, 28
 - in Chloridoideae analysis 342, 574
 - in Cyperaceae analysis 72–85
 - in *Eriachne* analysis 497
 - versus *ndhF* 4
 - in *Oxychloe* analysis 85, 89–92
 - in Poaceae analysis 240, 242, 245, 248–249, 252, 255, 259–260, 337–343
 - in *Rhodocoma* analysis 213–214, 217–218
 - in *Tristachya* analysis 497, 505, 533, 541
 - in Triticeae analysis 363–366
- Redfieldia*
 - biogeography of 593
 - in Chloridoideae phylogeny 567, 570–571, 573, 577, 581, 584–585
 - ITS analysis of 567, 571
 - ITS and *trnL*–*F* analysis of, combined 573
 - morphological analysis of 577, 584–585
 - New World 581, 584–585, 593
 - trnL*–*F* analysis of 567, 570
- Reederochloa*
 - biogeography of 593
- in Chloridoideae phylogeny 567, 570–571, 573, 575, 577, 581, 584
- ITS analysis of 567, 571
- ITS and *trnL*–*F* analysis of, combined 573
- morphological analysis of 584
- New World 581, 584, 593
- trnL*–*F* analysis of 567, 570
- Reedia*, photosynthetic pathway in 103, 138
- Refulio-Rodriguez, Nancy F. 565–579
- Rehia*, biogeography of 310
- Reimarochloa*
 - biogeography of 509, 528
 - New World 505, 508–509, 528
 - in Panicoideae phylogeny 505, 508, 528
 - photosynthetic pathway of 528
- Renvoize, Stephen A. 248–258
- Renvoize, Steve et al.—An electronic world grass flora 267–270
- Reproductive isolation, chromosome evolution and 62
- Reproductive plasticity
 - in *Aegilops triuncialis* invasion 295–300
 - chasmogamous and cleistogamous 286–294
 - of *Panicum* 282–284
 - of *Schizachyrium scoparium* 273–285
 - of *Sporobolus* 282–284
- Reproductive systems
 - in *Bouteloua* 605–614
 - evolution of dioecy 605–614
 - phylogenetic study of 606
- Restio*, as outgroup in Poaceae study 241
- Restionaceae
 - phylogeny and historical evolution of *Rhodocoma* 213–226
 - plastid genome analysis of 336
 - relationship with Bromeliaceae 27–28
 - relationship with Poaceae 255
- Reynaudia*
 - biogeography of 509–510, 517
 - New World 505, 507, 509–510, 517
 - in Panicoideae phylogeny 503, 505, 507, 517
 - photosynthetic pathway of 517
- Reznicek, Anton A. 72–83, 193–203
- Rheochloa*
 - biogeography of 587, 593
 - in Chloridoideae phylogeny 565–567, 570–571, 573, 581
 - New World 565–567, 581, 587, 593
- Rhipidium, possible presence in *Schoenus* 204–209
- Rhipidocladum*
 - in Andes 311
 - in Arthrostylidiinae (Bambuseae) phylogeny 308–309
 - biogeography of 308–309, 311
 - branching of 308
 - buds of 308
 - internode lengths of 308
 - in Poaceae phylogeny 251, 254

- Rhizocephalus*, tribal/subtribal placement of 424
- Rhizomes
 of Arthrostylidiinae 308
 of Bambuseae 306, 315, 320
 of Hickelinae 330
 of Juncaceae 55
- Rhodocoma*
 altitudinal variation among 215–216
 ancestral ecology reconstructions in 217, 220–221
atpB analysis of 213–214, 217–218
 bedrock-derived soils for 215–216
 ecological, genetic, and phylogenetic distances among species of 217
 ecological parameters in study of 215–216
 ecological similarity among species of 216–217
 fire survival mode of 215–216
 flowering times of 223–224
 groundwater availability for 215–216
 macroecology of 215–216, 218–220
matK analysis of 213–214, 218
 phylogeny and historical evolution of 213–226
 rainfall patterns for 215–216
rbcL analysis of 213–214, 217–218
 relationship with *Thamnochortus* 213, 218, 222
 soil rockiness for 215–216
 sympatric versus parapatric speciation in 213, 223–224
trnK analysis of 213–214
trnL analysis of 213–214
trnL-F analysis of 213–214, 217
- Rhombolytrum*, tribal/subtribal placement of 424
- Rhynchelytrum*, in Panicoideae phylogeny 503
- Rhynchocladium*, photosynthetic pathway in 103, 138
- Rhynchoryza*, in Oryzinae (Oryzeae) phylogeny 306
- Rhynchospora*
 agmatoploidy in 193
 BAC studies of 68
 chromosome evolution in 193
 in Cyperaceae phylogeny 58, 77–78
 5.8S rDNA insertion in 88, 92, 94
 genome size of 66
 photosynthesis in, conflicting data on 148
 photosynthetic pathways in (C₃ and C₄) 101, 103–105, 139–142
 polyploidy in 193
rbcL analysis of 77–78
rps16 analysis of 57–58
Syntrinema inclusion in 104
trnL-F analysis of 57–58
- Rhynchosporeae, inflorescence (spikelet) of 204
- Rhytachne*
 biogeography of 515
 New World 505, 515
 in Panicoideae phylogeny 505, 515
 photosynthetic pathway of 515
- Ribosomal RNA, in Poaceae analysis 259
- Roalson, Eric H. 72–83
- Roalson, Eric H. et al.—Chromosome evolution in Cyperales 62–71
- Ronnbergia*
 in Bromeliaceae phylogeny 36
trnL and *trnL*-F analysis of, combined 36
- Rostkovia*
 5.8S rDNA insertion in 89
 flower of 55, 60
 genome size of 66
 inflorescence of 55
 in Juncaceae phylogeny 55, 58–60
 relationship with *Oxychloe* 90
rps16 analysis of 57, 59
trnL-F analysis of 57, 59
- Rostraria*
ndhF analysis of 341
 in Poaceae phylogeny 341
 in Pooideae phylogeny 343, 346
- Rothrock, Paul E. 193–203
- Rottboellia*
 in Andropogoneae phylogeny 530, 532, 535, 538–540, 542
 ITS analysis of 535, 538, 542
 ITS and *trnL*-F analysis of, combined 540
 New World 504–505
 in Panicoideae phylogeny 504–505
trnL-F analysis of 535, 539
- Rottboellieae, in Andropogoneae phylogeny 532
- Rottboelliinae
 in Andropogoneae phylogeny 262, 531–532, 538–540, 542
 biogeography of 515
 ITS analysis of 538
 ITS and *trnL*-F analysis of, combined 540
 New World 504–506, 515
 in Panicoideae phylogeny 503–505, 515
 photosynthetic pathways in 515
trnL-F analysis of 539
- Royal Botanic Gardens, Kew, electronic world grass database of 267–270
- rpl16*
 in Arthrostylidiinae analysis 308
 in Bambuseae analysis 315–316, 319–330
 in Centothecoideae + Panicoideae clade analysis 487–502
 in Chusqueinae analysis 316
 in Poaceae analysis 248, 259–260
- rpoA*
 in *Elymus* analysis 373–375
 in Triticeae analysis 363–366
- rpoC2*
 in Chloridoideae analysis 574
 in *Paspalum-Thrasya* analysis 547
 in Poaceae analysis 241–245, 248–249, 252, 254, 259–260
- rps4*, in Poaceae analysis 248, 259–260

- rps15*, in Poaceae analysis 336
- rps16*
- in Bromeliaceae analysis 4, 28
 - in Chloridoideae analysis 572–574, 576
 - in Cyperaceae analysis 56–60, 81
 - in *Eragrostis* analysis 597–602
 - in Juncaceae analysis 55–61
 - in *Oxychloe* analysis 59, 90
 - in Thurniaceae analysis 58
- Rytidosperma*, in Danthonioideae phylogeny 261
- Saarela, Jeffery M. et al.—Molecular phylogenetics of *Bromus* (Poaceae: Pooideae) based on chloroplast and nuclear DNA sequence data 450–467
- Sabin-Badereau, Alexandru 349–361
- Sacchareae, in Andropogoneae phylogeny 532
- Saccharinae
- in Andropogoneae phylogeny 262, 532, 538–540, 542–543
 - biogeography of 515
 - ITS analysis of 538
 - ITS and *trnL*-F analysis of, combined 540
 - New World 504–506, 515
 - in Panicoideae phylogeny 503–505, 515
 - photosynthetic pathways in 515
 - trnL*-F analysis of 539
- Saccharum*
- in Andropogoneae phylogeny 530, 532, 535, 538–540, 542–543
 - biogeography of 515
 - economic and ecological importance of 530, 545
 - hybridization of 543
 - ITS analysis of 535, 538
 - ITS and *trnL*-F analysis of, combined 540, 542–543
 - matK* analysis of 342
 - ndhF* analysis of 342
 - ndhH* analysis of 342
 - New World 505, 515
 - in Panicoideae phylogeny 505, 515
 - photosynthetic pathway of 515
 - in Poaceae phylogeny 341–342
 - rbcL* analysis of 342
 - spikelets of 530
 - trnL*-F analysis of 535, 539
- Sacciolepis*
- biogeography of 521
 - New World 505, 508, 521
 - in Panicoideae phylogeny 505, 508, 521
 - photosynthetic pathway of 521
- Sahuquillo, Elvira 472–484
- Salamin, Nicolas 248–258
- Sánchez-Ken, J. Gabriel 237–247, 315–332
- Sánchez-Ken, J. Gabriel and Lynn G. Clark—Phylogenetic relationships within the Centothecoideae + Panicoideae clade (Poaceae) based on *ndhF* and *rpl16* intron sequences and structural data 487–502
- Sarga*, in Andropogoneae phylogeny 262
- Saugetia*
- biogeography of 589, 593
 - in Chloridoideae phylogeny 565–567, 570–571, 573, 581
 - New World 565–567, 581, 589, 593
- Savolainen, Vincent 248–258
- Saxofridericia* 7
- Schaffnerella*
- biogeography of 593
 - in Chloridoideae phylogeny 567, 570–571, 573–574, 577, 581, 584–585
 - ITS analysis of 567, 571
 - ITS and *trnL*-F analysis of, combined 573
 - morphological analysis of 584–585
 - New World 581, 584–585, 593
 - trnL*-F analysis of 567, 570
- Schedonnardus*
- biogeography of 588, 593
 - in Chloridoideae phylogeny 567, 570–571, 573–575, 577, 581, 584–585
 - ITS analysis of 567, 571
 - ITS and *trnL*-F analysis of, combined 573
 - morphological analysis of 575, 584–585
 - New World 581, 584–585, 588, 593
 - trnL*-F analysis of 567, 570
- Schedonorus*
- in Loliinae phylogeny 381, 400
 - molecular analysis of 414–415
 - relationship with *Festuca* 381, 406–413
- Schenopus*
- morphological analysis of 387
 - relationship with Loliinae 382
- Schizachne*
- matK* analysis of 342
 - ndhF* analysis of 342
 - ndhH* analysis of 342
 - in Poaceae phylogeny 342
 - rbcL* analysis of 342
- Schizachyrium*
- in Andropogoneae phylogeny 530, 532, 535–542
 - biogeography of 514–515
 - ITS analysis of 535–536, 538, 541
 - ITS and *trnL*-F analysis of, combined 537, 540–542
 - morphological analysis of 541
 - New World 504–505, 514–515
 - in Panicoideae phylogeny 504–505, 514–515
 - photosynthetic pathways in 514–515
 - PHYB* analysis of 541
 - trnL*-F analysis of 535–536, 539, 541
- Schizachyrium scoparium*
- as climax species 273
 - ecological history of 281–284
 - light experiment with 274–278
 - light-nutrient experiment with 274–276, 279–283
 - New Jersey versus Oklahoma populations of 274–284

- nutrient experiment with 274–281
 reproductive plasticity of 273–285
 restoration use of 273–274, 284
 as successional species 273
- Schizostachyum*
- in Bambuseae phylogeny 326–328
 - morphological analysis of 324, 326
 - rpl16* analysis of 327
 - rpl16* and morphological analysis of, combined 328
- Schmidtia*, in Chloridoideae phylogeny 583
- Schoeneae
- Arthrostylis* placement in 81
 - clades of 77
 - in Cyperaceae phylogeny 73, 77–79, 204
 - 5.8S rDNA insertion in 88, 94
 - geographic distribution of 77
 - polytomy in 73
 - rbcL* analysis of 73, 77–79
 - relationship with Arthrostylideae 160
- Schoenefeldia*, in Chloridoideae phylogeny 580
- Schoenocephalum* 7
- Schoenoplectiella* 81
- Schoenoplectus*
- absence of 5.8S rDNA insertion in 88, 92, 94, 96
 - in Cyperaceae phylogeny 72, 79, 81
 - in Cyperales phylogeny 68
 - genome size of 66
 - ITS analysis of 156
 - in New Zealand 418
 - photosynthesis in, conflicting data on 104, 148
 - photosynthetic pathway in 103–104, 142–143
 - rbcL* analysis of 79, 81
 - rbcL* and *trnL* analysis of, combined 81
 - relationship with *Actinoscirpus* 81
 - relationship with *Isolepis humillima* 81
 - trnL*–F analysis of 157–158
 - trnL*–F and ITS analysis of, combined 159
- Schoenoxiphium*
- in Cariceae phylogeny 175–179, 183, 188–189
 - ITS, ETS-1f, and *trnL*–F analysis of, combined 166, 175–178
 - ITS analysis of 179
 - photosynthetic pathway in 103
- Schoenus*
- in Cyperaceae phylogeny 78, 204
 - 5.8S rDNA insertion in 88, 92, 94
 - geographic distribution of 204
 - inflorescence unit of (spikelet) 204–209
 - photosynthetic pathway in 103, 143–145
 - rbcL* analysis of 78
- Schulte, Katharina 27–43
- Scirpeae
- in Cyperaceae phylogeny 73, 77–82
 - 5.8S rDNA insertion in, absence/presence of 88, 94, 96
 - polyphyly of 73, 78
 - rbcL* analysis of 73, 77–81
 - relationship with Abildgaardieae 150
 - relationship with Cariceae 77–81
- Scirpodendron*
- in Cyperaceae phylogeny 73, 78
 - 5.8S rDNA insertion in 88, 92, 94
 - in Juncaceae phylogeny 58
 - photosynthetic pathway in 103, 145
 - rbcL* analysis of 73, 78
 - rps16* analysis of 57–58
 - trnL*–F analysis of 57–58
- Scirpoides*
- absence of 5.8S rDNA insertion in 87, 94
 - in Cyperaceae phylogeny 58, 73, 80–81
 - photosynthetic pathway in 103, 145
 - rbcL* analysis of 73, 80–81
 - rps16* analysis of 57–58
 - trnL*–F analysis of 57–58
- Scirpus*
- in Cyperaceae phylogeny 79, 84
 - in Cyperales phylogeny 68
 - 5.8S rDNA insertion in 88, 94, 96
 - economic uses of 84
 - genome size of 66
 - as outgroup in Cariceae study 172–173, 177–178
 - photosynthetic pathway in 103, 145–146
 - rbcL* analysis of 79
- Scleria*
- in Cyperaceae phylogeny 78
 - 5.8S rDNA insertion in 88, 94
 - genome size of 66
 - photosynthesis in, conflicting data on 148
 - photosynthetic pathway in 103, 105
 - rbcL* analysis of 78
- Sclerieae
- in Cyperaceae phylogeny 73, 77–78
 - 5.8S rDNA insertion in 88, 94
 - monophyly of 73
 - rbcL* analysis of 73, 77–78
 - relationship with Bisboeckelereae 77
- Sclerioideae
- in Cyperaceae phylogeny 73, 77–78, 82
 - 5.8S rDNA insertion in 88
 - rbcL* analysis of 73, 77–78
 - relationship with Schoeneae 77
- Sclerochloa*
- morphological analysis of 387, 394
 - morphological-molecular analysis of, combined 397
 - as outgroup in Loliinae study 382, 387, 394, 397
 - in Pooideae phylogeny 343, 346, 381
 - tribal/subtribal placement of 424
- Scleropoa*, in Pooideae phylogeny 380
- Scleropogon*
- biogeography of 588, 593
 - in Chloridoideae phylogeny 567, 570–571, 573, 581
 - ITS analysis of 567, 571
 - ITS and *trnL*–F analysis of, combined 573

- New World 581, 588, 593
trnL-F analysis of 567, 570
- Scolochloinae, in Poaceae phylogeny 421
- Scutachne*
 biogeography of 509–510, 518
 New World 505, 507, 509–510, 518
 in Panicoideae phylogeny 505, 507, 518
 photosynthetic pathway of 518
- Seberg, Ole and Gitte Petersen—Phylogeny of Triticeae (Poaceae) based on three organelle genes, two single-copy nuclear genes, and morphology 362–371
- Secale*
 chloroplast DNA analysis of 366, 368
DMCI analysis of 365, 367, 370
EF-G analysis of 367, 370
 mitochondrial genome of 366
 molecular analysis of 390–392
 morphological analysis of 365, 387, 394
 morphological-molecular analysis of, combined 397
 as outgroup in Loliinae study 382, 387, 390–392, 394, 397
 total evidence (combined) analysis of 367–369
 in Triticeae phylogeny 362, 365–370
- Sehima*, in Poaceae phylogeny 251, 254
- Self-fertilization. *See also* Cleistogamy
 in Poaceae 286
- Sequencia* 18
 in Bromeliaceae phylogeny 8–9, 16, 18
 CAM photosynthesis by 11
 character states of 18
 epiphytism of 12
 geographic origin and distribution of 10
 leaf and trichome anatomy of 14, 18
 morphology of 18
ndhF analysis of 8–9, 16, 18
 pollination of 13
- Sequencia serrata* 18
 in Bromeliaceae phylogeny 7–9, 14–15, 17–18
ndhF analysis of 7–9, 14–15, 17–18
- Sesleria*
 geographic distribution of 380
 molecular analysis of 390–392
 morphological analysis of 387, 394
 morphological-molecular analysis of, combined 397
 as outgroup in Loliinae study 382, 387, 390–392, 394, 397
 in Poaceae phylogeny 380
 in Pooideae phylogeny 343, 346, 380
 relationship with *Poa* 421–422, 424
- Seslerieae
 as outgroup in Loliinae study 382
 in Pooideae phylogeny 380
- Sesleriinae
 in Poaceae phylogeny 421
 in Pooideae phylogeny 346
 relationship with *Poa* 421–422
- Setaria*
 biogeography of 509, 528–529
 economic and ecological importance of 545
 New World 504–509, 528–529
 in Panicoideae phylogeny 504–505, 507–508, 528–529
 photosynthetic pathway of 511, 528–529
- Setariinae
 biogeography of 510–511, 528–529
 New World 504–508, 510–511, 528–529
 in Panicoideae phylogeny 504–505, 507–508, 528–529
 photosynthetic pathway of 508, 528–529
- Setariopsis*
 biogeography of 529
 New World 505, 508, 529
 in Panicoideae phylogeny 505, 508, 529
 photosynthetic pathway of 529
- Sheath fertilization, in Poaceae 286
- Shibataeinae
 in Bambuseae phylogeny 306–307, 315–317, 326–328
 morphological analysis of 326
rpl16 analysis of 327
rpl16 and morphological analysis of, combined 328
- Siberia, *Poa* variability in 435–449
- Simon, Bryan K.—Grass phylogeny and classification: conflict of morphology and molecules 259–266
- Simpson, David A. 55–61
- Simpson, David A. et al.—Phylogeny of Cyperaceae based on DNA sequence data—a new *rbcL* analysis 72–83
- Sinochasea*
matK analysis of 342
ndhF analysis of 342
ndhH analysis of 342
 in Poaceae phylogeny 342
 in Pooideae phylogeny 345–346
rbcL analysis of 342
 relationship with *Bromus* 450
- Siqueiros-Delgado, Maria Elena 565–579
- Sirochloa*
 in Bambuseae phylogeny 315, 326–328, 330
 morphological analysis of 324, 326, 330
rpl16 analysis of 327
rpl16 and morphological analysis of, combined 328
- Skendzic, Elizabeth M. et al.—Phylogenetics of Andropogoneae (Poaceae: Panicoideae) based on nuclear ribosomal internal transcribed spacer and chloroplast *trnL-F* sequences 530–544
- Smets, Erik 72–83, 204–209
- Snowdenia*, in Panicoideae phylogeny 503
- Sohnsia*
 biogeography of 587, 593
 breeding system of 608
 in Chloridoideae phylogeny 567, 570–571, 573, 581
 ITS analysis of 567, 571
 ITS and *trnL-F* analysis of, combined 573

- New World 581, 587, 593
trnL-F analysis of 567, 570
- Soreng, Robert J. 335–348, 420–434
- Sorghastrum*
 in Andropogoneae phylogeny 530, 536, 538–540, 542–543
 biogeography of 515
 ITS analysis of 536, 538, 543
 ITS and *trnL*-F analysis of, combined 540, 543
 New World 505, 515
 in Panicoideae phylogeny 505, 515
 photosynthetic pathway of 515
trnL-F analysis of 536, 539, 543
- Sorghinae
 in Andropogoneae phylogeny 262, 532, 538–540, 542
 biogeography of 515
 ITS analysis of 538
 ITS and *trnL*-F analysis of, combined 540
 New World 504–506, 515
 in Panicoideae phylogeny 503–505, 515
 photosynthetic pathways in 515
trnL-F analysis of 539
- Sorghum*
 in Andropogoneae phylogeny 262, 530, 532, 536–540, 542
 biogeography of 515
 in Centothecoideae + Panicoideae phylogeny 494–496
 economic and ecological importance of 530, 545
 ITS analysis of 536, 538, 542
 ITS and *trnL*-F analysis of, combined 537, 540
 morphological analysis of 494
ndhF analysis of 494–496, 542
 New World 505, 515
 in Panicoideae phylogeny 505, 515
 photosynthetic pathway of 515
 in Poaceae phylogeny 238, 242–245
rpl16 analysis of 494, 496
tb1 analysis of 542
trnL-F analysis of 536, 539
- Sorghum* subgen. *Eu-Sorghum* 542
- South America
 Bromeliaceae origin and distribution in 5, 7, 10–12, 18–19
 Chloridoideae in 585–590
 Panicoideae in 509–512
- Sparganiaceae, relationship with Bromeliaceae 8
- Sparganium*, molecular clock analysis of 5
- Spartina*
 biogeography of 589, 594
 in Chloridoideae phylogeny 567, 570–571, 573–576, 580–581, 583–584
 ITS analysis of 567, 571, 574, 576
 ITS and *trnL*-F analysis of, combined 573
 morphological analysis of 575, 583–584
ndhF and *rbcL* analysis of, combined 574
 New World 581, 583–584, 589, 594
- in Poaceae phylogeny 242–244
 relationship with *Eragrostis* 600
trnL-F analysis of 567, 570
- Spartochloa*
 in Centothecoideae + Panicoideae phylogeny 489, 494–496, 498–499
 morphological analysis of 494
ndhF analysis of 494–496
 in Poaceae phylogeny 254
rpl16 analysis of 494, 496
- Spathanthus* 7
- Speciation. *See also specific taxa*
 chromosome evolution and 62
- Sphaerocyperus*
 absence of 5.8 rDNA insertion in 87, 94
 in Cyperaceae phylogeny 80
 photosynthetic pathway in 103, 146
rbcL analysis of 80
- Sphenaria*
 biogeography of 509–511, 528
 New World 505, 508–511, 528
 in Panicoideae phylogeny 505, 508, 528
 photosynthetic pathway of 528
- Sphenopus*
 molecular analysis of 390–392
 morphological analysis of 387, 394
 morphological-molecular analysis of, combined 397
 in Poeae phylogeny 380, 403
 in Pooideae phylogeny 343, 346
 relationship with Loliinae 380, 387, 394, 397, 401
 tribal/subtribal placement of 424
- Spikelet(s)
 of *Aegilops triuncialis* 296
 of Andropogoneae 261–262, 504, 530–531, 543
 of *Anthoxanthum odoratum* versus *A. alpinum* 474–477
 of Arthropogoninae 507
 of *Arundinella* 262
 of Arundinelleae 505
 of Bambuseae 320–321, 325
 of *Bouteloua* 606–613
 of *Bromus* 451, 468
 of Centothecoideae 506
 of Centothecoideae + Panicoideae clade 488–489, 491–492
 of Chloridoideae 565, 583–585
 of *Eragrostis* 595–596
 of Isachneae 506
 of *Microstegium* 288
 of Olyreae 310, 315
 of Paniceae 507–508
 of Panicoideae 503
 of Parianinae 310
 of Paspalinae 508
 of *Paspalum* 545, 552–555
 of *Poa* 438–440
 of Poaceae 259, 261, 286

- of Pooideae 380
- of *Schoenus* 204–209
- of Steyermarkochloae 509
- of *Thrasya* 546, 552–555
- of *Thrasypopsis* 546, 552
- of Triodieae 262–263
- of *Zeugites* clade 498–499
- Spikelet clade, of Poaceae 252, 255
- Spinifex grasses 262–263. *See also* Triodieae
- Sporobolinae
 - biogeography of 586, 594
 - in Chloridoideae phylogeny 575–576, 581–584
 - morphological analysis of 583–584
 - New World 581–584, 586, 594
- Sporobolus*
 - biogeography of 585, 587–588, 594
 - in Chloridoideae phylogeny 567, 570–576, 581, 583–584
 - ITS analysis of 567, 571, 574
 - ITS and *trnL*–F analysis of, combined 573
 - matK* analysis of 342
 - morphological analysis of 494, 575–576, 583–584
 - ndhF* analysis of 342, 494–496, 574
 - ndhH* analysis of 341–342
 - New World 581, 583–585, 587–588, 594
 - in Poaceae phylogeny 341–342
 - rbcL* analysis of 342
 - relationship with *Thellungia* 599–600
 - reproductive plasticity in 282–284
 - rpl16* analysis of 494, 496, 498
 - trnL*–F analysis of 567, 570
- SPOSTA clade, of *Poa* 421, 428–430, 432
- Stace, Clive A. 380–405
- Starr, Julian R. 72–83, 165–192
- Starr, Julian R. et al.—Phylogenetic implications of a unique 5.8S rDNA insertion in Cyperaceae 84–98
- Stegolepis* 7
- Steinchisma*
 - biogeography of 509, 528
 - New World 505, 508–509, 528
 - in Panicoideae phylogeny 505, 508, 528
 - photosynthetic pathways in 504, 528
 - relationship with *Paspalum* and *Thrasya* 547, 549–551
- Steirachne*
 - biogeography of 587, 594
 - in Chloridoideae phylogeny 565–567, 570–571, 573, 581, 583
 - morphological analysis of 583
 - New World 565–567, 581, 583, 587, 594
- Stenotaphrum*
 - New World 505, 507–508
 - in Panicoideae phylogeny 505, 507–508
- Stephanachne*, in Pooideae phylogeny 346
- Stevenson, Dennis Wm. 227–233
- Stewart, Alan V. 406–419
- Steyerbromelia*
 - in Bromeliaceae phylogeny 4, 16–17, 39–40
 - ndhF* analysis of 4, 16–17
- Steyermarkochloa*
 - biogeography of 509–510, 529
 - in Centothecoideae + Panicoideae phylogeny 497
 - New World 505, 508–510, 529
 - in Panicoideae phylogeny 505, 508–509, 529
 - photosynthetic pathway of 529
- Steyermarkochloae
 - in Arundinoideae phylogeny 503, 508
 - biogeography of 510, 529
 - in Centothecoideae + Panicoideae phylogeny 488, 491–500
 - ndhF* analysis of 490–500
 - New World 504–506, 508–510, 529
 - in Panicoideae phylogeny 503–505, 508–509, 529
 - photosynthetic pathway of 529
 - rpl16* analysis of 490–500
 - spikelets of 509
- Stiburus*
 - in Chloridoideae phylogeny 576
 - relationship with *Eragrostis* 595–600, 602
- Stipa*
 - geographic distribution of 360
 - ITS analysis of 351, 355–357, 360
 - matK* analysis of 342
 - ndhF* analysis of 342
 - ndhH* analysis of 342
 - in Poaceae phylogeny 242–244, 256, 342
 - rbcL* analysis of 342
 - in Stipeae phylogeny 349–351, 355–357, 360
 - uncertain circumscription of 349
- Stipa* s.s. 350, 360
- Stipa* sect. *Lasiagrostis* 350
- Stipa* subgen. *Jarava* 350
- Stipa* subgen. *Pappostipa* 354, 360
- Stipagrostis*
 - matK* analysis of 342
 - morphological analysis of 494
 - ndhF* analysis of 342, 494–496
 - ndhH* analysis of 342
 - in Poaceae phylogeny 242–244, 342
 - rbcL* analysis of 342
 - rpl16* analysis of 494, 496
- Stipeae
 - core genera of 351
 - early diverging 360
 - Eurasian origin of 360
 - fluid circumscription of 349
 - ITS analysis of 349–360
 - morphological analysis of 349
 - phylogeny of 349–361
 - in Poaceae phylogeny 239, 252–253
 - in Pooideae phylogeny 345–346

- Streptocalyx*
 in Bromeliaceae phylogeny 27
trnL and *trnL-F* analysis of, combined 36
- Streptochoaeta*
matK analysis of 342
ndhF analysis of 241–242, 248–249, 341–342, 344
ndhH analysis of 341–342
 in Poaceae phylogeny 239–244, 248–249, 251, 254, 337, 341–342, 344
rbcL analysis of 342
- Streptochoaeteae
 in Anomochloideae phylogeny 303
 in Poaceae phylogeny 252–253
- Streptogyna*
 morphological analysis of 324, 494
ndhF analysis of 494–496
 as outgroup in Bambuseae study 319, 322, 326–328
 in Poaceae phylogeny 237–239, 242–246, 260
rpl16 analysis of 494, 496
- Streptogynaeae, in Poaceae phylogeny 252–253
- Streptogynaeae
 biogeography of 306
 in Centothecoideae + Panicoideae phylogeny 499
 in Ehrhartoideae phylogeny 303, 306
 morphology of 306
 in Poaceae phylogeny 245
- Streptostachys*
 biogeography of 509, 511, 528
 New World 505, 508–509, 511, 528
 in Panicoideae phylogeny 505, 508, 528
 photosynthetic pathway of 528
- Sucrea*
 in Bambuseae phylogeny 326–328
 biogeography of 310
 morphological analysis of 324, 326
 in Olyrinae (Olyreae) phylogeny 311
rpl16 analysis of 322, 327
rpl16 and morphological analysis of, combined 328
 spikelets and glumes of 311
- Sumatrosclirpus*, photosynthetic pathway in 103, 146
- Swallenia*
 biogeography of 587, 593
 in Chloridoideae phylogeny 567, 570–571, 573, 581
 ITS analysis of 567, 571
 ITS and *trnL-F* analysis of, combined 573
 New World 581, 587, 593
trnL-F analysis of 567, 570
- Symplectrodia*
 “hard” anatomy of 263
 in Triodieae phylogeny 262–263
- Synapomorphy
 of Chloridoideae 580
 of Panicoideae 261, 503
 of Poaceae 260–261
- Synonym database, grasses 267–269
- Syntrinema*, inclusion in *Rhynchospora* 104
- Sytsma, Kenneth J. 3–26
- Taeniatherum*
 chloroplast DNA analysis of 366, 368
DMC1 analysis of 367, 370
EF-G analysis of 367, 370
 mitochondrial genome of 366
 morphological analysis of 365–366
 relationship with *Elymus* 374, 377
 total evidence (combined) analysis of 367–369
 in Triticeae phylogeny 365–370
- Tanzania, Chloridoideae in 585
- Tatianyx*
 biogeography of 509, 511, 528
 New World 505, 508–509, 511, 528
 in Panicoideae phylogeny 505, 508, 528
 photosynthetic pathway of 528
- Taxonomic Database Working Group (TDWG) 267
- Taxonomy (grasses)
 DELTA system for 267–268
 electronic world database for 267–270
 future developments in 269
- tb1*, in Andropogoneae analysis 531, 541–543
- Telomere(s), of Cyperales 62–64
- Temburongia*
 in Bambuseae phylogeny 323, 326–328
 morphological analysis of 324, 326
rpl16 analysis of 327
rpl16 and morphological analysis of, combined 328
- Temochloa*
 in Bambuseae phylogeny 315, 317, 323, 326–329
 morphological analysis of 317, 324, 326, 329
rpl16 analysis of 327
rpl16 and morphological analysis of, combined 328–329
- Tetrachne*
 in Chloridoideae phylogeny 576, 581, 583
 morphological analysis of 583
 New World 581, 583
- Tetralia*
 photosynthetic pathway in 103, 146
 in Schoeneae phylogeny 77–78
- Thamnocalamus*, in Arundinariinae (Bambuseae) phylogeny 307
- Thamnochortus*, relationship with *Rhodocoma* 213, 218, 222
- Thellungia*
 in Chloridoideae phylogeny 574, 576
 relationship with *Eragrostis* 595–596, 599–600, 602
 relationship with *Sporobolus* 599–600
- Themeda*
 in Andropogoneae phylogeny 530, 532, 536–540
 ITS analysis of 536, 538
 ITS and *trnL-F* analysis of, combined 537, 540, 542
 New World 504–505
 in Panicoideae phylogeny 504–505
trnL-F analysis of 536, 539

Thinopyrum

- chloroplast DNA analysis of 366, 368
- DMCI* analysis of 367
- EF-G* analysis of 367
- genome transfer to *Elymus* 372, 374–378
- mitochondrial genome of 366
- morphological analysis of 365
- total evidence (combined) analysis of 367–369
- in Triticeae phylogeny 362, 365–369

Thomas, William W. 72–83

Thrasya

- axillary penduncles of 553
- biogeography of 509, 528
- cladistic analysis of 547–552
- inflorescence of 546, 552, 555–556
- leaf anatomy of 546, 552, 555
- lemmas of 546, 553–554
- lower glumes of 546, 552–554
- lower palea of 546, 553
- morphological analysis of 545–562
- ndhF* analysis of 546–547
- New World 505, 508–509, 528
- in Panicoideae phylogeny 505, 508, 528
- papillae of 553, 555
- pedicels of 546, 555
- photosynthetic pathway of 528
- racemes of 552–553, 556
- rachis of 546, 555–556
- relationship with *Panicum* 546
- relationship with *Paspalum* 545–556
- rpoC2* analysis of 547
- spikelets of 546, 552–555
- upper antherium of 546, 555
- upper glumes of 553–554

Thrasypopsis

- biogeography of 509, 511, 528
- inflorescence of 546, 552, 556
- lower glumes of 546, 552–553
- lower palea of 546, 553
- morphological analysis of 546
- New World 505, 508–509, 511, 528
- in Paniceae phylogeny 556
- in Panicoideae phylogeny 505, 508, 528
- photosynthetic pathway of 528
- racemes of 546, 552, 556
- rachis of 546, 556
- relationship with *Panicum* 546
- relationship with *Paspalum* 545–547, 549–556
- spikelets of 546, 552
- upper glumes of 546

Thuarea, in Poaceae phylogeny 254

Thurnia

- geographic distribution of 55
- habitat of 55
- in Juncaceae phylogeny 56
- relationship with Cyperaceae 56, 58, 60

- relationship with Juncaceae 55–56, 58, 60
- rps16* analysis of 57–58
- in Thurniaceae phylogeny 55–56, 58
- trnL-F* analysis of 57–58

Thurniaceae

- absence of 5.8S rDNA insertion in 94
- in Cyperales phylogeny 55
- flowers of 55–56
- phylogeny of 55–56, 58, 60
- in Poales phylogeny 55
- relationship with Bromeliaceae 27–28
- relationship with Juncaceae 55–56, 60
- rps16* analysis of 58
- trnL-F* analysis of 58

Thysanolaena

- in Centothecoideae + Panicoideae phylogeny 488–489, 494–497
- geographic distribution of 489
- morphological analysis of 494
- ndhF* analysis of 494–496, 498
- New World 505, 509
- in Panicoideae phylogeny 504–505, 509
- in Poaceae phylogeny 241–244, 251, 254
- relationship with *Steyermarkochloa* 508
- rpl16* analysis of 494, 496
- spikelets of 489

Thysanolaena-Centotheca-Cyperochloa clade 497–498

Thysanolaeneae

- base chromosome number of 509
- in Centothecoideae + Panicoideae phylogeny 488–489, 491–500
- ndhF* analysis of 490–500
- New World 504–506, 509
- in Panicoideae phylogeny 504–505, 509
- in Poaceae phylogeny 252–253
- rpl16* analysis of 490–500
- Thysanolaena* placement in 509

Tillandsia

- in Bromeliaceae phylogeny 7–9, 28, 34–36, 41
- CAM photosynthesis by 11
- chlorophyll fluorescence of 45
- epiphytism of 12, 50
- geographic origin and distribution of 5, 10–11, 18
- growth form of 22
- leaf traits of 14
- ndhF* analysis of 7–9
- pollination of 13, 22
- trichome(s) of 14, 44–52
- trichome photoprotection of 44–51
- trichome reflectance in 44–45, 48–51
- trichome structure and density in 45–48
- trichome wing attenuation in 50–51
- trichome wing facets in 50–51
- trnL*, *trnL-F*, and *trnT-F* analysis of, combined 34
- trnL* and *trnL-F* analysis of, combined 36

- water vapor condensation on leaf surfaces of 48, 50–51
- Tillandsia* subgen. *Pseudalcantarea* 13
- Tillandsioideae
- atmospherics 20–21, 44
 - in Bromeliaceae phylogeny 3–4, 7–9, 14–16, 28, 34, 40, 44
 - CAM photosynthesis by 11, 19–20
 - chlorophyll fluorescence of 45
 - epiphytism of 12, 19–21, 50
 - geographic origin and distribution of 10–11, 18–19
 - growth form of 22
 - leaf traits of 14, 19–21
 - matK-rps16* analysis of, combined 28
 - ndhF* analysis of 3–4, 7–9, 14–16
 - as outgroup in Bromelioideae study 33, 35
 - pitcairnioid ancestor of 18–19
 - plumose seeds of 3
 - pollination of 12–13, 22
 - trichome function in 44
 - trichome photoprotection of 44–51
 - trichome reflectance in 44–45, 48–51
 - trichome structure and density in 45–48
 - trichome wing attenuation in 50
 - trichome wing facets in 50–51
 - trichome(s) of 14, 19–21, 44–52
 - trnL* analysis of 28
 - trnL* and *trnL-F* analysis of, combined 35
 - water vapor condensation on leaf surfaces of 48, 50
- Timouria*
- matK* analysis of 342
 - ndhF* analysis of 342
 - ndhH* analysis of 342
 - in Poaceae phylogeny 342
 - rbcL* analysis of 342
- Torrecilla, Pedro 380–405
- Torres, Amelia 349–361
- Torreyochloa*, in Pooideae phylogeny 343, 346
- Tovarochloa*, tribal/subtribal placement of 424
- Trachyeae, in Panicoideae phylogeny 503
- Trachypogon*
- in Andropogoneae phylogeny 530, 532, 536, 538–540
 - biogeography of 515
 - ITS analysis of 536, 538
 - ITS and *trnL-F* analysis of, combined 540
 - New World 505, 515
 - in Panicoideae phylogeny 505, 515
 - photosynthetic pathway of 515
 - spikelets of 530
 - trnL-F* analysis of 536, 539
- Trachys*, in Panicoideae phylogeny 503
- Trachystylis*
- in Cyperaceae phylogeny 81, 149, 160–161
 - geographic distribution of 149
 - photosynthetic pathway in 103, 146
- Traginae** 592
- biogeography of 588, 593
 - in Chloridoideae phylogeny 581–582, 585
 - morphological analysis of 585, 592
 - New World 581–582, 585, 588, 593
- Tragus*
- breeding system of 608–610
 - in Chloridoideae phylogeny 567–568, 570–571, 573, 577, 581, 585
 - ITS analysis of 567–568, 571
 - ITS and *trnL-F* analysis of, combined 573
 - morphological analysis of 577, 585
 - New World 581, 585
 - trnL-F* analysis of 567, 570
- Tremburongia*, in Bambuseae phylogeny 315
- Trianoptiles*, photosynthetic pathway in 103, 146
- Tribolium*, in Danthonioideae phylogeny 261
- Tribuloideae 194
- Trichloris*
- biogeography of 588, 593
 - in Chloridoideae phylogeny 567, 570–571, 573, 577, 581, 584
 - ITS analysis of 567, 571
 - ITS and *trnL-F* analysis of, combined 573
 - morphological analysis of 584
 - New World 581, 584, 588, 593
 - trnL-F* analysis of 567, 570
- Tricholaena*
- in Panicoideae phylogeny 503
 - in Poaceae phylogeny 254
- Trichome(s)
- of Bromeliaceae 5, 13–14, 19–22, 27, 44–52
 - function of 44
 - photoprotection by 44
 - reflectance by 44–45, 48–51
 - of *Tillandsia* 14, 44–52
 - wet versus dry 44–45
- Trichoneura*
- biogeography of 593
 - in Chloridoideae phylogeny 567, 570–571, 573, 581
 - ITS analysis of 567, 571
 - ITS and *trnL-F* analysis of, combined 573
 - New World 581, 593
 - trnL-F* analysis of 567, 570
- Trichophorum*
- in Cyperaceae phylogeny 79, 81
 - 5.8S rDNA insertion in, absence/presence of 88, 92, 94, 96
 - northern temperate versus Andean 81
 - as outgroup in Cariceae study 172–173, 177–178
 - photosynthetic pathway in 103, 146
 - rbcL* analysis of 79, 81
- Trichoschoenus*
- in Cyperaceae phylogeny 81, 149, 160–161
 - geographic distribution of 149
 - photosynthetic pathway in 103, 146

- Tricostularia*
 in Cyperaceae phylogeny 78
 photosynthetic pathway in 103, 146
rbcL analysis of 78
- Tridens*
 biogeography of 587, 593
 in Chloridoideae phylogeny 567, 570–571, 573, 575–577, 581
 ITS analysis of 567, 571, 577
 ITS and *trnL*-F analysis of, combined 573, 575–576
 morphological analysis of 576
 New World 581, 587, 593
trnL-F analysis of 567, 570, 577
- Trikeriaia*
matK analysis of 342
ndhF analysis of 342
ndhH analysis of 342
 in Poaceae phylogeny 342
rbcL analysis of 342
- Trilepideae
 in Cyperaceae phylogeny 73, 78
 5.8S rDNA insertion in 94
 monophyly of 73
rbcL analysis of 73, 78
- Trilepis*
 in Cyperaceae phylogeny 78
 photosynthetic pathway in 103, 146
rbcL analysis of 78
- Trimonoecy 606
- Triodia*
 in Chloridoideae phylogeny 567, 570–571, 573
 ITS analysis of 567, 571
 ITS and *trnL*-F analysis of, combined 573
 spikelets of 262
 in Triodieae phylogeny 262–263
trnL-F analysis of 567, 570
- Triodieae 262–263
 in Chloridoideae phylogeny 575
 ITS analysis of 263
 leaf anatomy of 263
 phylogeny of 262–263
 “soft” and “hard” types of 263
 spikelets of 262–263
- Triodiinae, in Chloridoideae phylogeny 574–575
- Triplachne*, in Pooideae phylogeny 343, 346
- Triplasis*
 biogeography of 587, 593
 in Chloridoideae phylogeny 567, 570–571, 573, 577, 580–581
 cleistogamy and chasmogamy in 287–293
 ITS analysis of 567, 571
 ITS and *trnL*-F analysis of, combined 573
 New World 581, 587, 593
trnL-F analysis of 567, 570
- Triplett, Jimmy 315–332
- Tripogon*
 biogeography of 593
 in Chloridoideae phylogeny 567, 570–571, 573, 581
 ITS analysis of 567, 571
 ITS and *trnL*-F analysis of, combined 573
 New World 581, 593
trnL-F analysis of 567, 570
- Tripsacinae
 in Andropogoneae phylogeny 262, 531–533, 538–540, 543
 biogeography of 510, 516
 ITS analysis of 538, 543
 ITS and *trnL*-F analysis of, combined 540, 543
 morphological analysis of 543
 New World 504–506, 510, 516
 in Panicoideae phylogeny 504–505, 516
 photosynthetic pathways in 516
trnL-F analysis of 539
- Tripsacum*
 in Andropogoneae phylogeny 262, 530–533, 536–540, 543
 biogeography of 509–510, 516
 ITS analysis of 536, 538
 ITS and *trnL*-F analysis of, combined 537, 540
 New World 504–505, 509–510, 516
 in Panicoideae phylogeny 504–505, 516
 photosynthetic pathway of 516
 spikelets of 530
trnL-F analysis of 536, 539
- Triraphis*
 biogeography of 589, 593
 in Chloridoideae phylogeny 572, 581
 New World 581, 589, 593
- Triscenia*
 biogeography of 509–510
 New World 505, 508–510
 in Panicoideae phylogeny 505, 508
- Trisetum*
ndhF analysis of 341
 in Poaceae phylogeny 341
 in Pooideae phylogeny 343, 346
- Tristachya*
 biogeography of 516
 in Centothecoideae + Panicoideae phylogeny 489, 494–498
 ITS analysis of 533, 536, 538, 541
 morphological analysis of 494
ndhF analysis of 494–498
 New World 504–505, 516
 in Panicoideae phylogeny 504–505, 516
 photosynthetic pathway of 516
rbcL analysis of 497, 505, 533, 541
 relationship with Andropogoneae 533, 536–541
rpl16 analysis of 494, 496–498
- Tristeginae, *Thysanolaena* placement in 509

Triticeae

- allopolyploids of 372–379
- chloroplast DNA analysis of 363, 365–366, 368–369
- coxII* analysis of 363–366, 369
- DMC1* analysis of 363, 365–367, 369–370
- economic importance of 372
- EF-G* analysis of 363–367, 369–370
- GBSSI* analysis of 369
- mitochondrial genome of 363, 365–366, 369
- molecular analysis of 362–363
- monophyly of 362
- morphological analysis of 362–363, 365–368
- nuclear DNA analysis of 363, 365–367, 369–370
- as outgroup in Loliinae study 382
- phylogeny of 362–371
- plastid genome analysis of 363, 366
- in Poaceae phylogeny 252–253, 255–256
- in Pooideae phylogeny 345
- rbcL* analysis of 363–366
- relationship with Bromeeae 450
- reproductive plasticity in *Aegilops triuncialis* invasion 295–300
- reticulate history of 372
- rpoA* analysis of 363–366
- total evidence (combined) analysis of 362, 365, 367–369
- trnF* analysis of 368
- trnL* analysis of 368
- trnT* analysis of 368

Triticum

- BAC studies of 68
- chloroplast DNA analysis of 366, 368
- DMC1* analysis of 367
- EF-G* analysis of 367
- mitochondrial genome of 366
- morphological analysis of 365, 494
- ndhF* analysis of 336–340, 344–345, 494–496
- ndhH* analysis of 336–337, 344–345
- plastid genome analysis of 336–340
- in Poaceae phylogeny 242–244, 344–345
- in Pooideae phylogeny 343–345
- rpl16* analysis of 494, 496
- total evidence (combined) analysis of 367–369
- in Triticeae phylogeny 362, 365–369

- Triticum aestivum*, as outgroup in *Bromus* study 453, 455
- trnE–D*, in *Carex* subgen. *Vignea* analysis 166–174, 176, 180–181, 189

trnF

- in *Elymus* analysis 373–375
- in Triticeae analysis 368

- trnK*, in *Rhodocoma* analysis 213–214

trnL

- in Andropogoneae analysis 530–544
- in Bromeliaceae analysis 16, 28–29, 33–36
- in Bromelioideae analysis 29, 33–36
- in *Bromus* analysis 450–467

- in Chloridoideae analysis 565–579

- in *Elymus* analysis 373–375

- in Juncaceae analysis 55–61

- in Poaceae analysis 248, 252, 254

- in *Rhodocoma* analysis 213–214

- in Triticeae analysis 368

trnL–F

- in Abildgaardieae analysis 151–162

- in Andropogoneae analysis 262, 530–544

- in Bambuseae analysis 329–330

- in Bromeliaceae (Bromelioideae) analysis 29, 33–36

- in Chloridoideae analysis 565–579, 582

- in Chusqueinae analysis 316

- in Cyperaceae analysis 56–60, 81

- in *Festuca* analysis 381, 386, 389, 391, 397–398, 406–419

- in Juncaceae analysis 55–61

- in Loliinae analysis 382, 385–399

- in *Oxychloe* analysis 57, 59, 90

- in Poaceae analysis 254

- in *Rhodocoma* analysis 213–214, 217

- in Thurniaceae analysis 58

trnT

- in *Elymus* analysis 373–375

- in Triticeae analysis 368

trnT inversion, in Poaceae 237–238*trnT–F*

- in Bromeliaceae (Bromelioideae) analysis 29, 33–36

- in *Poa* analysis 420–434

trnT–L

- in Bromeliaceae (Bromelioideae) analysis 29, 33–35

- in Cariceae analysis 166

Trochiliade. *See* Hummingbird(s); Hummingbird pollination*Tuctoria*

- biogeography of 593

- in Chloridoideae phylogeny 567, 570–571, 573, 577, 581, 585

- ITS analysis of 567, 571

- ITS and *trnL–F* analysis of, combined 573

- morphological analysis of 585

- New World 581, 585, 593

- trnL–F* analysis of 567, 570

Tylocarya

- in Abildgaardieae phylogeny 149–151

- inclusion in *Fimbristylis* 150

Typha

- molecular clock analysis of 5

- as outgroup in Juncaceae/Cyperaceae study 57–58

- rps16* analysis of 57–58

- trnL–F* analysis of 57–58

Typhaceae, relationship with Bromeliaceae 4, 8

Tzvelevia, tribal/subtribal placement of 424*Uncinia*

- in Cariceae phylogeny 175, 177–179, 183

- in Cyperaceae phylogeny 79, 84

- 5.8S rDNA insertion in 86–87

- genome size of 66
 ITS, ETS-1f, and *trnL*-F analysis of, combined 166, 175, 177–178
 ITS analysis of 179
 photosynthetic pathway in 103, 146–147
rbcL analysis of 79
Uncinia subgen. *Eu-Uncinia* 172
Uncinia subgen. *Eu-Uncinia* sect. *Platyandrae* 172
Uniola
 biogeography of 587, 594
 in Chloridoideae phylogeny 567, 570–571, 573–574, 576, 581, 583
 ITS analysis of 567, 571
 ITS and *trnL*-F analysis of, combined 573
matK analysis of 342
 morphological analysis of 583
ndhF analysis of 342
ndhF and *rbcL* analysis of, combined 574
ndhH analysis of 342
 New World 581, 583, 587, 594
 in Poaceae phylogeny 242–244, 342
rbcL analysis of 342
 relationship with *Eragrostis* 597
trnL-F analysis of 567, 570
 Unioliinae
 biogeography of 594
 in Chloridoideae phylogeny 574–576, 581–583
 morphological analysis of 583
 New World 581–583, 594
 Unisexual flowers, evolution of 605–614
 United States
 Chloridoideae in 583, 585–588
Microstegium invasion in 288
 Panicoideae in 509–510, 512
 Uplands of Central Brazil Province, Panicoideae of 511–512
Urochloa
 biogeography of 510, 518–519
 New World 505, 507, 510, 518–519
 in Panicoideae phylogeny 505, 507, 518–519
 photosynthetic pathway of 511, 518–519
Ursulaea
 AFLP analysis of 39, 41
 in Bromeliaceae phylogeny 34, 36, 39, 41
trnL, *trnL*-F, and *trnT*-F analysis of, combined 34
trnL and *trnL*-F analysis of, combined 36
Vacoparis, in Andropogoneae phylogeny 262
Vahlodea, in Pooideae phylogeny 343, 346–347
Valiha
 in Bambuseae phylogeny 315, 326–328
 morphological analysis of 324, 326
rpl16 analysis of 327
rpl16 and morphological analysis of, combined 328
Vaseyochloa
 biogeography of 593
 in Chloridoideae phylogeny 567, 570–571, 573, 577, 581, 584
 ITS analysis of 567, 571
 ITS and *trnL*-F analysis of, combined 573
 morphological analysis of 584
 New World 581, 584, 593
trnL-F analysis of 567, 570
 Vázquez, Francisco 349–361
 Venezuela, Panicoideae in 509–510
Vetiveria
 in Andropogoneae phylogeny 530, 532, 536–542
 economic and ecological importance of 530
 ITS analysis of 536, 538, 541–542
 ITS and *trnL*-F analysis of, combined 537, 540–542
 morphological analysis of 542
trnL-F analysis of 536, 539, 541–542
Virescentia, in Stipeae phylogeny 350
Volkiella, photosynthetic pathway in 103, 147
Vriesea
 in Bromeliaceae phylogeny 7–9, 28, 41
 CAM photosynthesis by 11
 epiphytism of 12
 geographic origin and distribution of 5, 10–11, 18
 leaf traits of 14
ndhF analysis of 7–9
 pollination of 13, 22
 trichome photoprotection of 44–51
 trichome reflectance in 44–45, 48–51
 trichome structure and density in 45–48
 trichome(s) of 14, 44–52
 water vapor condensation on leaf surfaces of 48
Vriesea subgen. *Xiphion* 13
 Vrijdaghs, Alexander et al.—The *Schoenus* spikelet: a rhipidium? A floral ontogenetic answer 204–209
Vulpia
 in Loliinae phylogeny 380–405
 molecular analysis of 386, 389–392, 414–415
 morphological analysis of 382–385, 388–389, 394–395
 morphological-molecular analysis of, combined 397–398
 morphology of 381
 phylogeny of 381, 401, 403
 in Pooideae phylogeny 343, 346, 380–381
 relationship with *Festuca* 381, 406–407, 413
Vulpia sect. *Apalochloa* 381–382, 384, 386, 399, 401, 403
Vulpia sect. *Loretia* 381–382, 384, 386, 389, 401, 403
Vulpia sect. *Monachne* 381–382, 384, 386, 389, 401, 403
Vulpia sect. *Spirachne* 381, 401, 403
Vulpia sect. *Vulpia* 381–382, 384, 386, 389, 399–400, 403
Vulpiella, in Pooideae phylogeny 381
Wangenheimia
 in Loliinae phylogeny 380–405
 molecular analysis of 386, 389–392, 414–415
 morphological analysis of 389, 394–396
 morphological-molecular analysis of, combined 397
 relationship with *Festuca* 406–407, 413
 Waterway, Marcia J. and Julian R. Starr—Phylogenetic relationships in tribe Cariceae (Cyperaceae)

- based on nested analysis of four molecular data sets 165–192
- waxy* (*GBSSI*)
- in Bambuseae analysis 330
 - in Chloridoideae analysis 572–574, 576
 - in *Elymus* analysis 374, 376–378
 - in *Eragrostis* analysis 597–602
 - in Poaceae analysis 248–249, 252, 260, 337, 343–344
 - in Pooideae analysis 343
 - in Triticeae analysis 369
- Websteria*, photosynthetic pathway in 103, 147
- Weising, Kurt 27–43
- Werauhia*, in Bromeliaceae phylogeny 28
- West Indian Province, Panicoideae of 510–511
- Whitkus, Richard 62–71
- Williamson, Helen 267–270
- Willkommia*
- biogeography of 588, 593
 - in Chloridoideae phylogeny 567, 570–571, 573, 577, 581, 585
 - ITS analysis of 567, 571
 - ITS and *trnL*–F analysis of, combined 573
 - morphological analysis of 577, 585
 - New World 581, 585, 588, 593
 - trnL*–F analysis of 567, 570
- Wilson, Karen L. 72–83, 99–148, 149–164
- Wind pollination, of Cyperaceae 77
- Windsorina* 7
- Wittrockia*
- in Bromeliaceae phylogeny 29, 34, 36
 - trnL*, *trnL*–F, and *trnT*–F analysis of, combined 34
 - trnL* and *trnL*–F analysis of, combined 36
- Woody bamboos. *See* Bambuseae
- Xerochloa*, in Poaceae phylogeny 254
- Xyridaceae
- androecium of 230
 - gynoecium of 230–232
 - inflorescence of 227–233
 - monophyly of, debate over 232–233
 - perianth of 228–230
 - in Poales phylogeny 227
 - relationship with Bromeliaceae 27–28
 - relationship with Eriocaulaceae 227
- Xyris*
- androecium of 230
 - gynoecium of 231
 - inflorescence of 228–233
 - perianth of 228–230
- Zameioscirpus*
- in Cyperaceae phylogeny 72–73, 79, 81
 - photosynthetic pathway in 103, 147
 - rbcL* analysis of 73, 79, 81
- Zea*
- in Andropogoneae phylogeny 262, 530–533, 536–540, 543
 - biogeography of 509–510, 516
 - in Centothecoideae + Panicoideae phylogeny 494–496
 - economic and ecological importance of 530, 545
 - ITS analysis of 536, 538
 - ITS and *trnL*–F analysis of, combined 537, 540
 - matK* analysis of 342
 - morphological analysis of 494
 - ndhF* analysis of 336–337, 342, 344, 494–496
 - ndhH* analysis of 336–337, 342, 344
 - New World 504–505, 509–510, 516
 - in Panicoideae phylogeny 504–505, 516
 - photosynthetic pathway of 516
 - in Poaceae phylogeny 242–245, 342, 344
 - rbcL* analysis of 342
 - rpl16* analysis of 494, 496
 - spikelets of 530
 - trnL*–F analysis of 536, 539
- Zeugiteae, in Centothecoideae phylogeny 488
- Zeugites*
- biogeography of 509–510, 516
 - in Centothecoideae + Panicoideae phylogeny 489, 493–500
 - geographic distribution of 489
 - morphological analysis of 494, 498–499
 - ndhF* analysis of 493–496
 - New World 505–506, 509–510, 516
 - in Panicoideae phylogeny 505–506, 516
 - photosynthetic pathway of 516
 - in Poaceae phylogeny 242–244
 - rpl16* analysis of 493–494, 496, 498
 - spikelets of 489, 498–499
- Zeugites* clade 497–499
- Zhang, Xiufu 72–83
- Zingeria*, tribal/subtribal placement of 424
- Zizania*
- biogeography of 306
 - in Zizaniinae (Ehrhartoideae) phylogeny 306
- Zizaniinae
- biogeography of 306
 - in Oryzeae (Ehrhartoideae) phylogeny 306
- Zizaniopsis*, in Poaceae phylogeny 251, 254
- Zizka, Georg 27–43
- Zoysia*
- in Chloridoideae phylogeny 567, 570–571, 573–574, 576, 581, 584
 - ITS analysis of 567, 571, 576, 582
 - ITS and *trnL*–F analysis of, combined 573, 582
 - morphological analysis of 576, 584
 - ndhF* analysis of 574
 - ndhF* and *rbcL* analysis of, combined 574
 - New World 581, 584
 - in Poaceae phylogeny 242–244
 - trnL*–F analysis of 567, 570
- Zoysiaceae
- biogeography of 589, 594
 - in Chloridoideae phylogeny 576, 581–584
 - morphological analysis of 583–584

New World 581–584, 589, 594
Zoysiinae
biogeography of 590
in Chloridoideae phylogeny 576, 581–582, 584
morphological analysis of 584

New World 581–582, 584, 590
Zuloaga, Fernando O. 545–562
Zuloaga, Fernando O. et al.—Classification and biogeography of Panicoideae (Poaceae) in the New World 503–529